

FLOQUET MULTIPLIERS AND THE STABILITY OF PERIODIC LINEAR DIFFERENTIAL EQUATIONS: A UNIFIED ALGORITHM AND ITS COMPUTER REALIZATION

Mengda Wu¹, Yonghui Xia^{1,†} and Ziyi Xu¹

Abstract Floquet multipliers (characteristic multipliers) play significant role in the stability of the periodic equations. Based on the iterative method, we provide a unified algorithm to compute the Floquet multipliers (characteristic multipliers) and determine the stability of the periodic linear differential equations on time scales unifying discrete, continuous, and hybrid dynamics. Our approach is based on calculating the value of $\mathcal{A}$ and $\mathcal{B}$ (see Theorem 3.1), which are the sum and product of all Floquet multipliers (characteristic multipliers) of the system, respectively. We obtain an explicit expression of $\mathcal{A}$ (see Theorem 4.1) by the method of variation and approximation theory (iterative method), and an explicit expression of $\mathcal{B}$ by Liouville's formula. Furthermore, a computer program is designed to realize our algorithm. Specifically, you can determine the stability of a second order periodic linear system, whether they are discrete, continuous or hybrid, as long as you enter the program codes associated with the parameters of the equation. In fact, few literatures have dealt with the algorithm to compute the Floquet multipliers, not mention to design the program for its computer realization. Our algorithm gives the explicit expressions of all Floquet multipliers and our computer program is based on the approximations of these explicit expressions. In particular, on an arbitrary discrete periodic time scale, we can do a finite number of calculations to get the explicit value of Floquet multipliers (see Theorem 4.2). Therefore, for any discrete periodic system, we can accurately determine the stability of the system by our algorithm even without computer! Finally, in Section 6, several examples are presented to illustrate the effectiveness of our algorithm.

Keywords Time scales, periodic differential equations, stability, Floquet multipliers, Hill equations.

MSC(2010) 34N05, 34L16, 26E70, 34L15, 65L15.

[†]The corresponding author. Email: yhxia@zjnu.cn (Y. Xia)

¹School of Mathematical Science, Zhejiang Normal University, Jinhua, 321004, China

1. Introduction

1.1. Theory of time scales unifying the continuous and discrete calculus

In 1988, Hilger [31] introduced the theory of time scales for the propose of unifying discrete and continuous calculus ([32, 33]). The systematic works of dynamic equations on time scales, one can refer to Bohner and Peterson [12], Agarwal and Bohner [1], Agarwal et al [3], and Bohner et al. [11]. In particular, the theory of the exponential dichotomy, reducibility, linearization, Hyers-Ulam stability and Sturmian theory are well studied, one can refer to Pötzche [6, 44–46] and Siegmund [50], Doan et al. [23, 24, 47], Zhang et al [62, 63], Erbe and Peterson [26]. It was also generalized to the measure differential equations on time scales (Federson et al. [27, 28]), fuzzy-valued differential equations on time scales (Wang et al. [56–60]), quaternion-valued differential equations on time scales (Li et al. [39], Cheng et al. [16]) and Califford-valued differential equations (Li et al. [40]). Recently, DaCunha and Davis [21], DaCunha [20] extend the Floquet theory to a more general case of an arbitrary periodic time scale which unifies discrete, continuous, and hybrid periodic cases. Adivar and Koyuncuoğlu [4] constructs a unified Floquet theory for homogeneous and nonhomogeneous hybrid periodic systems on domains having continuous, discrete or hybrid structure using the new periodicity concept based on shifts.

1.2. Floquet theory and Floquet multipliers

Floquet theory indicates that a nonautonomous T -periodic linear system of differential equations can be reducible to a corresponding autonomous linear system of differential equations by a periodic Lyapunov transformation [17]. Floquet theory is a powerful tool to study the stability and periodic solutions of dynamic systems. Mathematicians have extended Floquet theory in different directions. We can classify the results of Floquet theory into some types: ODEs (almost Floquet systems [29], almost-periodic systems [34], periodic Euler-Bernoulli equations [43], delay differential equations [51], linear systems with meromorphic solutions [61]), PDEs (parabolic differential equations [18], periodic evolution problems [37]), DAEs [22, 38], integro-differential equations [2], Volterra equations [8], discrete dynamical systems (countable systems [53]) and systems on time scales [5]. More details for the Floquet theory and applications, one can also refer to the monograph [35] and the works [7, 30].

Floquet multipliers (characteristic multipliers) play great role in the Floquet theory and the stability of the periodic equations. Thus, usually, to determine the stability, it suffices to calculate the characteristic multipliers. More specifically, if all of the characteristic multipliers have modulus less than or equal to one, and if, for each characteristic multiplier with modulus equal to one, the algebraic multiplicity equals the geometric multiplicity, the system is stable, otherwise the system is unstable. Then a natural question is how to compute the characteristic multipliers of the periodic systems. To this end, mathematicians have proposed some methods to compute the characteristic multipliers of periodic differential equations. For examples, Kotsis [36] studied the approximation of the characteristic multipliers based on a theorem of Demidovič; Shi [49] estimated the periodic Hill equation; some very nice results were obtained for the delay differential equations (functional

differential equations), see Breda, Mast and Vermiglio [13], Chow and Walther [19], Val'ter and Skubachevskii [54], Skubachevskii and Walther [52], Walther [55], Luzyanina and Engelborghs [41], Dormayer et al. [25], Mallet-Paret and Sell [42]. Floquet multipliers also can be used to discuss the existence of periodic solution of dynamic system. For more study of periodic solutions of dynamic systems, one can see [9, 10].

1.3. Motivation, novelty and contributions

There are a few works considering the Floquet theory and characteristic multipliers as mentioned above. However, few existing literatures have dealt with the algorithm to compute the Floquet multipliers (characteristic multipliers), not mention to design the program for its computer realization. In this paper, we provide a unified algorithm to compute the Floquet multipliers (characteristic multipliers) and determine the stability of the second order periodic linear equations on periodic time scales. We claim that the periodic system is stable if

$$\left| \frac{\mathcal{A}}{2} + \sqrt{\left(\frac{\mathcal{A}}{2}\right)^2 - \mathcal{B}} \right| < 1 \quad \text{and} \quad \left| \frac{\mathcal{A}}{2} - \sqrt{\left(\frac{\mathcal{A}}{2}\right)^2 - \mathcal{B}} \right| < 1,$$

and system is unstable if

$$\left| \frac{\mathcal{A}}{2} + \sqrt{\left(\frac{\mathcal{A}}{2}\right)^2 - \mathcal{B}} \right| > 1 \quad \text{or} \quad \left| \frac{\mathcal{A}}{2} - \sqrt{\left(\frac{\mathcal{A}}{2}\right)^2 - \mathcal{B}} \right| > 1.$$

To determine the stability of the periodic system, it is sufficient to know the modulus of characteristic multipliers, which can be derived from $\mathcal{A}$ and $\mathcal{B}$. Our main task is to calculate the value of $\mathcal{A}$ and $\mathcal{B}$ (see Theorem 3.1), which are the sum and product of all characteristic multipliers of the system, respectively. We obtain an explicit expression of $\mathcal{A}$ (see Theorem 4.1) by the method of variation and approximation theory (iterative method) and an explicit expression of $\mathcal{B}$ by Liouville's formula. Finally, in Section 6, several examples are presented to illustrate the effectiveness of our algorithm. The illustrative examples show good performance of our computer program. We summarize the contributions of this paper as follows.

- (1) Based on the iterative method, we provide a unified algorithm to compute Floquet multipliers and determine the stability of the periodic linear differential equations on time scales unifying discrete, continuous, and hybrid dynamics.
- (2) A computer program is designed to realize our algorithm. Therefore, you can determine the stability of a second order periodic linear system, whether they are discrete, continuous or hybrid, as long as you enter the program codes associated with the parameters of the equation.
- (3) Few existing literatures have dealt with the algorithm to compute the Floquet multipliers, not mention to design the program for its computer realization. Our algorithm gives the explicit expressions of all Floquet multipliers and our computer program is based on the approximations of these explicit expressions.
- (4) We provide an estimate of the error between $\mathcal{A}(n)$ and $\mathcal{A}$. And a computer program is given for calculating the value of $\mathcal{A}(n)$, $\mathcal{B}$ and $\rho(n)$, where $\mathcal{A}(n)$ is the n -th approximation of $\mathcal{A}$ and $\rho(n)$ is the n -th approximations of modulus of characteristic multipliers.
- (5) In particular, on an arbitrary discrete periodic time scale, there is a constant $k \in \mathbb{N}$, such that $\mathcal{A} = \mathcal{A}(k)$. Consequently, we can do a finite number of calculations to get the explicit value of Floquet multipliers (see Theorem 4.2). Therefore,

for any discrete periodic system, we can accurately determine the stability of the system by our algorithm even without computer!

(6) We obtain an explicit expression of $\mathcal{A}$ (see Theorem 4.1) by the method of variation and approximation theory and an explicit expression of $\mathcal{B}$ by Liouville's formula.

1.4. Outline of the paper

The rest of this paper is organized as follows. In Section 2, we introduce some notations and lemmas. Section 3 gives the stability criteria for the systems we studied. Section 4 introduces the processes of getting the expression of $\mathcal{A}$. Our main results on the expression of $\mathcal{A}$ are collected in three theorems (Theorem 4.1–Theorem 4.3). In Section 5, a computer program is given. Finally, in Section 6, we give some examples to show the effectiveness of our algorithm and verify our computer program.

2. Preliminaries

For completeness, we recall the following notations and concepts for the theory of time scales from [12]. A time scale $\mathbb{T}$ is a nonempty closed subset of $\mathbb{R}$. We denote $[a, b] \cap \mathbb{T}$ by $[a, b]_{\mathbb{T}}$. The forward jump operator is defined by $\sigma(t) := \inf\{s \in \mathbb{T} : s > t\}$. The backward jump operator is defined by $\rho(t) := \sup\{s \in \mathbb{T} : s < t\}$. We put $\inf \emptyset = \sup \mathbb{T}$ and $\sup \emptyset = \inf \mathbb{T}$. A point $t \in \mathbb{T}$ is said to be right-dense if $\sigma(t) = t$, right-scattered if $\sigma(t) > t$, left-dense if $\rho(t) = t$, left-scattered if $\rho(t) < t$, isolated if $\rho(t) < t < \sigma(t)$, and dense if $\rho(t) = t = \sigma(t)$. A set $\mathbb{T}^\kappa$ is defined as $\mathbb{T}^\kappa = \mathbb{T} - \{m\}$ if $\mathbb{T}$ has a left-scattered maximum, $\mathbb{T}^\kappa = \mathbb{T}$ otherwise. A time scale $\mathbb{T}$ is said to be discrete if t is scattered for all $t \in \mathbb{T}$, and it is said to be continuous if t is dense for all $t \in \mathbb{T}$. A function $f : \mathbb{T} \rightarrow \mathbb{R}$ is called regulated provided its right-sided limits exist (finite) at all right-dense points in $\mathbb{T}$ and its left-sided limits exist (finite) at all left-dense points in $\mathbb{T}$. A function $f : \mathbb{T} \rightarrow \mathbb{R}$ is called rd-continuous provided it is continuous at right-dense points in $\mathbb{T}$ and its left-sided limits exist (finite) at left-dense points in $\mathbb{T}$. The set of rd-continuous functions $f : \mathbb{T} \rightarrow \mathbb{R}$ is denoted by $C_{rd}(\mathbb{T}, \mathbb{R})$. The graininess function μ is defined by $\mu(t) := \sigma(t) - t$.

We say that a function $p : \mathbb{T} \rightarrow \mathbb{R}$ is regressive provided $1 + \mu(t)p(t) \neq 0$ holds for all $t \in \mathbb{T}^\kappa$. The set of all regressive and rd-continuous functions $f : \mathbb{T} \rightarrow \mathbb{R}$ is denoted by $\mathcal{R}$. The delta derivative of a function $f : \mathbb{T} \rightarrow \mathbb{R}$ at a point $t \in \mathbb{T}^\kappa$, denoted by $f^\Delta(t)$, is defined by

$$f^\Delta(t) = \lim_{s \searrow \mu(t)} \frac{f(t+s) - f(t)}{s}.$$

For a function $f : \mathbb{T} \rightarrow \mathbb{R}$ we shall talk about the second derivative $f^{\Delta\Delta}$ provided f^Δ is differentiable on $(\mathbb{T}^\kappa)^\kappa$ with $f^{\Delta\Delta} = (f^\Delta)^\Delta : (\mathbb{T}^\kappa)^\kappa \rightarrow \mathbb{R}$. A continuous function $f : \mathbb{T} \rightarrow \mathbb{R}$ is called pre-differentiable with D , provided $D \subseteq \mathbb{T}^\kappa, \mathbb{T}^\kappa \setminus D$ is countable and contains no right-scattered elements of $\mathbb{T}$, and f is differentiable at each $t \in D$. A pre-differentiable function $F : \mathbb{T} \rightarrow \mathbb{R}$ is called the pre-antiderivative of f if $F^\Delta(t) = f(t)$ holds for all $t \in D$, where D is the region of differentiation. Then we define the Cauchy integral by

$$\int_r^s f(t) \Delta t = F(s) - F(r), \quad \text{for } r, s \in \mathbb{T},$$

where F is the pre-antiderivative of f .

If $p \in \mathcal{R}$, we define the exponential function by

$$e_p(t, s) = \exp \left(\int_s^t \lim_{s \searrow \mu(\tau)} \frac{\text{Log}(1 + p(\tau)s)}{s} \Delta\tau \right) \quad \text{for } s, t \in \mathbb{T}.$$

One can see that $e_p(t, s)$ is a solution of the equation $x^\Delta = p(t)x$. The delta derivative of a vector-valued (matrix-valued) function is given by taking the derivative of each components. The integral of a vector-valued (matrix-valued) function can be given in a similar manner. Let A be an $m \times n$ -matrix-valued function on $\mathbb{T}$. We say that A is rd-continuous on $\mathbb{T}$ if each entry of A is rd-continuous on $\mathbb{T}$, and the class of all such rd-continuous $m \times n$ -matrix-valued functions on $\mathbb{T}$ is denoted by $C_{rd} = C_{rd}(\mathbb{T}, \mathbb{R}^{m \times n})$. An $n \times n$ -matrix-valued function A on a time scale $\mathbb{T}$ is called regressive provided $I + \mu(t)A(t)$ is invertible for all $t \in \mathbb{T}^\kappa$, and the class of all such regressive and rd-continuous functions is denoted by $\mathcal{R}$.

Definition 2.1 ([12], p.92). If $p \in C_{rd}$ and $\mu p^2 \in \mathcal{R}$, then we define the trigonometric functions $\cos_p$ and $\sin_p$ by

$$\cos_p = \frac{e_{ip} + e_{-ip}}{2} \quad \text{and} \quad \sin_p = \frac{e_{ip} - e_{-ip}}{2i}.$$

For trigonometric functions on time scales, we have some formulas, which can be found in ([12], Exercise 3.27).

Definition 2.2 ([21]). Let $T \in (0, \infty)$. Then the time scale $\mathbb{T}$ is T -periodic if for all $t \in \mathbb{T}$,

1. $t \in \mathbb{T}$ implies $t + T \in \mathbb{T}$;
2. $\mu(t) = \mu(t + T)$.

Definition 2.3 ([21]). $A : \mathbb{T} \rightarrow \mathbb{R}^{n \times n}$ is T -periodic if $A(t) = A(t + T)$ for all $t \in \mathbb{T}$.

Consider the regressive time varying linear dynamic initial value problem

$$x^\Delta(t) = A(t)x(t), \quad x(t_0) = x_0, \quad (2.1)$$

where $A(t)$ is T -periodic for $t \in \mathbb{T}$ and the time scale $\mathbb{T}$ is also T -periodic.

Definition 2.4 ([21]). Let $x_0 \in \mathbb{R}^n$ be a nonzero vector and $\Psi(t)$ be any fundamental matrix for the system (2.1). The vector solution of the system with initial condition $x(t_0) = x_0$ is given by $\Phi_A(t, t_0)x_0$. The operator $M : \mathbb{R}^n \rightarrow \mathbb{R}^n$ given by $M(x_0) := \Phi_A(t_0 + T, t_0) = \Psi(t_0 + T)\Psi^{-1}(t_0)x_0$, is called a monodromy operator. The eigenvalues of the monodromy operator are called the Floquet (or characteristic) multipliers of the system (2.1).

Lemma 2.1 ([21], Corollary 7.10). Consider the p -periodic system (2.1).

1. If all the Floquet multipliers have modulus less than one, then the system (2.1) is exponentially stable.
2. If all of the Floquet multipliers have modulus less than or equal to one, and if, for each Floquet multiplier with modulus equal to one, the algebraic multiplicity equals the geometry multiplicity, then the system (2.1) is stable; otherwise the system (2.1) is unstable, growing at rates of generalized polynomials of t .

3. If at least one Floquet multiplier has modulus greater than one, then the system (2.1) is unstable.

Lemma 2.2 ([12], p.23). Every regulated function on a compact interval is bounded.

Lemma 2.3. Assume that D is a compact subset of $\mathbb{R}$ and $f_n \in C_{rd}(D, \mathbb{R})$ for each $n \in \mathbb{N}$. If $\{f_n\}$ uniformly converges to f on D , then f is rd-continuous and

$$\int_a^b f(t) \Delta t = \lim_{n \rightarrow \infty} \int_a^b f_n(t) \Delta t.$$

where $a, b \in D$.

Lemma 2.4. Let $\mathbb{T}$ be an arbitrary time scale. Suppose $f : [a, b] \rightarrow \mathbb{R}$ is an increasing function, where $a, b \in \mathbb{T}$ (b may be ∞). If f is rd-continuous when it is restricted on $[a, b]_{\mathbb{T}}$, then we have

$$\int_a^b f(s) ds \geq \int_a^b f(s) \Delta s.$$

Proof. Note that f is an increasing function on $[a, b]$, hence f is integrable on $[a, b]$. Let $\varepsilon > 0$. We now show by induction that

$$S(t) : \int_a^t f(s) + \varepsilon ds - \int_a^t f(s) \Delta s \geq 0$$

holds for all $t \in [a, b]_{\mathbb{T}}$.

1. The statement $S(a)$ is trivially satisfied.
2. Let t be right-scattered and assume that $S(t)$ holds. Then we have

$$\begin{aligned} & \int_a^{\sigma(t)} f(s) + \varepsilon ds - \int_a^{\sigma(t)} f(s) \Delta s \\ & \geq \int_t^{\sigma(t)} f(s) + \varepsilon ds - \int_t^{\sigma(t)} f(s) \Delta s \geq \int_t^{\sigma(t)} f(t) + \varepsilon ds - \mu(t)f(t) = \mu(t)\varepsilon > 0. \end{aligned}$$

Therefore $S(\sigma(t))$ holds.

3. Assume that $S(t)$ holds and $t \neq a$ is right-dense. Since $f(t) \in C_{rd}([a, b]_{\mathbb{T}}, \mathbb{R})$, $f(t)$ is continuous (on $\mathbb{T}$) at t . Then there exists $\delta = \delta(\varepsilon, t)$, such that $|f(s) - f(t)| \leq \varepsilon/2$ holds for all $s \in (t - \delta, t + \delta)_{\mathbb{T}}$. Hence we have for all $\tau \in (t, t + \delta)_{\mathbb{T}}$,

$$\begin{aligned} & \int_a^{\tau} f(s) + \varepsilon ds - \int_a^{\tau} f(s) \Delta s \\ & \geq \int_t^{\tau} f(s) + \varepsilon ds - \int_t^{\tau} f(s) \Delta s \geq (\tau - t)(\varepsilon + f(t) - f(\tau)) \geq \frac{\varepsilon(\tau - t)}{2} > 0. \end{aligned}$$

Therefore $S(\tau)$ holds for all $\tau \in (t, t + \delta)_{\mathbb{T}}$.

4. Now let t be left-dense and suppose $S(\tau)$ is true for all $\tau \in [a, t)_{\mathbb{T}}$, then $S(t)$ holds since the function

$$F(t, \varepsilon) := \int_a^t f(s) + \varepsilon ds - \int_a^t f(s) \Delta s$$

is continuous (on $\mathbb{T}$) with respect to t .

By induction principle ([12], p.4), $S(b)$ is true (i.e. $F(b, \varepsilon) \geq 0$). Moreover, it can be seen that $F(b, \varepsilon)$ is continuous with respect to ε , then $F(b, 0) = \lim_{\varepsilon \rightarrow 0^+} F(b, \varepsilon) \geq 0$.

The proof is completed. $\square$

Corollary 2.1. *Let $\mathbb{T}$ be an arbitrary time scale. Suppose $f : [a, b] \rightarrow \mathbb{R}$ is a decreasing function, where $a, b \in \mathbb{T}$ (b may be ∞). If f is rd-continuous when it is restricted on $[a, b]_{\mathbb{T}}$, then we have*

$$\int_a^b f(s) \Delta s \leq \int_a^b f(s) \Delta s.$$

Corollary 2.2. *Let $\mathbb{T}$ be an arbitrary time scale and c be an arbitrary nonnegative constant. Then we have*

$$\int_a^b \int_a^{t_1} \cdots \int_a^{t_{n-1}} c \Delta t_n \cdots \Delta t_1 \leq \frac{c(b-a)^n}{n!},$$

where $a, b \in \mathbb{T}, a \leq t_{n-1} \leq \cdots \leq t_1 \leq b$.

Proof. Let $b = t_0$. We now show by induction that

$$S(k) : \int_a^{t_{n-k}} \cdots \int_a^{t_{n-1}} c \Delta t_n \cdots \Delta t_{n-k+1} \leq \frac{c(t_{n-k} - a)^k}{k!}$$

holds for all $k \in \{1, 2, \dots, n\}$

1. Clearly, $S(1)$ holds.
2. Now suppose $k \leq n-1$ and that $S(k)$ holds. Then

$$\begin{aligned} & \int_a^{t_{n-(k+1)}} \cdots \int_a^{t_{n-1}} c \Delta t_n \cdots \Delta t_{n-k} \\ & \leq \int_a^{t_{n-(k+1)}} \frac{c(t_{n-k} - a)^k}{k!} \Delta t_{n-k} \leq \int_a^{t_{n-(k+1)}} \frac{c(t_{n-k} - a)^k}{k!} dt_{n-k} \\ & = \frac{c(t_{n-(k+1)} - a)^{k+1}}{(k+1)!}. \end{aligned}$$

Thus, $S(k+1)$ holds.

By induction principle, the proof is completed. $\square$

Corollary 2.3. *Let $\mathbb{T}$ be an arbitrary time scale and c be an arbitrary nonpositive constant. Then we have*

$$\int_a^b \int_a^{t_1} \cdots \int_a^{t_{n-1}} c \Delta t_n \cdots \Delta t_1 \geq \frac{c(b-a)^n}{n!},$$

where $a, b \in \mathbb{T}, a \leq t_{n-1} \leq \cdots \leq t_1 \leq b$.

3. Stability Criteria

Now we start our main work. Let $\mathbb{T}$ be a T -periodic time scale and unbounded above. Consider the stability of the regressive time varying linear dynamic system

$$x^{\Delta\Delta} + p(t)x^{\Delta} + q(t)x = 0, \quad (3.1)$$

where $p(t+T) = p(t)$, $q(t+T) = q(t)$, $p(t), q(t) \in C_{rd}(\mathbb{T}, \mathbb{R})$, $1 - \mu(t)p(t) + \mu^2(t)q(t) \neq 0$, $q(t) \neq 0$ for all $t \in \mathbb{T}$. We assume that $q(t) > 0$ if t is right-dense, and the equation

$$x^\sigma x = q(t) \quad (3.2)$$

has a solution $\phi(t) \in C_{rd}^1(\mathbb{T}, \mathbb{R})$, where x^σ denotes $x(\sigma(t))$.

Remark 3.1. The assumption that Eq. (3.2) exists a solution $\phi(t) \in C_{rd}^1(\mathbb{T}, \mathbb{R})$ can be satisfied for some time scales, such as discrete time scales, continuous time scales and the combination of them.

Note that Eq. (3.1) can be written in the form

$$\begin{pmatrix} x^\Delta \\ y^\Delta \end{pmatrix} = \begin{pmatrix} 0 & 1 \\ -q(t) & -p(t) \end{pmatrix} \begin{pmatrix} x \\ y \end{pmatrix}. \quad (3.3)$$

We assume that $S(t) = \begin{pmatrix} 0 & 1 \\ -q(t) & -p(t) \end{pmatrix}$ and $Y(t) = \begin{pmatrix} x(t) & \bar{x}(t) \\ y(t) & \bar{y}(t) \end{pmatrix} = \Phi_S(t, t_0)$, then the eigenvalues of $Y(t_0 + T)$ are the characteristic multipliers of (3.3). It can be seen that

$$\det Y(t_0 + T) = e_{-p+\mu q}(t_0 + T, t_0) \det Y(t_0) = e_{-p+\mu q}(t_0 + T, t_0).$$

Let ρ_1, ρ_2 denote the characteristic multipliers of (3.3) and

$$\begin{aligned} \mathcal{A} &= x(t_0 + T) + \bar{y}(t_0 + T), \\ \mathcal{B} &= e_{-p+\mu q}(t_0 + T, t_0). \end{aligned} \quad (3.4)$$

Hence ρ_1, ρ_2 satisfy

$$\rho^2 - \mathcal{A}\rho + \mathcal{B} = 0.$$

Obviously,

$$\rho_{1,2} = \frac{\mathcal{A}}{2} \pm \sqrt{\left(\frac{\mathcal{A}}{2}\right)^2 - \mathcal{B}}. \quad (3.5)$$

Note that the value of $\mathcal{B}$ can be easily calculated, then if we can get the value of $\mathcal{A}$, the stability of system (3.1) can be studied by Lemma 2.1.

Theorem 3.1. *We claim that system (3.1) is stable if*

$$\left| \frac{\mathcal{A}}{2} + \sqrt{\left(\frac{\mathcal{A}}{2}\right)^2 - \mathcal{B}} \right| < 1 \quad \text{and} \quad \left| \frac{\mathcal{A}}{2} - \sqrt{\left(\frac{\mathcal{A}}{2}\right)^2 - \mathcal{B}} \right| < 1,$$

and system (3.1) is unstable if

$$\left| \frac{\mathcal{A}}{2} + \sqrt{\left(\frac{\mathcal{A}}{2}\right)^2 - \mathcal{B}} \right| > 1 \quad \text{or} \quad \left| \frac{\mathcal{A}}{2} - \sqrt{\left(\frac{\mathcal{A}}{2}\right)^2 - \mathcal{B}} \right| > 1.$$

Theorem 3.2. *Assume that $\mathcal{B} = 1$. Then we have*

1. *if $|\mathcal{A}| < 2$, system (3.1) is stable;*

2. if $|\mathcal{A}| > 2$, system (3.1) is unstable.

Proof. It follows from (3.5) that $|\rho_1| = |\rho_2| = 1$ and $\rho_1 \neq \rho_2$ as $|\mathcal{A}| < 2$, $\mathcal{B} = 1$, which implies that system (3.1) is stable. The proof of (ii) is similar. $\square$

Remark 3.2. If $\mathbb{T} = \mathbb{R}$, system (3.1) reduces to $x'' + p(t)x' + q(t)x = 0$. If $\mathbb{T} = \mathbb{Z}$, system (3.1) reduces to $\Delta\Delta x + p(t)\Delta x + q(t)x = 0$. In fact, the explicit expression of $\mathcal{A}$ is important to study the stability of the system. Thus, the next section is devoted to presenting an algorithm for the expression of $\mathcal{A}$.

4. Algorithm for the Expression of $\mathcal{A}$

In this section, we are going to focus on the algorithm for $\mathcal{A}$. Note that system (3.3) can be written as

$$\begin{pmatrix} x^\Delta \\ y^\Delta \end{pmatrix} = \begin{pmatrix} 0 & 1 \\ -q(t) & \frac{\phi^\Delta(t)}{\phi(t)} \end{pmatrix} \begin{pmatrix} x \\ y \end{pmatrix} + \begin{pmatrix} 0 \\ (-p(t) - \frac{\phi^\Delta(t)}{\phi(t)})y \end{pmatrix}. \quad (4.1)$$

Let

$$h(t) = -p(t) - \frac{\phi^\Delta(t)}{\phi(t)}, \quad (4.2)$$

thus Eq. (4.1) can be rewritten as

$$\begin{pmatrix} x^\Delta \\ y^\Delta \end{pmatrix} = \begin{pmatrix} 0 & 1 \\ -q(t) & \frac{\phi^\Delta(t)}{\phi(t)} \end{pmatrix} \begin{pmatrix} x \\ y \end{pmatrix} + \begin{pmatrix} 0 \\ h(t)y \end{pmatrix}. \quad (4.3)$$

Let $\cos_\phi(t, t_0) = \cos_\phi(t)$, $\sin_\phi(t, t_0) = \sin_\phi(t)$, hence it can be verified that

$$X(t) = \begin{pmatrix} \cos_\phi(t) & \frac{1}{\phi(t_0)} \sin_\phi(t) \\ -\phi(t) \sin_\phi(t) & \frac{\phi(t)}{\phi(t_0)} \cos_\phi(t) \end{pmatrix} \quad (4.4)$$

is the fundamental matrix solution of the system

$$\begin{pmatrix} x^\Delta \\ y^\Delta \end{pmatrix} = \begin{pmatrix} 0 & 1 \\ -q(t) & \frac{\phi^\Delta(t)}{\phi(t)} \end{pmatrix} \begin{pmatrix} x \\ y \end{pmatrix}. \quad (4.5)$$

Remark 4.1. Let $A(t) = \begin{pmatrix} 0 & 1 \\ -q(t) & \frac{\phi^\Delta(t)}{\phi(t)} \end{pmatrix}$ and we claim that $A(t) \in \mathcal{R}$. On the one hand, $q(t), \phi^\Delta(t)$ are rd-continuous and $\phi(t) \neq 0$, so $A(t) \in C_{rd}(\mathbb{T}, \mathbb{R}^{2 \times 2})$. On the other hand,

$$\det(I + \mu(t)A(t)) = \frac{\phi^\sigma(t)}{\phi(t)} + \mu^2(t)q(t) = \frac{\phi^\sigma(t)(1 + \mu^2(t)\phi^2(t))}{\phi(t)} \neq 0, \quad \text{for all } t \in \mathbb{T},$$

hence $A(t)$ is regressive. Besides we have to consider the rationality of the function $\sin_\phi(t)$ and $\cos_\phi(t)$. We assert that $\sin_\phi(t)$ and $\cos_\phi(t)$ are well defined, since

$$(1 + i\mu(t)\phi(t))(1 - i\mu(t)\phi(t)) = 1 + \mu^2(t)\phi^2(t) \neq 0$$

holds for all $t \in \mathbb{T}$.

The solution of system (4.3) satisfying $\begin{pmatrix} x(t_0) \\ y(t_0) \end{pmatrix} = \begin{pmatrix} x_0 \\ y_0 \end{pmatrix}$ can be represented as

$$\begin{pmatrix} x(t) \\ y(t) \end{pmatrix} = X(t) \begin{pmatrix} x_0 \\ y_0 \end{pmatrix} + \int_{t_0}^t X(t)X^{-1}(s)(I + \mu(s)A(s))^{-1} \begin{pmatrix} 0 \\ h(s)y(s) \end{pmatrix} \Delta s. \quad (4.6)$$

Note that

$$X^{-1}(s) = \begin{pmatrix} \frac{\cos_\phi(s)}{e_{\mu\phi^2}(s)} & -\frac{\sin_\phi(s)}{\phi(s)e_{\mu\phi^2}(s)} \\ \frac{\phi(0)\sin_\phi(s)}{e_{\mu\phi^2}(s)} & \frac{\phi(0)\cos_\phi(s)}{\phi(s)e_{\mu\phi^2}(s)} \end{pmatrix}, \quad I + \mu(s)A(s) = \begin{pmatrix} 1 & \mu(s) \\ -\mu(s)q(s) & \frac{\phi^\sigma(s)}{\phi(s)} \end{pmatrix},$$

$$\det(I + \mu(s)A(s)) = \frac{\phi^\sigma(s)}{\phi(s)} + \mu^2(s)q(s) = \frac{\phi^\sigma(s)(1 + \mu^2(s)\phi^2(s))}{\phi(s)},$$

and

$$(I + \mu(s)A(s))^{-1} = \begin{pmatrix} \frac{1}{1 + \mu^2(s)\phi^2(s)} & \frac{-\mu(s)\phi(s)}{\phi^\sigma(s)(1 + \mu^2(s)\phi^2(s))} \\ \frac{\mu(s)\phi^2(s)}{1 + \mu^2(s)\phi^2(s)} & \frac{\phi(s)}{\phi^\sigma(s)(1 + \mu^2(s)\phi^2(s))} \end{pmatrix}.$$

Substituting them in Eq. (4.6), then we have

$$\begin{pmatrix} x(t) \\ y(t) \end{pmatrix} = \begin{pmatrix} \cos_\phi(t) & \frac{1}{\phi(t_0)}\sin_\phi(t) \\ -\phi\sin_\phi(t) & \frac{\phi(t)}{\phi(t_0)}\cos_\phi(t) \end{pmatrix} \begin{pmatrix} x_0 \\ y_0 \end{pmatrix} + \int_{t_0}^t \begin{pmatrix} h(s)\frac{-\mu(s)\phi(s)\cos_\phi(t,s) + \sin_\phi(t,s)}{\phi^\sigma(s)(1 + \mu^2(s)\phi^2(s))}y(s) \\ h(s)\frac{\mu(s)\phi(s)\phi(t)\sin_\phi(t,s) + \phi(t)\cos_\phi(t,s)}{\phi^\sigma(s)(1 + \mu^2(s)\phi^2(s))}y(s) \end{pmatrix} \Delta s. \quad (4.7)$$

Let $\begin{pmatrix} x(t) \\ y(t) \end{pmatrix}, \begin{pmatrix} \bar{x}(t) \\ \bar{y}(t) \end{pmatrix}$ denote the solutions of system (4.3)(i.e. (3.3)) that satisfy the initial condition $\begin{pmatrix} x(t_0) \\ y(t_0) \end{pmatrix} = \begin{pmatrix} 1 \\ 0 \end{pmatrix}, \begin{pmatrix} \bar{x}(t_0) \\ \bar{y}(t_0) \end{pmatrix} = \begin{pmatrix} 0 \\ 1 \end{pmatrix}$, respectively. By Eq. (3.4) we get

$$\mathcal{A} = x(t_0 + T) + \bar{y}(t_0 + T). \quad (4.8)$$

Now let's use the approximation method to calculate $\mathcal{A}$. We assume that

$$\begin{pmatrix} x_0(t) \\ y_0(t) \end{pmatrix} = X(t) \begin{pmatrix} 1 \\ 0 \end{pmatrix} = \begin{pmatrix} \cos_\phi(t) \\ -\phi(t) \sin_\phi(t) \end{pmatrix}.$$

And if $\begin{pmatrix} x_{n-1}(t) \\ y_{n-1}(t) \end{pmatrix}$ was given, then we define $\begin{pmatrix} x_n(t) \\ y_n(t) \end{pmatrix}$ inductively by

$$\begin{pmatrix} x_n(t) \\ y_n(t) \end{pmatrix} = X(t) \begin{pmatrix} 1 \\ 0 \end{pmatrix} + \int_{t_0}^t \begin{pmatrix} h(s) \frac{-\mu(s)\phi(s) \cos_\phi(t, s) + \sin_\phi(t, s)}{\phi^\sigma(s)(1 + \mu^2(s)\phi^2(s))} y_{n-1}(s) \\ h(s) \frac{\mu(s)\phi(s)\phi(t) \sin_\phi(t, s) + \phi(t) \cos_\phi(t, s)}{\phi^\sigma(s)(1 + \mu^2(s)\phi^2(s))} y_{n-1}(s) \end{pmatrix} \Delta s. \quad (4.9)$$

Similarly, we assume that

$$\begin{pmatrix} \bar{x}_0(t) \\ \bar{y}_0(t) \end{pmatrix} = X(t) \begin{pmatrix} 0 \\ 1 \end{pmatrix} = \begin{pmatrix} \frac{1}{\phi(t_0)} \sin_\phi(t) \\ \frac{\phi(t)}{\phi(t_0)} \cos_\phi(t) \end{pmatrix}.$$

And if $\begin{pmatrix} \bar{x}_{n-1}(t) \\ \bar{y}_{n-1}(t) \end{pmatrix}$ was given, then we define $\begin{pmatrix} \bar{x}_n(t) \\ \bar{y}_n(t) \end{pmatrix}$ inductively by

$$\begin{pmatrix} \bar{x}_n(t) \\ \bar{y}_n(t) \end{pmatrix} = X(t) \begin{pmatrix} 0 \\ 1 \end{pmatrix} + \int_{t_0}^t \begin{pmatrix} h(s) \frac{-\mu(s)\phi(s) \cos_\phi(t, s) + \sin_\phi(t, s)}{\phi^\sigma(s)(1 + \mu^2(s)\phi^2(s))} \bar{y}_{n-1}(s) \\ h(s) \frac{\mu(s)\phi(s)\phi(t) \sin_\phi(t, s) + \phi(t) \cos_\phi(t, s)}{\phi^\sigma(s)(1 + \mu^2(s)\phi^2(s))} \bar{y}_{n-1}(s) \end{pmatrix} \Delta s. \quad (4.10)$$

It is easy to see that

$$\begin{cases} x_1(t) = \cos_\phi(t) - \int_{t_0}^t h(s)\phi(s)\sin_\phi(s) \frac{-\mu(s)\phi(s)\cos_\phi(t,s) + \sin_\phi(t,s)}{\phi^\sigma(s)(1+\mu^2(s)\phi^2(s))} \Delta s, \\ y_1(t) = -\phi(t)\sin_\phi(t) - \phi(t) \int_{t_0}^t h(s)\phi(s)\sin_\phi(s) \frac{\mu(s)\phi(s)\sin_\phi(t,s) + \cos_\phi(t,s)}{\phi^\sigma(s)(1+\mu^2(s)\phi^2(s))} \Delta s, \\ \bar{y}_1(t) = \frac{\phi(t)}{\phi(t_0)} \cos_\phi(t) + \frac{\phi(t)}{\phi(t_0)} \int_{t_0}^t h(s)\phi(s)\cos_\phi(s) \frac{\mu(s)\phi(s)\sin_\phi(t,s) + \cos_\phi(t,s)}{\phi^\sigma(s)(1+\mu^2(s)\phi^2(s))} \Delta s. \end{cases} \quad (4.11)$$

Remark 4.2. Note that $\bar{x}_1(t)$ doesn't work for recursion, so we don't have to figure it out. For the same reason, $\bar{x}_n(t)$ also needn't to be calculated.

Let

$$\begin{aligned} P(t,s) &= \frac{-\mu(s)\phi(s)\cos_\phi(t,s) + \sin_\phi(t,s)}{\phi^\sigma(s)(1+\mu^2(s)\phi^2(s))}, \\ Q(t,s) &= \frac{\mu(s)\phi(s)\phi(t)\sin_\phi(t,s) + \phi(t)\cos_\phi(t,s)}{\phi^\sigma(s)(1+\mu^2(s)\phi^2(s))}. \end{aligned} \quad (4.12)$$

It can be seen that

$$\begin{aligned} \sin_\phi(\sigma(s), t) &= \frac{e_{i\phi}(\sigma(s), t) - e_{-i\phi}(\sigma(s), t)}{2i} \\ &= \frac{(1 + i\mu(s)\phi(s))e_{i\phi}(s, t) - (1 - i\mu(s)\phi(s))e_{-i\phi}(s, t)}{2i} \\ &= \sin_\phi(s, t) + \mu(s)\phi(s)\cos_\phi(s, t), \end{aligned}$$

and

$$\sin_\phi(t, s) = -e_{\mu\phi^2}(t, s)\sin_\phi(s, t).$$

Similarly, we have

$$\cos_\phi(\sigma(s), t) = \cos_\phi(s, t) - \mu(s)\phi(s)\sin_\phi(s, t),$$

and

$$\cos_\phi(t, s) = e_{\mu\phi^2}(t, s)\cos_\phi(s, t).$$

Then the function P, Q can be simplified as

$$P(t,s) = \frac{1}{\phi^\sigma(s)} \sin_\phi(t, \sigma(s)), \quad Q(t,s) = \frac{\phi(t)}{\phi^\sigma(s)} \cos_\phi(t, \sigma(s)). \quad (4.13)$$

Using Eq. (4.9), (4.10), (4.11) we obtain

$$\left\{ \begin{array}{l} x_2(t) = \cos_\phi(t) - \int_{t_0}^t h(s) \sin_\phi(s) P(t, s) \phi(s) \Delta s \\ \quad - \int_{t_0}^t \int_{t_0}^{t_1} h(t_1) h(t_2) \sin_\phi(t_2) P(t, t_1) Q(t_1, t_2) \phi(t_2) \Delta t_2 \Delta t_1, \\ y_2(t) = -\phi(t) \sin_\phi(t) - \int_{t_0}^t h(s) \sin_\phi(s) Q(t, s) \phi(s) \Delta s \\ \quad - \int_{t_0}^t \int_{t_0}^{t_1} h(t_1) h(t_2) \sin_\phi(t_2) Q(t, t_1) Q(t_1, t_2) \phi(t_2) \Delta t_2 \Delta t_1, \\ \bar{y}_2(t) = \frac{\phi(t)}{\phi(t_0)} \cos_\phi(t) + \frac{1}{\phi(t_0)} \int_{t_0}^t h(s) \cos_\phi(s) Q(t, s) \phi(s) \Delta s \\ \quad + \frac{1}{\phi(t_0)} \int_{t_0}^t \int_{t_0}^{t_1} h(t_1) h(t_2) \cos_\phi(t_2) Q(t, t_1) Q(t_1, t_2) \phi(t_2) \Delta t_2 \Delta t_1. \end{array} \right. \quad (4.14)$$

Let

$$\left\{ \begin{array}{l} u_k(t) = - \int_{t_0}^t \int_{t_0}^{t_1} \cdots \int_{t_0}^{t_{k-1}} \phi(t_k) \sin_\phi(t_k) Q(t_{k-1}, t_k) \\ \quad \cdots Q(t_1, t_2) P(t, t_1) \prod_{i=1}^k h(t_i) \Delta t_k \cdots \Delta t_1, \\ v_k(t) = - \int_{t_0}^t \int_{t_0}^{t_1} \cdots \int_{t_0}^{t_{k-1}} \phi(t_k) \sin_\phi(t_k) Q(t_{k-1}, t_k) \\ \quad \cdots Q(t_1, t_2) Q(t, t_1) \prod_{i=1}^k h(t_i) \Delta t_k \cdots \Delta t_1, \\ \bar{v}_k(t) = \frac{1}{\phi(t_0)} \int_{t_0}^t \int_{t_0}^{t_1} \cdots \int_{t_0}^{t_{k-1}} \phi(t_k) \cos_\phi(t_k) Q(t_{k-1}, t_k) \\ \quad \cdots Q(t_1, t_2) Q(t, t_1) \prod_{i=1}^k h(t_i) \Delta t_k \cdots \Delta t_1, \\ (t_0 \leq t_k \leq t_{k-1} \leq \cdots \leq t_1 \leq t, \quad k = 1, 2, \dots). \end{array} \right. \quad (4.15)$$

For Eq. (4.11), we have

$$\left\{ \begin{array}{l} x_1(t) = \cos_\phi(t) + u_1(t), \\ y_1(t) = -\phi(t) \sin_\phi(t) + v_1(t), \\ \bar{y}_1(t) = \frac{\phi(t)}{\phi(t_0)} \cos_\phi(t) + \bar{v}_1(t). \end{array} \right. \quad (4.16)$$

For Eq. (4.14), we have

$$\left\{ \begin{array}{l} x_2(t) = \cos_\phi(t) + u_1(t) + u_2(t), \\ y_2(t) = -\phi(t) \sin_\phi(t) + v_1(t) + v_2(t), \\ \bar{y}_2(t) = \frac{\phi(t)}{\phi(t_0)} \cos_\phi(t) + \bar{v}_1(t) + \bar{v}_2(t). \end{array} \right. \quad (4.17)$$

Now we take an inductive assumption that

$$\begin{cases} x_k(t) = \cos_\phi(t) + u_1(t) + \cdots + u_k(t), \\ y_k(t) = -\phi(t) \sin_\phi(t) + v_1(t) + \cdots + v_k(t), \\ \bar{y}_k(t) = \frac{\phi(t)}{\phi(t_0)} \cos_\phi(t) + \bar{v}_1(t) + \cdots + \bar{v}_k(t). \end{cases} \quad (4.18)$$

According to Eq. (4.9) and (4.10),

$$\begin{cases} x_{k+1}(t) = \cos_\phi(t) + \int_{t_0}^t h(s)P(t, s)y_k(s)\Delta s, \\ y_{k+1}(t) = -\phi(t) \sin_\phi(t) + \int_{t_0}^t h(s)Q(t, s)y_k(s)\Delta s, \\ \bar{y}_{k+1}(t) = \frac{\phi(t)}{\phi(t_0)} \cos_\phi(t) + \int_{t_0}^t h(s)Q(t, s)\bar{y}_k(s)\Delta s. \end{cases} \quad (4.19)$$

Substituting Eq. (4.18) into Eq. (4.19), we get

$$\begin{cases} x_{k+1}(t) = \cos_\phi(t) + u_1(t) + \cdots + u_{k+1}(t), \\ y_{k+1}(t) = -\phi(t) \sin_\phi(t) + v_1(t) + \cdots + v_{k+1}(t), \\ \bar{y}_{k+1}(t) = \frac{\phi(t)}{\phi(t_0)} \cos_\phi(t) + \bar{v}_1(t) + \cdots + \bar{v}_{k+1}(t). \end{cases} \quad (4.20)$$

This implies that Eq.(4.18) holds for all $k \in \mathbb{N}$.

Let $[t_0, t_0 + T]_{\mathbb{T}} := [t_0, t_0 + T] \cap \mathbb{T}$. For the bounded closed interval $[t_0, t_0 + T]_{\mathbb{T}}$, consider the series

$$y_0(t) + \sum_{k=1}^{\infty} [y_k(t) - y_{k-1}(t)], \quad t \in [t_0, t_0 + T]_{\mathbb{T}}, \quad (4.21)$$

and the partial sum

$$y_0(t) + \sum_{k=1}^n [y_k(t) - y_{k-1}(t)] = y_n(t).$$

So if we want to prove the sequence $\{y_n(t)\}$ is uniformly convergent on $[t_0, t_0 + T]_{\mathbb{T}}$, just show that series (4.21) converges uniformly on $[t_0, t_0 + T]_{\mathbb{T}}$. Note that $\sin_\phi(t, s), \cos_\phi(t, s), \phi(t), \mu(t), h(t)$ are rd-continuous. By lemma 2.2, we have the functions

$$|\phi(t)|, |\sin_\phi(t)|, |\cos_\phi(t)|, |h(t)|$$

are all bounded on compact set $[t_0, t_0 + T]_{\mathbb{T}}$. By Eq. (4.13), since $\phi(t) \neq 0$, it can be seen that $|P(t, s)|, |Q(t, s)|$ are all bounded on $[t_0, t_0 + T]_{\mathbb{T}} \times [t_0, t_0 + T]_{\mathbb{T}}$. Let M denote their common upper bound, so we have

$$\begin{aligned} & |y_k(t) - y_{k-1}(t)| = |v_k(t)| \\ & = \left| \int_{t_0}^t \int_{t_0}^{t_1} \cdots \int_{t_0}^{t_{k-1}} \phi(t_k) \sin_\phi(t_k) Q(t_{k-1}, t_k) \cdots Q(t_1, t_2) Q(t, t_1) \prod_{i=1}^k h(t_i) \Delta t_k \cdots \Delta t_1 \right| \end{aligned}$$

$$\begin{aligned}
&\leq \int_{t_0}^t \int_{t_0}^{t_1} \cdots \int_{t_0}^{t_{k-1}} \left| \phi(t_k) \sin_\phi(t_k) Q(t_{k-1}, t_k) \cdots Q(t_1, t_2) Q(t, t_1) \prod_{i=1}^k h(t_i) \right| \Delta t_k \cdots \Delta t_1 \\
&\leq \int_{t_0}^t \int_{t_0}^{t_1} \cdots \int_{t_0}^{t_{k-1}} M^{2k+2} \Delta t_k \cdots \Delta t_1 \\
&\leq \frac{M^{2k+2} (t - t_0)^k}{k!} \leq \frac{M^{2k+2} T^k}{k!}, \quad t_0 \leq t \leq t_0 + T.
\end{aligned} \tag{4.22}$$

The third inequality in (4.22) is derived from Corollary (2.2). According to Weierstrass Discriminane, series (4.21) is uniformly convergent on $[t_0, t_0 + T]_{\mathbb{T}}$, thus the sequence $\{y_k(t)\}$ is uniformly convergent on $[t_0, t_0 + T]_{\mathbb{T}}$. Now assume

$$\lim_{k \rightarrow \infty} y_k(t) = y^*(t).$$

By lemma 2.3 we get $y^*(t)$ is rd-continuous on $[t_0, t_0 + T]_{\mathbb{T}}$. Hence

$$\begin{aligned}
\lim_{k \rightarrow \infty} y_k(t) &= -\phi(t) \sin_\phi(t) + \lim_{k \rightarrow \infty} \int_{t_0}^t h(s) Q(t, s) y_{k-1}(s) \Delta s \\
&= -\phi(t) \sin_\phi(t) + \int_{t_0}^t \lim_{k \rightarrow \infty} h(s) Q(t, s) y_{k-1}(s) \Delta s,
\end{aligned} \tag{4.23}$$

i.e.,

$$y^*(t) = -\phi(t) \sin_\phi(t) + \int_{t_0}^t h(s) Q(t, s) y^*(s) \Delta s.$$

In the same way, the sequence $\{x_k(t)\}$ uniformly converges to $x^*(t)$ which satisfies

$$x^*(t) = \cos_\phi(t) + \int_{t_0}^t h(s) P(t, s) y^*(s) \Delta s.$$

That is to say $\begin{pmatrix} x^*(t) \\ y^*(t) \end{pmatrix}$ is the solution of system (4.3) with the initial condition

$$\begin{pmatrix} x^*(t_0) \\ y^*(t_0) \end{pmatrix} = \begin{pmatrix} 1 \\ 0 \end{pmatrix}.$$

For the theorem of existence and uniqueness of solution, $x^*(t) = x(t)$, $y^*(t) = y(t)$.

Let's do the same things for $\bar{y}_n(t)$. Finally we have $\begin{pmatrix} x_n(t) \\ y_n(t) \end{pmatrix}$ uniformly converges

to $\begin{pmatrix} x(t) \\ y(t) \end{pmatrix}$ and $\bar{y}_n(t)$ uniformly converges to $\bar{y}(t)$. Let

$$\begin{cases} \mathcal{A}_0 = x_0(t_0 + T) + \bar{y}_0(t_0 + T), \\ \mathcal{A}_1 = u_1(t_0 + T) + \bar{v}_1(t_0 + T), \\ \dots \\ \mathcal{A}_n = u_n(t_0 + T) + \bar{v}_n(t_0 + T). \end{cases}$$

By $\mathcal{A} = x(t_0 + T) + \bar{y}(t_0 + T)$ and Eq. (4.18), we get

$$\mathcal{A} = \sum_{n=0}^{\infty} \mathcal{A}_n. \quad (4.24)$$

Now we evaluate $\mathcal{A}_n (n = 0, 1, 2, 3, \dots)$:

$$\begin{aligned} \mathcal{A}_0 &= \left(1 + \frac{\phi(t_0 + T)}{\phi(t_0)}\right) \cos_{\phi}(t_0 + T) \\ \mathcal{A}_1 &= \int_{t_0}^{t_0+T} \left(\frac{1}{\phi(t_0)} \cos_{\phi}(t_1) Q(t_0 + T, t_1) - \sin_{\phi}(t_1) P(t_0 + T, t_1)\right) \phi(t_1) h(t_1) \Delta t_1 \\ \mathcal{A}_n &= - \int_{t_0}^{t_0+T} \int_{t_0}^{t_1} \cdots \int_{t_0}^{t_{n-1}} \phi(t_n) \sin_{\phi}(t_n) Q(t_{n-1}, t_n) \\ &\quad \cdots Q(t_1, t_2) P(t_0 + T, t_1) \prod_{i=1}^n h(t_i) \Delta t_n \cdots \Delta t_1 \\ &\quad + \frac{1}{\phi(t_0)} \int_{t_0}^{t_0+T} \int_{t_0}^{t_1} \cdots \int_{t_0}^{t_{n-1}} \phi(t_n) \cos_{\phi}(t_n) Q(t_{n-1}, t_n) \\ &\quad \cdots Q(t_1, t_2) Q(t_0 + T, t_1) \prod_{i=1}^n h(t_i) \Delta t_n \cdots \Delta t_1 \\ &= \int_{t_0}^{t_0+T} \int_{t_0}^{t_1} \cdots \int_{t_0}^{t_{n-1}} \left(\frac{1}{\phi(t_0)} \cos_{\phi}(t_n) Q(t_0 + T, t_1) - \sin_{\phi}(t_n) P(t_0 + T, t_1)\right) \cdot \\ &\quad \phi(t_n) Q(t_{n-1}, t_n) \cdots Q(t_1, t_2) \prod_{i=1}^n h(t_i) \Delta t_n \cdots \Delta t_1, \quad n \geq 2. \end{aligned} \quad (4.25)$$

Thus we have

$$\begin{aligned} \mathcal{A} &= \left(1 + \frac{\phi(t_0 + T)}{\phi(t_0)}\right) \cos_{\phi}(t_0 + T) \\ &\quad + \int_{t_0}^{t_0+T} \left(\frac{1}{\phi(t_0)} \cos_{\phi}(t_1) Q(t_0 + T, t_1) - \sin_{\phi}(t_1) P(t_0 + T, t_1)\right) \phi(t_1) h(t_1) \Delta t_1 \\ &\quad + \sum_{n=2}^{\infty} \int_{t_0}^{t_0+T} \int_{t_0}^{t_1} \cdots \int_{t_0}^{t_{n-1}} \left(\frac{1}{\phi(t_0)} \cos_{\phi}(t_n) Q(t_0 + T, t_1) - \sin_{\phi}(t_n) P(t_0 + T, t_1)\right) \cdot \\ &\quad \phi(t_n) Q(t_{n-1}, t_n) \cdots Q(t_1, t_2) \prod_{i=1}^n h(t_i) \Delta t_n \cdots \Delta t_1. \end{aligned} \quad (4.26)$$

The formula above can be used for approximations and error estimates. Let

$$h(t, s) = \left(\frac{1}{\phi(t_0)} \cos_{\phi}(t) Q(t_0 + T, s) - \sin_{\phi}(t) P(t_0 + T, s)\right) \cdot \phi(t).$$

Then we have

$$|\mathcal{A}_n| \leq \int_{t_0}^{t_0+T} \int_{t_0}^{t_1} \cdots \int_{t_0}^{t_{n-1}} K_1 K_2^{n-1} K_3^n \Delta t_n \cdots \Delta t_1 \leq \frac{K_1 K_2^{n-1} K_3^n T^n}{n!},$$

where K_1, K_2, K_3 are upper bounds of $|h(t, s)|$, $|Q(t, s)|$ and $|h(t)|$ respectively. Let

$$\mathcal{A}(n) = \mathcal{A}_0 + \mathcal{A}_1 + \cdots + \mathcal{A}_n, \quad (4.27)$$

and we have the following error estimate

$$|\mathcal{A} - \mathcal{A}(n)| \leq \sum_{k=n+1}^{\infty} \frac{K_1}{K_2} \frac{(K_2 K_3 T)^k}{k!} = \frac{K_1}{K_2} \left(e^{K_2 K_3 T} - \sum_{k=0}^n \frac{(K_2 K_3 T)^k}{k!} \right). \quad (4.28)$$

Theorem 4.1. *The expression of $\mathcal{A}$ mentioned in Theorem 3.1 is*

$$\begin{aligned} \mathcal{A} = & \left(1 + \frac{\phi(t_0 + T)}{\phi(t_0)} \right) \cos_{\phi}(t_0 + T) \\ & + \int_{t_0}^{t_0+T} \left(\frac{1}{\phi(t_0)} \cos_{\phi}(t_1) Q(t_0 + T, t_1) - \sin_{\phi}(t_1) P(t_0 + T, t_1) \right) \phi(t_1) h(t_1) \Delta t_1 \\ & + \sum_{n=2}^{\infty} \int_{t_0}^{t_0+T} \int_{t_0}^{t_1} \cdots \int_{t_0}^{t_{n-1}} \left(\frac{1}{\phi(t_0)} \cos_{\phi}(t_n) Q(t_0 + T, t_1) - \sin_{\phi}(t_n) P(t_0 + T, t_1) \right) \\ & \times \phi(t_n) Q(t_{n-1}, t_n) \cdots Q(t_1, t_2) \prod_{i=1}^n h(t_i) \Delta t_n \cdots \Delta t_1, \end{aligned} \quad (4.29)$$

and the expression of $\mathcal{B}$ is

$$\mathcal{B} = e_{-p+\mu q}(t_0 + T, t_0).$$

Theorem 4.2. *Let $\mathbb{T}$ be an arbitrary discrete time scale and there are k points in $[t_0, t_0 + T]_{\mathbb{T}}$, then equation (4.29) can be simplified as*

$$\begin{aligned} \mathcal{A} = \mathcal{A}(k) = & \left(1 + \frac{\phi(t_0 + T)}{\phi(t_0)} \right) \cos_{\phi}(t_0 + T) \\ & + \int_{t_0}^{t_0+T} \left(\frac{1}{\phi(t_0)} \cos_{\phi}(t_1) Q(t_0 + T, t_1) - \sin_{\phi}(t_1) P(t_0 + T, t_1) \right) \phi(t_1) h(t_1) \Delta t_1 \\ & + \sum_{n=2}^k \int_{t_0}^{t_0+T} \int_{t_0}^{t_1} \cdots \int_{t_0}^{t_{n-1}} \left(\frac{1}{\phi(t_0)} \cos_{\phi}(t_n) Q(t_0 + T, t_1) - \sin_{\phi}(t_n) P(t_0 + T, t_1) \right) \\ & \times \phi(t_n) Q(t_{n-1}, t_n) \cdots Q(t_1, t_2) \prod_{i=1}^n h(t_i) \Delta t_n \cdots \Delta t_1, \end{aligned} \quad (4.30)$$

where $\sum_{n=2}^1 (\cdot) := 0$.

Proof. Now we show that $\mathcal{A}_n = 0$ if $n \geq k + 1$, where $\mathcal{A}_n$ is defined in equation (4.25). Let's abbreviate $\mathcal{A}_n$ as $\int_{t_0}^{t_0+T} \int_{t_0}^{t_1} \cdots \int_{t_0}^{t_{n-1}} R(\cdot) \Delta t_n \cdots \Delta t_1$, where $t_0 \leq t_{n-1} < t_{n-2} < \cdots < t_1 < t_0 + T$. Note that the number of the points in $[t_0, t_0 + T]_{\mathbb{T}}$ is k , which is less than n . Hence there must exists an element of the set $\{t_i | i = 1, 2, \dots, n-1\}$ equal to t_0 , which implies that $\mathcal{A}_n = 0$. The proof is completed. $\square$

Theorem 4.3. Consider the Hill's equation ([14, 15])

$$x^{\Delta\Delta}(t) + q(t)x(t) = 0, \quad (4.31)$$

where $q(t)$ and $\mathbb{T}$ are both T -periodic, then the expression of $\mathcal{A}$ of (4.31) can be simplified as

$$\begin{aligned} \mathcal{A} = & \left(1 + \frac{\phi(t_0 + T)}{\phi(t_0)}\right) \cos_\phi(t_0 + T) \\ & + \int_{t_0}^{t_0+T} \left(\frac{1}{\phi(t_0)} \cos_\phi(t_1) Q(t_0 + T, t_1) - \sin_\phi(t_1) P(t_0 + T, t_1) \right) \phi(t_1) h(t_1) \Delta t_1 \\ & + \sum_{n=2}^{\infty} \int_{t_0}^{t_0+T} \int_{t_0}^{t_1} \cdots \int_{t_0}^{t_{n-1}} \left(\frac{1}{\phi(t_0)} \cos_\phi(t_n) Q(t_0 + T, t_1) - \sin_\phi(t_n) P(t_0 + T, t_1) \right) \\ & \times (-1)^n \phi^\Delta(t_1) \prod_{i=2}^n \frac{\sin_\phi(t_{i-1}, \sigma(t_i)) \phi^\Delta(t_i)}{\phi^\sigma(t_i)} \Delta t_n \cdots \Delta t_1. \end{aligned} \quad (4.32)$$

Proof. The proof is an algebraic process, so we omit it. $\square$

Theorem 4.4 ([48]). If the time scale $\mathbb{T} = \mathbb{R}$ and $\mathcal{B} = 1$, then equation (4.29) can be simplified as

$$\begin{aligned} \mathcal{A} = & 2 \cos \Phi(t_0 + T) \\ & + \sum_{n=1}^{\infty} \frac{1}{2^{2n-1}} \int_{t_0}^{t_0+T} \int_{t_0}^{t_1} \cdots \int_{t_0}^{t_{2n-1}} \cos \Psi(t_1, \dots, t_{2n}) \cdot \prod_{i=1}^{2n} h(t_i) dt_{2n} \cdots dt_1, \end{aligned}$$

where

$$\begin{aligned} \Phi(t) &= \int_{t_0}^t \phi(\tau) d\tau, \quad \Phi(t, s) = \int_s^t \phi(\tau) d\tau, \\ \Psi(t_1, \dots, t_{2n}) &= \Phi(t_0 + T) - 2\Phi(t_1, t_2) - 2\Phi(t_3, t_4) - \cdots - 2\Phi(t_{2n-1}, t_{2n}). \end{aligned}$$

Remark 4.3. Theoretically, we show that this approach is also valid for critical case: the system has the same characteristic multipliers with modulus equal to one. In a similar manner, we can get an expression of $\bar{x}(t_0 + T)$ in the form of a series. That is, combined with the previous discussion, the matrix $\Phi_A(t_0, t_0 + T)$ also has an expression in the form of a convergent series. Note that the system we studied in critical case is stable if and only if $\Phi_A(t_0, t_0 + T) - \rho I = 0$, where ρ is the characteristic multipliers. Then we can get the error estimate like (4.27) and (4.28) to analyse the stability. Moreover, we see that the stability of the nonhomogeneous system $x^{\Delta\Delta} + p(t)x^\Delta + q(t)x = f(t)$ is equivalent to the system $x^{\Delta\Delta} + p(t)x^\Delta + q(t)x = 0$.

5. Program for the algorithm

The following Matlab program is designed for calculating the value of $\mathcal{A}(n)$ and $\mathcal{B}$ mentioned above. One can run the following program by Matlab R2018a.

Program 1.

```

% This program was designed for calculating the value of
% A(n) and B mentioned in this paper.

%=====
% Users should set the functions p(t), q(t) and q_diff in
% advance in section 2 of this script, where q_diff is
% the derivative function of q(t) in continuous part(If
% there is no continuous part, take q_diff=0).
%=====
% discrete part: Input the discrete points in the form
% of a row vector from small to large.
%=====
% continuous part: Input the ends of continuous intervals
% in the form of a matrix, and its first and second row
% record the left and right ends from small to large,
% respectively.
clc

global discrete_part continuous_part time_scale;
discrete_part=input('Enter the discrete point: ');
continuous_part=input('Enter the continuous interval: ');

if isequal(continuous_part,[])
time_scale=discrete_part;
else
time_scale=sort([discrete_part,continuous_part(1,:),...
continuous_part(2,:)]);
end
B=exp_fun(@(t) -p(t)+mu(t)*q(t),time_scale(end),time_scale(1));

if isequal(continuous_part,[])
A=valueOfDelta;
fprintf('The value of A is %f \n',A);
fprintf('The value of B is %f \n',B);
fprintf('The modulus of multipliers are %f %f\n',...
abs((A-sqrt(A^2-4*B))/2),abs((A+sqrt(A^2-4*B))/2));
elseif isequal(discrete_part,[])
n=input('n:');
A=Consum(n);
fprintf(['The value of A(',num2str(n),') is %f \n'],A);
fprintf('The value of B is %f \n',B);
fprintf(['The ',num2str(n),'th approximate modulus are %f %f\n'],...
abs((A-sqrt(A^2-4*B))/2),abs((A+sqrt(A^2-4*B))/2));
else
n=input('n:');
A=Delta_H(n);
fprintf(['The value of A(',num2str(n),') is %f \n'],A);
fprintf('The value of B is %f \n',B);
fprintf(['The ',num2str(n),'th approximate modulus are %f %f\n'],...

```

```

abs((A-sqrt(A^2-4*B))/2),abs((A+sqrt(A^2-4*B))/2));
end
clear global;

%%
%Users should define the following functions:p,q,q_diff
function f=p(t)
if t==pi
f=0.25;
else
f=0;
end
end

function f=q(t)
f=1;
end

%the derivative function of q(t) in continuous part
function f=q_diff(t)
f=0;
end

%%
function f=mu(t)
global discrete_part continuous_part time_scale;
if ismember(t,discrete_part) || ismember(t,continuous_part(2,:))
if t==time_scale(end)
f=mu(time_scale(1));
else
for i=1:length(time_scale)
if t==time_scale(i)
f=time_scale(i+1)-time_scale(i);
end
end
end
else
f=0;
end
end

function f=sigma(t)
f=t+mu(t);
end

function f=phi(t)
global discrete_part continuous_part time_scale;
if isequal(continuous_part,[])
exphi=NaN(1,length(discrete_part));

```

```

exphi(1)=1;
for i=2:length(discrete_part)
    exphi(i)=q(discrete_part(i-1))/exphi(i-1);
end
for i=1:length(discrete_part)
    if t==discrete_part(i)
        f=exphi(i);
    end
end

else
leftends=continuous_part(1,:);rightends=continuous_part(2,:);
if ~(ismember(t,discrete_part) || ismember(t,rightends))
f=sqrt(q(t));

elseif t<leftends(end)
n=1;tt=t;
while ~ismember(tt,leftends)
n=n+1;tt=sigma(tt);
end
temp=NaN(1,n);temp(n)=sqrt(q(tt));k=1;
while ~isequal(tt,time_scale(k))
k=k+1;
end
for i=n-1:-1:1
temp(i)=q(time_scale(k-n+i))./temp(i+1);
end
f=temp(1);

elseif t==time_scale(end)
f=phi(time_scale(1));

else
k=1;
while ~isequal(t,time_scale(k))
k=k+1;
end
n=length(time_scale)-k+1;
temp=NaN(1,n);
temp(n)=phi(time_scale(1));
for i=n-1:-1:1
temp(i)=q(time_scale(length(time_scale)-n+i))./temp(i+1);
end
f=temp(1);
end
end
end

function f=delta_int(g,t,s)

```

```

% where g is a function handle, t and s are up and low, respectively.
global continuous_part; ss=s; sum=0;
if isequal(continuous_part,[])
while ss<t
sum=sum+mu(ss).*g(ss);
ss=sigma(ss);
end
else
rightends=continuous_part(2,:);
while ss<t
if ss==sigma(ss)
k=1;
while ss>rightends(k)
k=k+1;
end
if rightends(k)>t
sum=sum+integral(@(x) arrayfun(@(x)g(x),x),ss,t);
else
sum=sum+integral(@(x) arrayfun(@(x)g(x),x),ss,rightends(k));
end
ss=rightends(k);
else
sum=sum+mu(ss).*g(ss);
ss=sigma(ss);
end
end
end
f=sum;
end

function f=cylinder_fun(g,t)
% where g is a function handle.
if mu(t)==0
f=g(t);
else
f=log(1+mu(t).*g(t))./mu(t);
end
end

function f=exp_fun(g,t,s)
% where g is a function handle, t and s are up and low, respectively.
cylinder_g=@(t)cylinder_fun(g,t);
f=exp(delta_int(cylinder_g,t,s));
end

function f=cos_phi(t,s)
f=(exp_fun(@(x) phi(x).*1i,t,s)+exp_fun(@(x) -phi(x).*1i,t,s))./2;
end

```

```

function f=sin_phi(t,s)
f=(exp_fun(@(x) phi(x).*1i,t,s)-exp_fun(@(x) -phi(x).*1i,t,s))./2i;
end

function f=P_H(t,s)
f=(-mu(s).*phi(s).*cos_phi(t,s)+sin_phi(t,s))./(phi(sigma(s)).*...
(1+mu(s).^2.*phi(s).^2));
end

function f=Q_H(t,s)
f=(mu(s).*phi(s).*phi(t).*sin_phi(t,s)+phi(t).*cos_phi(t,s))./(...
phi(sigma(s)).*(1+mu(s).^2.*phi(s).^2));
end

function f=phi_diff(t)
if mu(t)==0
f=q_diff(t)/(2*sqrt(q(t)));
else
f=(phi(sigma(t))-phi(t))/mu(t);
end
end

%need function q_diff(t)
function f=h_H(t)
f=-p(t)-phi_diff(t)/phi(t);
end

function funcn=funvec(n,m)
global time_scale;
t_0=time_scale(1);
T=time_scale(end)-time_scale(1);
if n==1
funcn= (1/phi(t_0)*cos_phi(m(n),t_0)*Q_H(t_0+T,m(1))...
-sin_phi(m(n),t_0)*P_H(t_0+T,m(1)))*phi(m(n))*h_H(m(1));
else
last=1;
for k=2:n
last=last*Q_H(m(k-1),m(k))*h_H(m(k));
end
funcn=last*(1/phi(t_0)*cos_phi(m(n),t_0)*Q_H(t_0+T,m(1))...
-sin_phi(m(n),t_0)*P_H(t_0+T,m(1)))*phi(m(n))*h_H(m(1));
end
end

function f=Delta(n)
global time_scale;
m=time_scale;
m(end)=[];
m=sort(m,'descend');

```

```

M=nchoosek(m,n);
[r,~]=size(M);
sum=0;
for i=1:r
prod=1;
for j=1:n
prod=prod*mu(M(i,j));
end
sum=sum+prod*funvec(n,M(i,1:n));
end
f=sum;
end

function f=valueOfDelta()
global time_scale;
t_0=time_scale(1);
T=time_scale(end)-time_scale(1);
sum=(1+phi(t_0+T)/phi(t_0))*cos_phi(t_0+T,t_0);
for i=1:(length(time_scale)-1)
sum=sum+Delta(i);
end
f=sum;
end

function f = nIntergrate(fun,n)
global time_scale;
t0=time_scale(1);N=n;
up=cell(1,N);
up{1}='time_scale(end)';
for i=2:N
up{i}=['t',num2str(i-1)];
end
expr = GenerateExpr_quadl(N);
function expr = GenerateExpr_quadl(n)
if n == 1
expr = ['delta_int(@(t',num2str(N),')',fun,',',up{N},',t0)'];
else
expr = ['delta_int(@(t',num2str(N-n+1),')',...
GenerateExpr_quadl(n-1),',',up{N-n+1},',t0)'];
end
end
f = eval(expr);
end

function f=func_ser(n)
last=['(cos_phi(t',num2str(n),',t0)*Q_H(time_scale(end),...
t1)/phi(t0)', '-sin_phi(t',num2str(n),',t0)*P_H(time_scale(end),...
t1))*phi(t',num2str(n),')*h_H(t1)'];
if n==1

```

```

f=last;
else
for i=2:n
last=[last,'*Q_H(t',num2str(i-1),'t',num2str(i),'*h_H(t',...
num2str(i),'')'];
end
f=last;
end
end

function f=Delta_H(n)
global time_scale;
t0=time_scale(1);
sum=(1+phi(time_scale(end))/phi(t0))*cos_phi(time_scale(end),t0);
for i=1:n
sum=sum+nIntergrate(func_ser(i),i);
end
f=sum;
end
function f=ConPhi(t,s)
f=integral(@(x) arrayfun(@(x) sqrt(q(x))+0*x,x),s,t);
end
function f=Conh(t)
f=-p(t)-0.5*q_diff(t)/q(t);
end
function f=Confun_sec(n)
temp='ConPhi(time_scale(end),time_scale(1))';
temp2='1';
for i=1:2:n-1
temp=[temp,'-2*ConPhi(x',num2str(i),'x',num2str(i+1),'')'];
end
for j=1:n
temp2=[temp2,'*Conh(x',num2str(j),'')'];
end
f=['cos(',temp,')','* ',temp2];
end
function f=Conint_fun_sec(n)
global B;
if B==1
if mod(n,2)==0
f=ConnIntergrate(Confun_sec(n),n)/(2^(n-1));
else
f=0;
end
else
f=ConnIntergrate(Confun_sec(n),n)/(2^(n-1));
end
end
end

```

```

function f=Consum(n)
global time_scale;
sum=2*cos(ConPhi(time_scale(end),time_scale(1)));
for i=1:n
sum=sum+Conint_fun_sec(i);
end
f=sum;
end
function f = ConnIntergrate(fun,N)
global time_scale;
t0=time_scale(1);
up=cell(N);low=cell(N);x0=time_scale(end);
for i=1:N
low{i}=['t0+0*x',num2str(i-1)];
up{i}=['x',num2str(i-1)];
end

if mod(N,2) == 0
expr = GenerateExpr_quad2d(N);
else
expr = ['quad1(@(x1) arrayfun(@(x1)',GenerateExpr_quad2d(N-1),...
',x1)',',low{1}',',',up{1}',')'];
end
function expr = GenerateExpr_quad2d(n)
if n == 2
expr = ['quad2d(@(x',num2str(N-1)',',x',num2str(N),')',...
'arrayfun(@(x',num2str(N-1)',',x',num2str(N),')',fun,...
',x',num2str(N-1)',',x',num2str(N),')',',low{N-1}',',',...
up{N-1}',',@(x',num2str(N-1),')',low{N}',',@(x',...
num2str(N-1),')',up{N},')'];
else
expr = ['quad2d(@(x',num2str(N-n+1)',',x',num2str(N-n+2),')',...
'arrayfun(@(x',num2str(N-n+1)',',x',num2str(N-n+2),')',...
GenerateExpr_quad2d(n-2)',',x',num2str(N-n+1)',',x',...
num2str(N-n+2),')',',low{N-n+1}',',',up{N-n+1}',',@(x',...
num2str(N-n+1),')',low{N-n+2}',',@(x',num2str(N-n+1),')',...
up{N-n+2},')'];
end
end
f = eval(expr);
end

```

6. Examples

Example 6.1 (Discrete Time Scale). Consider the time scale $\mathbb{T} = \mathbb{Z}$ and the regressive equation

$$\Delta\Delta x(t) + \frac{-17 + 15(-1)^t}{16}\Delta x(t) + \frac{1 - 15(-1)^t}{16}x(t) = 0, \quad (6.1)$$

which can be rewritten as

$$\Delta X(t) = \begin{pmatrix} 0 & 1 \\ -\frac{1-15(-1)^t}{16} & -\frac{-17+15(-1)^t}{16} \end{pmatrix} X(t). \quad (6.2)$$

$$\text{Let } A(t) = \begin{pmatrix} 0 & 1 \\ -q(t) & -p(t) \end{pmatrix} = \begin{pmatrix} 0 & 1 \\ -\frac{1-15(-1)^t}{16} & -\frac{-17+15(-1)^t}{16} \end{pmatrix}.$$

Obviously, the time scale $\mathbb{Z}$ and matrix $A(t)$ have periods of 2. Also, it can be verified that $\mathcal{B} = e_{-p+\mu q}(2, 0) = 1$ and then we are going to use formula (4.29) to calculate the value of $\mathcal{A}$. Taking

$$\phi(0) = 1, \phi(1) = -\frac{7}{8}, \phi(2) = -\frac{8}{7},$$

then we have

$$\begin{aligned} \cos_\phi(0) &= 1, \quad \sin_\phi(0) = 0, & \cos_\phi(1) &= 1, \quad \sin_\phi(1) = 1, \\ \cos_\phi(2) &= \frac{15}{8}, \quad \sin_\phi(2) = \frac{1}{8}, & \cos_\phi(2, 1) &= 1, \quad \sin_\phi(2, 1) = -\frac{7}{8}, \\ P(1, 0) &= 0, \quad Q(1, 0) = 1, & P(2, 0) &= 1, \quad Q(2, 0) = \frac{64}{49}, \\ P(2, 1) &= 0, \quad Q(2, 1) = 1, & h(0) &= 2, \quad h(1) = \frac{83}{49}, \end{aligned}$$

and

$$\begin{aligned} \mathcal{A} &= \left(1 + \frac{\phi(t_0 + T)}{\phi(t_0)}\right) \cos_\phi(t_0 + T) \\ &\quad + \int_0^2 (\cos_\phi(t_1)Q(2, t_1) - \sin_\phi(t_1)P(2, t_1)) \cdot \phi(t_1)h(t_1)\Delta t_1 \\ &\quad + \int_0^2 \int_0^{t_1} (\cos_\phi(t_2)Q(2, t_1) - \sin_\phi(t_2)P(2, t_1)) \cdot \phi(t_2)Q(t_1, t_2)h(t_1)h(t_2)\Delta t_2\Delta t_1 \\ &= -\frac{15}{56} + \frac{128}{49} - \frac{7}{8} \cdot \frac{83}{49} + \frac{166}{49} = \frac{17}{4}. \end{aligned}$$

Now we calculate the value of $\mathcal{A}$ using (3.4). It can be seen that the transition matrix of system (6.2) is given by

$$\begin{aligned} &\Phi_A(t, 0) \\ &= \begin{pmatrix} 2^t - 2^t \int_0^t \frac{5+3(-1)^s}{2^{2s+3}} \Delta s & 2^t \int_0^t \frac{5+3(-1)^s}{2^{2s+3}} \Delta s \\ 2^t - 2^t \int_0^t \frac{5+3(-1)^s}{2^{2s+3}} \Delta s - \frac{5+3(-1)^t}{2^{t+3}} & 2^t \int_0^t \frac{5+3(-1)^s}{2^{2s+3}} \Delta s + \frac{5+3(-1)^t}{2^{t+3}} \end{pmatrix}. \end{aligned} \quad (6.3)$$

Then we can obtain that $\mathcal{A} = \text{trace}(\Phi_A(2, 0)) = \frac{17}{4}$, which is consistent with the previous calculations, and we get system (6.1) is unstable. We also can use Program 1 given in Section 5 to calculate:

```
Enter the discrete point: [0,1,2]
Enter the continuous interval: []
The value of A is 4.250000
The value of B is 1.000000
The modulus of multipliers are 0.250000 4.000000.
```

Example 6.2 (Discrete Time Scale). Consider the time scale $\mathbb{T} = 2\mathbb{Z}$ and the regressive equation

$$x^{\Delta\Delta}(t) + \frac{\sin \frac{\pi}{3}t + 2}{10}x^{\Delta}(t) + \frac{\sin \frac{\pi}{3}t + 2}{20}x(t) = 0. \quad (6.4)$$

Obviously, the time scale $2\mathbb{Z}$ and the functions $p(t), q(t)$ have periods of 6. Also, it can be verified that $\mathcal{B} = e_{-p+\mu q}(6, 0) = 1$. Then we use Program 1 to calculate:

```
Enter the discrete point: [0,2,4,6]
Enter the continuous interval: []
The value of A is -0.752000
The value of B is 1.000000
The modulus of multipliers are 1.000000 1.000000.
```

Now we calculate the value of $\mathcal{A}$ using (3.4). Let $x_1(t), x_2(t)$ be solutions of (6.4) satisfying

$$x_1(0) = 1, \quad x_1^{\Delta}(0) = 0, \quad x_2(0) = 0, \quad x_2^{\Delta}(0) = 1.$$

Then we have

$$\begin{aligned} x_1^{\Delta\Delta}(0) &= -\frac{1}{10}, & x_1(2) &= 1, & x_1^{\Delta}(2) &= -\frac{1}{5}, \\ x_1^{\Delta\Delta}(2) &= -\frac{3\sqrt{3}-12}{200}, & x_1(4) &= \frac{3}{5}, & x_1^{\Delta}(4) &= -\frac{3\sqrt{3}+32}{100}, \\ x_1(6) &= -\frac{3\sqrt{3}+2}{50}, & x_2^{\Delta\Delta}(0) &= -\frac{1}{5}, & x_2(2) &= 2, \\ x_2^{\Delta}(2) &= \frac{3}{5}, & x_2^{\Delta\Delta}(2) &= -\frac{2\sqrt{3}+8}{25}, & x_2(4) &= \frac{16}{5}, \\ x_2^{\Delta}(4) &= -\frac{4\sqrt{3}+1}{25}, & x_2^{\Delta\Delta}(4) &= \frac{55\sqrt{3}-168}{500}, & x_2^{\Delta}(6) &= \frac{15\sqrt{3}-178}{250}. \end{aligned}$$

Thus, $\mathcal{A} = x_1(6) + x_2^{\Delta}(6) = -0.752$, which is consistent with the previous calculations and we get system (6.4) is stable.

Example 6.3 (Hybrid Time Scale). Consider the time scale $\mathbb{T} = [2k\pi, (2k+1)\pi], k \in \mathbb{Z}$ and the regressive equation

$$x^{\Delta\Delta}(t) + p(t)x^{\Delta}(t) + x(t) = 0, \quad (6.5)$$

where

$$p(t) = \begin{cases} 0, & t \in [2k\pi, (2k+1)\pi), \\ \frac{1}{4}, & t = (2k+1)\pi. \end{cases}$$

Obviously, $q(t) = 1$ and the time scale $\mathbb{T}$ and the function $p(t)$ have periods of 2π . Also, it can be verified that $\mathcal{B} = e_{-p+\mu q}(2\pi, 0) = \pi^2 - \frac{\pi}{4} + 1$ and then we are going to use formula (4.29) to calculate the value of $\mathcal{A}$. It can be seen that $\phi(t) = 1$ for all $t \in \mathbb{T}$ and

$$h(t) = -p(t) - \frac{\phi^\Delta(t)}{\phi(t)} = \begin{cases} 0, & t \in [2k\pi, (2k+1)\pi), \\ -\frac{1}{4}, & t = (2k+1)\pi. \end{cases}$$

Note that $h(t) = 0$ for all $t \in [0, \pi)$, then the expression of $\mathcal{A}$ given by (4.29) can be reduced to

$$\begin{aligned} \mathcal{A} &= 2\cos_1(2\pi) + \int_{\pi}^{2\pi} (\cos_1(t_1)Q(2\pi, t_1) - \sin_1(t_1)P(2\pi, t_1)) \cdot h(t_1) \Delta t_1 \\ &= 2\cos_1(2\pi) + \mu(\pi) \cdot (\cos_1(\pi)Q(2\pi, \pi) - \sin_1(\pi)P(2\pi, \pi)) \cdot h(\pi) \\ &= -2 + \pi \cdot (-1 - 0) \cdot (-\frac{1}{4}) = \frac{\pi}{4} - 2. \end{aligned} \quad (6.6)$$

Now we calculate the value of $\mathcal{A}$ using (3.4). Let $x_1(t), x_2(t)$ be solutions of (6.5) satisfying

$$x_1(0) = 1, \quad x_1^\Delta(0) = 0, \quad x_2(0) = 0, \quad x_2^\Delta(0) = 1.$$

For any $t \in [0, \pi]$, we have

$$x_1(t) = \cos t \quad \text{and} \quad x_2(t) = \sin t.$$

Hence, we get $x_1^\Delta(\pi) = 0, x_2^\Delta(\pi) = -1$ and

$$x_2^{\Delta\Delta}(\pi) = \frac{x_2^\Delta(2\pi) - x_2^\Delta(\pi)}{\pi} = -p(\pi)x_2^\Delta(\pi) - x_2(\pi).$$

Thus, $x_1(2\pi) = -1, x_2^\Delta(2\pi) = \frac{\pi}{4} - 1$. Finally, we have

$$\mathcal{A} = x_1(2\pi) + x_2^\Delta(2\pi) = \frac{\pi}{4} - 2,$$

which is consistent with the previous calculations and system (6.5) is unstable. We also can use Program 1 given in Section 5 to calculate $\mathcal{A}(n)$ given by (4.25):

```
Enter the discrete point: [2*pi]
Enter the continuous interval: [0;pi]
n:1
The value of A(1) is -1.214602
The value of B is 10.084206
The 1th approximate modulus are 3.175564 3.175564.
```

Example 6.4 (Continuous Time Scale). Consider the time scale $\mathbb{T} = \mathbb{R}$ and the equation

$$x'(t) + \frac{1}{2}\sin(2t)x'(t) + \frac{1}{4}x(t) = 0. \quad (6.7)$$

We can use Program 1 to calculate $\mathcal{A}(n)$:

```

Enter the discrete point: []
Enter the continuous interval: [0;pi]
n:3
The value of A(3) is -0.065450
The value of B is 1.000000
The 3th approximate modulus are 1.000000 1.000000.

```

Now we estimate the value of $|\mathcal{A}(3) - \mathcal{A}|$ by (4.28). A straightforward calculation leads to

$$|\mathcal{A}(3) - \mathcal{A}| \leq e^{\frac{\pi}{2}} - \left(1 + \frac{\pi}{2} + \frac{(\frac{\pi}{2})^2}{2} + \frac{(\frac{\pi}{2})^3}{6}\right) \approx 0.360016406528039.$$

It is clear that $0 < \mathcal{A} < 0.5$. It follows from Theorem 3.2 that system (6.7) is stable.

Example 6.5 ([49]). Consider Mathieu equation

$$x'' + (\lambda - h \cos 2t)x = 0. \quad (6.8)$$

Table 1. The values of parameters h and λ

h	λ			
	λ_1	λ_2	λ'_1	λ'_2
1	3.979	4.101	9.014	9.018
2	3.917	4.371	9.047	9.078
3	3.814	4.747	9.093	9.193

Table 2. 3-th approximate value of $\mathcal{A}$

Equation	3-th approximate value of $\mathcal{A}$
$x'' + (3.979 - \cos 2t)x = 0$	2.000049
$x'' + (4.101 - \cos 2t)x = 0$	2.000044
$x'' + (9.014 - \cos 2t)x = 0$	-2.000001
$x'' + (9.018 - \cos 2t)x = 0$	-2.000000
$x'' + (3.917 - 2 \cos 2t)x = 0$	2.000798
$x'' + (4.371 - 2 \cos 2t)x = 0$	2.000384
$x'' + (9.047 - 2 \cos 2t)x = 0$	-2.000009
$x'' + (9.078 - 2 \cos 2t)x = 0$	-2.000018
$x'' + (3.814 - 3 \cos 2t)x = 0$	1.998646
$x'' + (4.747 - 3 \cos 2t)x = 0$	1.998733
$x'' + (9.093 - 3 \cos 2t)x = 0$	-2.000103
$x'' + (9.193 - 3 \cos 2t)x = 0$	-2.000093

Let $\mathcal{A}[\lambda_i]$ and $\mathcal{A}[\lambda'_i]$ be the value of $\mathcal{A}$ of (6.8) as $\lambda = \lambda_i$ and $\lambda = \lambda'_i$, respectively (the values of parameters h , λ_i and λ'_i given by Table 1). It is well known that $\mathcal{A}[\lambda_i] \approx 2$ and $\mathcal{A}[\lambda'_i] \approx -2$, (see [49]). Now we are going to calculate the 3-th approximate value of $\mathcal{A}[\lambda_i]$ and $\mathcal{A}[\lambda'_i]$ by Program 1 and the results are shown in Table 2.

References

- [1] R. P. Agarwal and M. Bohner, *Basic calculus on time scales and some of its applications*, Results Math., 1999, 35(1–2), 3–22.
- [2] R. P. Agarwal, M. Bohner, A. Domoshnitsky and Y. Goltser, *Floquet theory and stability of nonlinear integro-differential equations*, Acta Math. Hungar., 2005, 109(4), 305–330.
- [3] R. P. Agarwal, M. Bohner and D. O'Regan, *Dynamic equations on time scales: a survey*, J. Comput Appl. Math., 2002, 141(1–2), 1–26.
- [4] M. Adivar and H. C. Koyuncuoğlu, *Floquet theory based on new periodicity concept for hybrid systems involving q -difference equations*, Appl. Math. Comput., 2016, 273, 1208–1233.
- [5] C. D. Ahlbrandt and J. Ridenhour, *Floquet theory for time scales and Putzer representations of matrix logarithms*, J. Difference Equ. Appl., 2003, 9(1), 77–92.
- [6] B. Aulbach and C. Pötzsche, *Reducibility of linear dynamic equations on measure chains*, J. Comput. Appl. Math., 2002, 141(1–2), 101–115.
- [7] S. R. Barone, M. A. Narcowich and F. J. Narcowich, *Floquet theory and applications*, Phys. Rev. A, 1977, 15(3), 1119–1125.
- [8] L. C. Becker, T. A. Burton and T. Krisztin, *Floquet theory for a Volterra equation*, J. London Math. Soc., 1988, 37(2), 141–147.
- [9] R. Benterki and J. Llibre, *Periodic solutions of a class of duffing differential equations*, J. Nonlinear Modeling Anal., 2019, 1(2), 167–177.
- [10] R. Benterki and J. Llibre, *Periodic solutions of the duffing differential equation revisited via the averaging theory*, J. Nonlinear Modeling Anal., 2019, 1(1), 11–26.
- [11] M. Bohner, G. Guseinov and A. Peterson, *Introduction to the Time Scales Calculus: Advances in Dynamic Equations on Time Scales*, Birkhäuser, Boston, 2003.
- [12] M. Bohner and A. Peterson, *Dynamic Equations on Time Scales: An Introduction with Applications*, Birkhäuser, Boston, 2001.
- [13] D. Breda, S. Mast and R. Vermiglio, *Numerical computation of characteristic multipliers for linear time periodic coefficients delay differential equations*, IFIC Proceedings Volumes, 2006, 39, 163–168.
- [14] H. Broer, *Resonance tongues in Hill's equations: a geometric approach*, J. Differential Equations, 2000, 166(2), 290–327.
- [15] R. Carlson, *Eigenvalue estimates and trace for the matrix Hill's equation*, J. Differential Equations, 2000, 167(1), 211–244.
- [16] D. Cheng, K. I. Kou and Y. Xia, *A unified annlysis of linear quaternion dynamic equations on time scales*, J. Appl. Anal. Comput., 2020, 10(5), 1869–1877.
- [17] C. Chicone, *Ordinary Differential Equations with Applications*, Springer-Verlag, New York, 1999.
- [18] S. N. Chow, K. Lu and J. Mallet-Paret, *Floquet theory for parabolic differential equations*, J. Differential Equations, 1994, 109(1), 147–200.

- [19] S. N. Chow and H. O. Walthers, *Characteristic multipliers and stability of symmetric periodic solutions of $\dot{x}(t) = g(x(t-1))$* , Trans. Amer. Math. Soc., 1988, 307(1), 127–142.
- [20] J. J. DaCunha, *Lyapunov Stability and Floquet Theory for Nonautonomous Linear Dynamic Systems on Time Scales*, Ph.D. Thesis, Baylor University, 2004. ISBN: 978-0496-82618-6.
- [21] J. J. DaCunha and J. M. Davis, *A unified Floquet theory for discrete, continuous, and hybrid periodic linear systems*, J. Differential Equations, 2011, 251(11), 2987–3027.
- [22] A. Demir, *Floquet theory and non-linear perturbation analysis for oscillators with differential-algebraic equations*, Int. J. Circuit Theory Appl., 2000, 28(2), 163–185.
- [23] T. S. Doan, A. Kalauch, S. Siegmund and F. R. Wirth, *Stability radii for positive linear time-invariant systems on time scales*, Systems Control Lett., 2010, 59(3–4), 173–179.
- [24] T. S. Doan, A. Kalauch and S. Siegmund, *Exponential stability of linear time-invariant systems on time scales*, Nonlinear Dyn. Syst. Theory, 2009, 9(1), 37–50.
- [25] P. Dormayer, A. F. Ivanov and B. Lani-Wayda, *Floquet multipliers of symmetric rapidly oscillating solutions of differential delay equations*, Tohoku Math. J., 2002, 54(3), 419–441.
- [26] L. Erbe and A. Peterson, *Green's functions and comparison theorems for differential equations on measure chains*, Dynam. Contin. Discrete Impuls. Systems, 1999, 6(1), 121–137.
- [27] M. Federson, J. G. Mesquita and A. Slavík, *Measure functional differential equations and functional dynamic equations on time scales*, J. Differential Equations, 2012, 252(6), 3816–3847.
- [28] M. Federson, R. Grau, J. G. Mesquita and E. Toon, *Lyapunov stability for measure differential equations and dynamic equations on time scales*, J. Differential Equations, 2019, 267(7), 4192–4223.
- [29] H. I. Freedman, *Almost Floquet systems*, J. Differential Equations, 1971, 10, 345–354.
- [30] F. Gesztesy and R. Weikard, *Floquet theory revisited*, in: Differential Equations and Mathematical Physics, Proceedings of the International Conference, Univ. of Alabama at Birmingham, March 13–17, 1994, International Press, Boston, MA, 1995.
- [31] S. Hilger, *Ein Maßkettenkalkül mit Anwendung auf Zentrumsmannigfaltigkeiten*, Ph.D. thesis, Universität Würzburg, 1988.
- [32] S. Hilger, *Analysis on measure chains—a unified approach to continuous and discrete calculus*, Result Math., 1990, 18(1–2), 18–56.
- [33] S. Hilger, *Differential and difference calculus-unified!*, Nonlinear Anal., 1997, 30(5), 2683–2694.
- [34] R. A. Johnson, *On a Floquet theory for almost-periodic, two-dimensional linear systems*, J. Differential Equations, 1980, 37(2), 184–205.

- [35] W. G. Kelley and A. C. Peterson, *Difference Equations: An Introduction with Applications*, Academic Press, San Diego, 2001.
- [36] D. Kotsis, *The approximation of the characteristic multipliers of periodic differential equations*, Alkalmaz. Mat. Lapok, 1976, 2(3–4), 269–276.
- [37] P. Kuchment, *On the behavior of Floquet exponents of a kind of periodic evolution problems*, J. Differential Equations, 1994, 109(2), 309–324.
- [38] R. Lamour, R. März and R. Winkler, *How Floquet theory applies to index 1 differential algebraic equations*, J. Math. Anal. Appl., 1998, 217(2), 372–394.
- [39] Z. Li, C. Wang, R. P. Agarwal and D. O'Regan, *Commutativity of quaternion-matrix-valued functions and quaternion matrix dynamic equations on time scales*, Stud. Appl. Math., 2021, 146(1), 139–210.
- [40] Y. Li, X. Wang and N. Huo, *Weyl almost automorphic solutions in distribution sense of Clifford-valued stochastic neural networks with time-varying delays*, Proc. A, 2022, 478(2257), 20210719.
- [41] T. Luzyanina and K. Engelborghs, *Computing Floquet multipliers for functional differential equations*, Internat. J. Bifur. Chaos Appl. Sci. Engrg., 2002, 12(12), 2977–2989.
- [42] J. Mallet-Paret and G. R. Sell, *Systems of differential delay equations: Floquet multipliers and discrete Lyapunov functions*, J. Differential Equations, 1996, 125(2), 385–440.
- [43] V. G. Papanicolaou and D. Kravvaritis, *The Floquet theory of the periodic Euler-Bernoulli equation*, J. Differential Equations, 1998, 150(1), 24–41.
- [44] C. Pötzsche, *Langsame Faserbündel dynamischer Gleichungen auf Maßketten*, Logos, Berlin, 2002.
- [45] C. Pötzsche, *Exponential dichotomies for dynamic equations on measure chains*, Nonlinear Anal., 2001, 47(2), 873–884.
- [46] C. Pötzsche, *Topological decoupling, linearization and perturbation on inhomogeneous time scales*, J. Differential Equation, 2008, 245(5), 1210–1242.
- [47] C. Pötzsche, S. Siegmund and F. Wirth, *A spectral characterization of exponential stability for linear time-invariant systems on time scales*, Discrete Contin. Dyn. Syst., 2003, 9(5), 1223–1241.
- [48] J. Shi, *On stability of two order linear differential equation with periodic coefficient*, Acta Math. Sci. Ser. A, 2000 20(1), 130–139.
- [49] J. Shi, M. Lin and J. Chen, *The calculation for characteristic multiplier of Hill's equation*, Appl. Math. Comput., 2004, 159(1), 57–77.
- [50] S. Siegmund, *A spectral notion for dynamic equations on time scales*, J. Comput. Appl. Math., 2002, 141(1–2), 255–265.
- [51] C. Simmendinger, A. Wunderlin and A. Pelster, *Analytical approach for the Floquet theory of delay differential equations*, Phys. Rev. E, 1999, 59(5), 5344–5353.
- [52] A. L. Skubachevskii and H. O. Walther, *On the Floquet multipliers of periodic solutions to non-linear functional differential equations*, J. Dynam. Differential Equations, 2006 18(2), 257–355.

- [53] Y. V. Teplinskiĭ and A. Y. Teplinskiĭ, *On the Erugin and Floquet-Lyapunov theorems for countable systems of difference equations*, Ukrainian Math. J., 1996, 48(2), 314–321.
- [54] K. O. Val'ter and A. L. Skubachevskii, *On Floquet multipliers for slowly oscillating periodic solutions of nonlinear functional differential equations*, Tr. Mosk. Mat. Obs., 2003, 64, 3–53.
- [55] H. O. Walther, *Hyperbolic periodic solutions, heteroclinic connections and transversal homoclinic points in autonomous differential delay equations*, Mem. Amer. Math. Soc., 1989, 79(402). DOI: 10.1090/memo/0402.
- [56] C. Wang, R. P. Agarwal and D. O'Regan, *Calculus of fuzzy vector-valued functions and almost periodic fuzzy vector-valued functions on time scales*, Fuzzy Sets and Systems, 2019, 375, 1–52.
- [57] C. Wang, R. P. Agarwal and D. O'Regan, *Almost periodic fuzzy multidimensional dynamic systems and applications on time scales*, Chaos Solitons Fractals, 2022, 156, 111781.
- [58] C. Wang, R. P. Agarwal, D. O'Regan and R. Sakthivel, *Theory of Translation Closedness for Time Scales*, Springer-Verlag, Switzerland, 2020.
- [59] C. Wang and R. P. Agarwal, *Almost automorphic functions on semigroups induced by complete-closed time scales and application to dynamic equations*, Discrete Contin. Dyn. Syst. Ser. B, 2020, 25(2), 781–798.
- [60] C. Wang, Z. Li and R. P. Agarwal, *Hyers-Ulam-Rassias stability of high-dimensional quaternion impulsive fuzzy dynamic equations on time scales*, Discrete Contin. Dyn. Syst. Ser. S, 2022, 15(2), 359–386.
- [61] R. Weikard, *Floquet theory for linear differential equations with meromorphic solutions*, Electron. J. Qual. Theory Differ. Equ., 2000, 8, 1–6. DOI: 10.14232/ejqtde.2000.1.8.
- [62] J. Zhang, M. Fan and H. Zhu, *Existence and roughness of exponential dichotomies of linear dynamic equations on time scales*, Comput. Math. Appl., 2010, 59(8), 2658–2675.
- [63] J. Zhang, M. Fan and H. Zhu, *Necessary and sufficient criteria for the existence of exponential dichotomy on time scales*, Comput. Math. Appl., 2010, 60(8), 2387–2398.