PSEUDO ALMOST PERIODIC SOLUTION OF FRACTIONAL-ORDER CLIFFORD-VALUED HIGH-ORDER HOPFIELD NEURAL NETWORKS*

Nina Huo¹ and Yongkun Li^{2,†}

Abstract In this work, based on the principle of contraction mapping, we deduce sufficient conditions ensuring the existence of pseudo almost periodic solutions of fractional-order Clifford-valued high-order Hopfield neural networks (FCHHNNs). In addition, we employ a kind of Gronwall inequality to study the finite-time stability of pseudo almost periodic solutions of FCHHNNs. The results and methods of our paper are new. Finally, we give a numerical example to illustrate the effectiveness of the results obtained.

Keywords Fractional-order calculus, high-order Hopfield neural network, Clifford algebra, pseudo almost periodic solution, finite-time stability.

MSC(2010) 34K14, 34K37, 93D40.

1. Introduction

Clifford-valued neural networks (CVNNs) have greater advantages in high-dimensional signal processing and storage capacity because they require fewer connection weight functions compared to real-valued, complex-valued, and quaternion-valued neural networks [5–7, 13]. In recent years, more and more scholars have devoted themselves to the theoretical and practical application research of CVNN, and have achieved many results [2,3,19,22,24,26,32,33]. Since the multiplication of Clifford algebra does not satisfy the commutative law, many results have been obtained by decomposing CVNNs into real-valued neural networks [4,31–33]. However, the results obtained by the decomposition method are essentially about the real-valued system, which is not easy to be directly applied to the CVNN system under consideration. Therefore, it is of great theoretical significance and potential application value to further explore the direct method for studying the qualitative behavior of CVNNs, namely, the non decomposition method.

[†]The corresponding author.

¹Key Laboratory of Applied Mathematics and Mechanism of Artificial Intelligence, Hefei University, Hefei 230601, China

²Department of Mathematics, Yunnan University, Kunming 650091, China *The authors were supported by National Natural Science Foundation of China (Grant Nos. 12261098 and 11861072), Natural Science Foundation of Anhui Province (Grant No. 2108085QA10), Natural Science Research Project of Colleges and Universities in Anhui Province (Grant No. 2022AH051782) and Talent Research Fund Project of Hefei University (Grant No. 20RC22). Email: huonn@hfuu.edu.cn(N. Huo), yklie@ynu.edu.cn(Y. Li)

In addition, it is well known that fractional order differential equation models can better describe some real processes with heredity, memory and nonlocality than integer order differential equation models in many aspects. Therefore, in recent decades, researchers have proposed a large number of fractional differential equation models, and the application fields of fractional differential equations are also expanding. In particular, the study of fractional order neural networks has received extensive attention and achieved some good results [1,8,18,20,28–30,34–39,41–43]. However, the research results on fractional order CVNNs are few. Especially, up to now, no results of almost periodic oscillation of fractional-order CVNNs have been published, but nevertheless, as is well known, almost periodic oscillations are one of the crucial dynamics of neural networks [15, 16, 18, 20, 21, 26]. Therefore, the study of almost periodic oscillations of fractional-order CVNNs has important theoretical and practical implications.

Moreover, it is well known that high-order Hopfield neural networks (HHNNs) have advantages over their corresponding low-order ones in approximation, convergence speed, storage capacity and fault tolerance. Therefore, HHNNs have always been the focus of research [9,11,12,25]. In addition, time delay is ubiquitous and inevitable in practical systems. Consequently, it is more reasonable to consider the neural network systems with time delay.

Inspired by the above observations, and noted that in a certain sense, it is more practical to consider the finite-time stability than the stability in Lyapunov's sense. Therefore, the main purpose of this paper is to study the existence and finite-time stability of pseudo almost periodic solutions of FCHHNN with time-varying delays.

The contributions of this work are

- (1) The result obtained in this paper is the first result regarding the pseudo almost periodic solutions of fractional-order CVNN.
- (2) The approaches used in the paper can be applied to study almost periodic and almost automorphic solutions to other types of fractional-order CVNNs.
- (3) Even when the network considered in this paper degenerates into a real-valued one, the results of this paper still remain new.

The remaining part of the paper is structured in this way: in Sect. 2, we introduce some concepts, notations and preliminary results, as well as the description of the model. In Sect. 3, we discuss the existence and finite time stability of pseudo almost periodic solutions for the network under consideration. In Sect. 4, we present a numerical example and computer simulation. Finally, we draw a brief conclusion in Sect. 5.

2. Preliminaries and model description

Let $\mathcal{A} = \left\{ \sum_{A \in \mathcal{P}} a^A e_A, a^A \in \mathbb{R} \right\}$ be a real Clifford algebra over \mathbb{R}^m (see [14]), where $\mathcal{P} = \{\emptyset, 1, 2, \dots, A, \dots, 12 \cdots m\}$ and $e_A = e_{h_1} e_{h_2} \cdots e_{h_\nu}, A \in \mathcal{P}, e_{\emptyset} = e_0 = 1$ and $e_h, h = 1, 2, \dots, m$ are its generators and satisfy $e_i^2 = 1, i = 1, 2, \dots, s, e_i^2 = -1, i = s + 1, s + 2, \dots, m, e_i e_j + e_j e_i = 0, i \neq j$, where $i, j = 1, 2, \dots, m$.

For $x = \sum_{A \in \mathcal{P}} x^A e_A \in \mathcal{A}$, define $|x|_{\mathcal{A}} = \sqrt{\sum_{A \in \mathcal{P}} (x^A)^2}$ and for $y = (y_1, y_2, \dots, y_n) \in \mathcal{A}^n$, define $||y||_{\mathcal{A}^n} = \max_{1 \le p \le n} \{|y_p|_{\mathcal{A}}\}$. Obviously, $(\mathcal{A}, |\cdot|_{\mathcal{A}})$ and $(\mathcal{A}^n, |\cdot|_{\mathcal{A}^n})$ are

Banach spaces, respectively.

Definition 2.1. [27] The fractional integral of order α for a function $f \in L^1([a, b], \mathbb{R})$ is defined as

$${}_aI^{\alpha}_tf(t) = \frac{1}{\Gamma(\alpha)}\int_a^t (t-s)^{\alpha-1}f(s)ds, \ t\in[a,b],$$

where $\alpha > 0$ and Γ is the Gamma function, a and b may take $-\infty$ and $+\infty$ as their values.

Definition 2.2. [27] The Caputo derivative of order α for a function $f \in C^{n-1}([a,b],\mathbb{R})$ and $f^{(n)} \in L^1([a,b],\mathbb{R})$ is defined by

$$_{a}D_{t}^{\alpha}f(t)=\frac{1}{\Gamma(n-\alpha)}\int_{a}^{t}\frac{f^{n}(s)}{(t-s)^{\alpha+1-n}}ds,\ t\in[a,b],$$

where n is a nature number such that $n-1 < \alpha < n$ and Γ is the Gamma function, a and b may take $-\infty$ and $+\infty$ as their values.

Definition 2.3. The fractional integral with fractional order $\alpha > 0$ of function $f = \sum_{A \in \mathcal{P}} f^A e_A \in L^1([a, b]), \mathcal{A})$ is defined as ${}_aI^{\alpha}_t(t)f(t) = \sum_{A \in \mathcal{P}} {}_aI^{\alpha}_tf^A(t)e_A$ and the Caputo fractional-order derivative of order α for function $f \in C^{n-1}([a, b], \mathcal{A})$ and $f^{(n)} \in L^1([a, b], \mathcal{A})$ is defined as ${}_aD^{\alpha}_tf(t) = \sum_{A \in \mathcal{P}} {}_aD^{\alpha}_tf^A(t)e_A$, where n is a nature number such that $n - 1 < \alpha < n$.

Let $BC(\mathbb{R}, \mathcal{A}^l)$ be the set of bounded continuous functions from \mathbb{R} to \mathcal{A}^l , where l is a positive integer. Then $(BC(\mathbb{R}, \mathcal{A}^l), \|\cdot\|_{\infty})$ is a Banach space, where $\|f\|_{\infty} := \sup_{t \in \mathbb{R}} \|f(t)\|_{\mathcal{A}^l}$ for $f \in BC(\mathbb{R}, \mathcal{A}^l)$.

Definition 2.4. [10] Function $f \in BC(\mathbb{R}, \mathcal{A}^l)$ is called to be almost periodic if for every $\varepsilon > 0$, there is a positive number $l = l(\varepsilon)$ such that in each interval with length l, there is a τ satisfying

$$||f(t+\tau) - f(t)||_{\mathcal{A}^l} < \varepsilon \text{ for all } t \in \mathbb{R}.$$

The space of all such functions will be denoted by $AP(\mathbb{R}, \mathcal{A}^l)$.

Define

$$PAP_0(\mathbb{R},\mathcal{A}^l) := \left\{ f \in BC(\mathbb{R},\mathcal{A}^l) : \lim_{L \to \infty} \frac{1}{2L} \int_{-L}^{L} \|f(s)\|_{\mathcal{A}^l} ds = 0 \right\}.$$

Definition 2.5. [10] Let $f \in BC(\mathbb{R}, \mathcal{A}^l)$, then f is called pseudo almost periodic if there exist $f_1 \in AP(\mathbb{R}, \mathcal{A}^l)$ and $f_2 \in PAP_0(\mathbb{R}, \mathcal{A}^l)$ such that $f = f_1 + f_2$. The collection of all such functions will be denoted by $PAP(\mathbb{R}, \mathcal{A}^l)$.

Lemma 2.1. [10] The set $PAP(\mathbb{R}, \mathcal{A})$ with the supremum norm is a Banach space. Lemma 2.2. [10] Let $\alpha \in \mathbb{R}$, $f, g \in PAP(\mathbb{R}, \mathcal{A})$, then $\alpha f, f+g, f \cdot g \in PAP(\mathbb{R}, \mathcal{A})$. Lemma 2.3. Let $f \in PAP(\mathbb{R}, \mathcal{A}), \tau \in AP(\mathbb{R}, \mathbb{R}^+) \cap C^1(\mathbb{R}, \mathbb{R})$ with $\inf_{t \in \mathbb{R}} |1-\dot{\tau}(t)| > 0$, then $f(\cdot - \tau(\cdot)) \in PAP(\mathbb{R}, \mathcal{A})$. m

Proof. In view of Definition 2.5, f can be expressed as $f = f_1 + f_0$, in which $f_1 \in AP(\mathbb{R}, \mathcal{A}^n)$ and $f_0 \in PAP_0(\mathbb{R}, \mathcal{A}^n) \subset BC(\mathbb{R}, \mathcal{A}^n)$, then one has

$$f(t - \tau(t)) = f_1(t - \tau(t)) + f_0(t - \tau(t)).$$

Similar to the proof of Lemma 5 in [17], one can easily get $f_1(\cdot - \tau(\cdot)) \in AP(\mathbb{R}, \mathcal{A}^n)$. In addition, noting that

$$\begin{split} &\lim_{T \to +\infty} \frac{1}{2T} \int_{-T}^{T} \|f_0(t - \tau(t))\|_{\mathcal{A}^n} dt \\ &\leq \lim_{T \to +\infty} \frac{1}{\inf_{t \in R} |1 - \dot{\tau}(t)|} \frac{1}{2T} \left| \int_{-T - \tau(-T)}^{T - \tau(T)} \|f_0(s)\|_{\mathcal{A}^n} ds \right| \\ &= \lim_{T \to +\infty} \frac{1}{\inf_{t \in R} |1 - \dot{\tau}(t)|} \frac{1}{2T} \left| \left(\int_{-T - \tau(-T)}^{-T + \tau(T)} + \int_{-T + \tau(T)}^{T - \tau(T)} \right) \|f_0(s)\|_{\mathcal{A}^n} ds \right| \\ &= \lim_{T \to +\infty} \frac{1}{\inf_{t \in R} |1 - \dot{\tau}(t)|} \frac{T - \tau(T)}{T} \frac{1}{2(T - \tau(T))} \int_{-T + \tau(T)}^{T - \tau(T)} \|f_0(s)\|_{\mathcal{A}^n} ds \\ &= 0, \end{split}$$

we arrive at $f_0(\cdot - \tau(\cdot)) \in PAP_0(\mathbb{R}, \mathcal{A}^n)$. Thus, $f(\cdot - \tau(\cdot)) \in PAP(\mathbb{R}, \mathcal{A}^n)$. This completes the proof.

The model we are concerned in this paper is the following Caputo FCHHNN with time-varying delays:

$$t_0 D_t^{\alpha} x_p(t) = -a_p x_p(t) + \sum_{q=1}^n a_{pq}(t) f_q(x_q(t - \tau_{pq}(t))) + \sum_{q=1}^n \sum_{l=1}^n b_{pql}(t) g_q(x_q(t - \sigma_{pql}(t))) \times g_l(x_l(t - \nu_{pql}(t))) + I_p(t), \ t > t_0, \ p = 1, 2, \dots, n,$$
(2.1)

where $0 < \alpha < 1$ is a constant, n denotes the number of units in the network; $x_p(t) \in \mathcal{A}$ is the state of the pth unit at time t; $a_p \geq 0$ is the self feedback connection weight; $a_{pq}(t) \in \mathcal{A}$ and $b_{pql}(t) \in \mathcal{A}$ are connection weights of the network; $\tau_{pq}(t) \geq 0$, $\sigma_{pql}(t) \geq 0$ and $\nu_{pql}(t) \geq 0$ are transmission delays at time t; $I_p(t) \in \mathcal{A}$ represents the external inputs at time t; $f_q, g_q : \mathcal{A} \to \mathcal{A}$ are activation functions of signal transmission.

We will use the following notations:

$$\hat{\tau} = \max_{1 \le p,q \le n} \Big\{ \sup_{t \in \mathbb{R}} \tau_{pq}(t) \Big\}, \ \hat{\sigma} = \max_{1 \le p,q,l \le n} \Big\{ \sup_{t \in \mathbb{R}} \sigma_{pql}(t) \Big\}, \ \hat{\nu} = \max_{1 \le p,q,l \le n} \Big\{ \sup_{t \in \mathbb{R}} \nu_{pql}(t) \Big\},$$

$$\rho = \max\{\hat{\tau}, \hat{\sigma}, \hat{\nu}\}, \ \hat{a}_{pq} = \sup_{t \in \mathbb{R}} |a_{pq}(t)|_{\mathcal{A}}, \ \hat{b}_{pql} = \sup_{t \in \mathbb{R}} |b_{pql}(t)|_{\mathcal{A}}, \ \hat{I}_p = \sup_{t \in \mathbb{R}} |I_p(t)|_{\mathcal{A}}.$$

System (2.1) is supplemented with the initial values:

$$x_p(s) = \psi_p(s), \ s \in [t_0 - \rho, t_0], \ p = 1, 2, \dots, n,$$

where $\psi_p \in C([t_0 - \rho, t_0], \mathcal{A}).$

In the next section, in order to get our main results, we need the following conditions:

- $(S_1) \text{ For } p, q, l = 1, \dots, n, a_{pq}, b_{pql}, I_p \in PAP(\mathbb{R}, \mathcal{A}), \tau_{pq}, \sigma_{pql}, \nu_{pql} \in AP(\mathbb{R}, \mathbb{R}^+) \cap C^1(\mathbb{R}, \mathbb{R}) \text{ satisfying } \inf_{t \in \mathbb{R}} |1 \tau_{pq}(t)| > 0, \inf_{t \in \mathbb{R}} |1 \sigma_{pql}(t)| > 0, \inf_{t \in \mathbb{R}} |1 \nu_{pql}(t)| > 0.$
- (S₂) There exist positive numbers L_q^f, L_q^g, M_q^g such that, for all $x, y \in \mathcal{A}, q = 1, \ldots, n$,

$$|f_q(x) - f_q(y)|_{\mathcal{A}} \le L_q^f |x - y|_{\mathcal{A}}, \ |g_q(x) - g_q(y)|_{\mathcal{A}} \le L_q^g |x - y|_{\mathcal{A}}, \ |g_q(x)|_{\mathcal{A}} \le M_q^g,$$

 $f_q, h_q \in C(\mathcal{A}, \mathcal{A})$; in addition, $f_q(0) = g_q(0) = 0$. (S₃)

$$\max_{1 \le p \le n} \left\{ \frac{C_p}{a_p} + \frac{\hat{I}_p}{a_p} \right\} \le r, \quad \xi = \max_{1 \le p \le n} \left\{ \frac{D_p}{a_p} \right\} < 1,$$

where for p = 1, 2, ..., n,

$$C_{p} = \left(\sum_{q=1}^{n} \hat{a}_{pq} L_{q}^{f} + \sum_{q=1}^{n} \sum_{l=1}^{n} \hat{b}_{pql} L_{q}^{g} M_{l}^{g}\right) r,$$
$$D_{p} = \sum_{q=1}^{n} \hat{a}_{pq} L_{q}^{f} + \sum_{q=1}^{n} \sum_{l=1}^{n} \hat{b}_{pql} \left(L_{q}^{g} M_{l}^{g} + L_{l}^{g} M_{q}^{g}\right).$$

3. Main results

Based on Definition 2.3 and Clifford algebra's multiplication rule as well as Definition 3.1 in [23] of solutions for real-valued equations, we can introduce the following definition:

Definition 3.1. A function $x = (x_1, x_2, \dots, x_n)^T \in C([t_0 - \rho, +\infty), \mathcal{A}^n)$ is called a mild solution of system (2.1), if it meets

$$\begin{cases} x_p(t) = U_p(t-t_0)x_0 + \int_{t_0}^t (t-s)^{\alpha-1}\varphi_p(t-s)[\Theta_p(s,x) + I_p(s)]ds, \ t > t_0, \\ x_p(t) = \psi_p(t), \ t \in [t_0 - \rho, t_0], \end{cases}$$
(3.1)

where

$$\begin{split} \Theta_p(s,x) &= \sum_{q=1}^n \left[a_{pq}(s) f_q(x_q(s-\tau_{pq}(s))) \right. \\ &+ \sum_{l=1}^n b_{pql}(s) g_q(x_q(s-\sigma_{pql}(s))) g_l(x_l(s-\nu_{pql}(s))) \right], \\ x_p(t_0) &= \psi_p(t_0) = x_0, \ \varrho_\alpha(\gamma) = \frac{1}{\alpha} \gamma^{-1-\frac{1}{\alpha}} \vartheta_\alpha(\gamma^{\frac{-1}{\alpha}}), \\ U_p(t) &= \int_0^{+\infty} \varrho_\alpha(\gamma) e^{-a_p t^\alpha \gamma} d\gamma, \ \varphi_p(t) = \alpha \int_0^{+\infty} \gamma \varrho_\alpha(\gamma) e^{-a_p t^\alpha \gamma} d\gamma, \end{split}$$

$$\vartheta_{\alpha}(\gamma) = \frac{1}{\pi} \sum_{n=1}^{\infty} (-1)^{n-1} \gamma^{-n\alpha-1} \frac{\Gamma(n\alpha+1)}{n!} \sin(n\pi\alpha), \ \gamma \in (0, +\infty),$$

and ρ_{α} satisfies

$$\varrho_{\alpha}(\gamma) \ge 0, \quad \gamma \in (0, +\infty), \quad \int_{0}^{+\infty} \varrho_{\alpha}(\gamma) d\gamma = 1, \quad \int_{0}^{+\infty} \gamma \varrho_{\alpha}(\gamma) d\gamma = \frac{1}{\Gamma(\alpha+1)}.$$

Letting $t_0 \longrightarrow -\infty$, we gain

$$x_p(t) = \int_{-\infty}^t (t-s)^{\alpha-1} \varphi_p(t-s) [\Theta_p(s,x) + I_p(s)] ds$$

which is a solution of system (2.1).

Let $\mathbb{B} = \{x = (x_1, x_2, \dots, x_n)^T \in BC(\mathbb{R}, \mathcal{A}^n) : x_p \in PAP(\mathbb{R}, \mathcal{A}), p = 1, 2, ..., n_n\}$ \ldots, n , then $(\mathbb{B}, \|\cdot\|_{\infty})$ is a Banach space.

Theorem 3.1. Let (S_1) - (S_3) be fulfilled, then system (2.1) possesses unique one pseudo almost periodic mild solution in $\mathbb{B}_r = \{x \mid x \in \mathbb{B}, \|x\|_{\infty} \leq r\}.$

Proof. Define a mapping $\Phi : \mathbb{B} \to BC(\mathbb{R}, \mathcal{A}^n)$ as follow:

$$\Phi x = ((\Phi x)_1, (\Phi x)_2, \cdots, (\Phi x)_n)^T,$$

where for $p = 1, 2, \ldots, n, t \in \mathbb{R}$,

$$(\Phi x)_p(t) = \int_{-\infty}^t (t-s)^{\alpha-1} \varphi_p(t-s) (\Theta_p(s,x) + I_p(s)) ds.$$

Firstly, we will show that $\Phi : \mathbb{B} \to \mathbb{B}$. For every $x = (x_1, x_2, \dots, x_n)^T \in \mathbb{B}_r$, from (S_1) - (S_3) it follows that $f_q, g_q, \tau_{pq}, \sigma_{pql}, \nu_{pql}$ meet all the conditions of Lemma 2.3, and hence, by Lemmas 2.2 and 2.3, we can derive that

$$\phi_p(\cdot) = \Theta_p(\cdot, x) + I_p(\cdot) \in PAP(\mathbb{R}, \mathcal{A})$$

So, there are $\phi_p^1 \in AP(\mathbb{R}, \mathcal{A})$ and $\phi_p^0 \in PAP_0(\mathbb{R}, \mathcal{A})$ such that $\phi_p = \phi_p^1 + \phi_p^0$ where $p = 1, 2, \ldots, n$. Consequently, we have

$$(\Phi x)_{p}(t) = \int_{-\infty}^{t} (t-s)^{\alpha-1} \varphi_{p}(t-s) \phi_{p}(s) ds$$

= $\int_{-\infty}^{t} (t-s)^{\alpha-1} \varphi_{p}(t-s) \phi_{p}^{1}(s) ds + \int_{-\infty}^{t} (t-s)^{\alpha-1} \varphi_{p}(t-s) \phi_{p}^{0}(s) ds$
:= $(\Phi x)_{p}^{1}(t) + (\Phi x)_{p}^{0}(t), \ p = 1, 2, \dots, n.$ (3.2)

We will prove that $(\Phi x)_p^1 \in AP(\mathbb{R}, \mathcal{A})$ and $(\Phi x)_p^0 \in PAP_0(\mathbb{R}, \mathcal{A})$, for p = 1, 2, ..., n. Since $\phi_p^1 \in AP(\mathbb{R}, \mathcal{A})$, for given $\varepsilon > 0$, there corresponds an $l = l(\varepsilon) > 0$ such that every interval of length l contains a point $\zeta \in (a, a + l)$ such that

 $|\phi_n^1(t+\zeta) - \phi_n^1(t)|_{\mathcal{A}} < \varepsilon, \ t \in \mathbb{R}, \ p = 1, 2, \dots, n.$

From this and (3.2) it follows that

$$|(\Phi x)_{p}^{1}(t+\zeta) - (\Phi x)_{p}^{1}(t)|_{\mathcal{A}} = \left| \int_{-\infty}^{t} (t-s)^{\alpha-1} \varphi_{p}(t-s) \phi_{p}^{1}(s+\zeta) ds \right|^{2}$$

$$\begin{split} &-\int_{-\infty}^{t} (t-s)^{\alpha-1} \varphi_p(t-s) \phi_p^1(s) ds \Big|_{\mathcal{A}} \\ &\leq \sup_{t \in \mathbb{R}} \left| \int_{-\infty}^{t} (t-s)^{\alpha-1} \varphi_p(t-s) (\phi_p^1(s+\zeta) - \phi_p^1(s)) ds \right|_{\mathcal{A}} \\ &\leq \varepsilon \int_{-\infty}^{t} (t-s)^{\alpha-1} \alpha \int_{0}^{+\infty} \gamma \varrho_\alpha(\gamma) e^{-a_p(t-s)^{\alpha} \gamma} d\gamma ds \\ &\leq \varepsilon \int_{0}^{+\infty} \int_{0}^{+\infty} \alpha \gamma \varrho_\alpha \sigma^{\alpha-1} e^{-a_p \sigma^{\alpha} \gamma} d\sigma d\gamma \\ &= \frac{\varepsilon}{a_p}, \ t \in \mathbb{R}, \end{split}$$

which implies $(\Phi x)_p^1 \in AP(\mathbb{R}, \mathcal{A})$ for p = 1, 2, ..., n.

Noting that

$$\lim_{L \to \infty} \frac{1}{2L} \int_{-L}^{L} \left| \int_{-\infty}^{t} (t-s)^{\alpha-1} \varphi_p(t-s) \phi_p^0(s) ds \right|_{\mathcal{A}} dt$$
$$= \lim_{L \to \infty} \frac{1}{2L} \int_{-L}^{L} \left| \int_{0}^{\infty} \sigma^{\alpha-1} \varphi_p(\sigma) \phi_p^0(t-\sigma) d\sigma \right|_{\mathcal{A}} dt$$
$$\leq \frac{1}{a_p} \lim_{L \to \infty} \frac{1}{2L} \int_{-L}^{L} |\phi_p^0(t-\sigma)|_{\mathcal{A}} dt$$
$$= 0, \ p = 1, 2, \dots, n,$$

we have $(\Phi x)_p^0 \in PAP_0(\mathbb{R}, \mathcal{A})$. Hence, for p = 1, 2, ..., n, $(\Phi x)_p \in PAP(\mathbb{R}, \mathcal{A})$, this yields that $\Phi : \mathbb{B} \to \mathbb{B}$.

Next, we show that $\Phi : \mathbb{B}_r \to \mathbb{B}_r$.

For any $x \in \mathbb{B}_r$, we have

$$\begin{aligned} |\Theta_{p}(t,x)|_{\mathcal{A}} &\leq \sum_{q=1}^{n} \hat{a}_{pq} \left(|f_{q}(x_{q}(t-\tau_{pq}(t))) - f_{q}(0)|_{\mathcal{A}} + |f_{q}(0)|_{\mathcal{A}} \right) \\ &+ \sum_{q=1}^{n} \sum_{l=1}^{n} \hat{b}_{pql} (|g_{q}(x_{q}(t-\sigma_{pql}(t))) - g_{q}(0)|_{\mathcal{A}} + |g_{q}(0)|_{\mathcal{A}}) M_{l}^{g} \\ &\leq \left(\sum_{q=1}^{n} \hat{a}_{pq} L_{q}^{f} + \sum_{q=1}^{n} \sum_{l=1}^{n} \hat{b}_{pql} L_{q}^{g} M_{l}^{g} \right) ||x||_{\infty} \\ &\leq C_{p}, \ p = 1, 2, \dots, n, \end{aligned}$$
(3.3)

which combined with (3.3) and (S_3) leads to

$$\begin{split} |(\Phi x)_p(t)|_{\mathcal{A}} &\leq \int_{-\infty}^t (t-s)^{\alpha-1} \varphi_p(t-s) \left(C_p + \hat{I}_p\right) ds \\ &\leq & \frac{C_p}{a_p} + \frac{\hat{I}_p}{a_p} \end{split}$$

$$\leq r, p=1,2,\ldots,n.$$

Hence,

$$\|\Phi x\|_{\infty} = \sup_{t \in \mathbb{R}} \{\|(\Phi x)(t)\|_{\mathcal{A}^n}\} \le r,$$

that is, $\Phi x \in \mathbb{B}_r$. Therefore, $\Phi : \mathbb{B}_r \to \mathbb{B}_r$.

Finally, we prove that Φ is a contraction mapping. For any $x, y \in \mathbb{B}_r$, we find

$$\begin{split} &|(\Phi x)_{p}(t) - (\Phi y)_{p}(t)|_{\mathcal{A}} \\ \leq &\|x - y\|_{\infty} \int_{-\infty}^{t} (t - s)^{\alpha - 1} \varphi_{p}(t - s) \sum_{q=1}^{n} \left[\hat{a}_{pq} L_{q}^{f} + \sum_{l=1}^{n} \hat{b}_{pql} \left(L_{q}^{g} M_{l}^{g} + L_{l}^{g} M_{q}^{g} \right) \right] ds \\ \leq & \frac{D_{p}}{a_{p}} \|x - y\|_{\infty} \\ \leq & \xi \|x - y\|_{\infty}, \ p = 1, 2, \dots, n, \end{split}$$

which implies that Φ is a contraction.

Consequently, Φ possesses a unique fixed point in \mathbb{B}_r . The proof is completed.

Remark 3.1. From the proof process of Theorem 3.1, it is easy to see that the proof method of Theorem 3.1 is also applicable to studying the existence of pseudo almost automorphic solutions for system (2.1).

In the following, we take the initial moment $t_0 = 0$ to discuss the finite-time stability of system (2.1), that is, we consider system (2.1) supplemented with the following initial condition:

$$x_p(\theta) = \psi_p(\theta), \ \theta \in [-\rho, 0], \ p = 1, 2, \dots, n,$$

where $\psi_p \in C([-\rho, 0], \mathcal{A}).$

Definition 3.2. A mild solution x^* of system (2.1) with initial value ψ^* is called finite-time stable with respect to $\{\delta, \varepsilon, T\}$, here $0 < \delta < \varepsilon$ and T > 0, if every mild solution x of system (2.1) with initial value ψ satisfies that if $\|\psi - \psi^*\| \leq \delta$, then, for $t \in [0, T]$,

$$||x(t) - x^*(t)||_{\mathcal{A}^n} \le \varepsilon,$$

where $\|\psi - \psi^*\|_{\rho} = \sup_{t \in [-\rho, 0]} \|\psi(t) - \psi^*(t)\|_{\mathcal{A}^n}.$

Theorem 3.2. Assume (S_1) - (S_3) hold. If

$$\delta E_{\alpha} \left(M T^{\alpha} \right) \le \varepsilon, \tag{3.4}$$

where $M = \max_{1 \le p \le n} \{D_p\}$ and $E_{\alpha}(\cdot)$ is the Mittag-Leffler function of one parameter, then system (2.1) possesses exactly one pseudo almost periodic mild solution that is finite-time stable with respect to $\{\delta, \varepsilon, T\}$. **Proof.** Denote by x^* the pseudo almost periodic mild solution of system (2.1) with initial value ψ^* and let x be any mild solution to system (2.1) with initial value ψ . Set $z = x - x^*$, invoking (3.1), one gets

$$\begin{aligned} z_p(t) = U_p(t)(\psi_p(0) - \psi_p^*(0)) + \int_0^t (t-s)^{\alpha-1} \varphi_p(t-s) \bigg[\sum_{q=1}^n a_{pq}(s) \\ & \times \left(f_q(x_q(s - \tau_{pq}(s))) - f_q(x_q^*(s - \tau_{pq}(s))) \right) \\ & + \sum_{q=1}^n \sum_{l=1}^n b_{pql}(s) \Big(g_q(x_q(s - \sigma_{pql}(s))) \\ & \times g_l(x_l(s - \nu_{pql}(s))) - g_q(x_q^*(s - \sigma_{pql}(s))) g_l(x_l^*(s - \nu_{pql}(s))) \Big) \bigg] ds, \quad t \ge 0 \end{aligned}$$

and

$$\begin{aligned} |U_p(t)(\psi_p(0) - \psi_p^*(0))|_{\mathcal{A}} &\leq \int_0^{+\infty} \varrho_\alpha(\gamma) e^{-a_p^-(t)^{\alpha}\gamma} d\gamma |\psi_p(0) - \psi_p^*(0)|_{\mathcal{A}} \\ &\leq \int_0^{+\infty} \varrho_\alpha(\gamma) d\gamma |\psi_p - \psi_p^*|_{\mathcal{A}} \\ &\leq |\psi_p - \psi_p^*|_{\mathcal{A}}. \end{aligned}$$

Denote $\lambda(t) = \max_{1 \le p \le n} \sup_{s \in [-\rho, t]} |z_p(s)|_{\mathcal{A}}$, then for $t \in [0, T]$ and $p = 1, 2, \ldots, n$, we find

$$\begin{split} |z_{p}(t)|_{\mathcal{A}} \leq & \|\psi - \psi^{*}\|_{\rho} + \int_{0}^{t} (t-s)^{\alpha-1} \varphi_{p}(t-s) \bigg[\sum_{q=1}^{n} \hat{a}_{pq} L_{q}^{f} \\ & + \sum_{q=1}^{n} \sum_{l=1}^{n} \hat{b}_{pql} \Big(M_{l}^{g} L_{q}^{g} + M_{q}^{g} L_{l}^{g} \Big) \bigg] \lambda(s) ds \\ \leq & \|\psi - \psi^{*}\|_{\rho} + M \int_{0}^{t} (t-s)^{\alpha-1} \alpha \int_{0}^{+\infty} \gamma \varrho_{\alpha}(\gamma) e^{-a_{p}(t-s)^{\alpha} \gamma} d\gamma \lambda(s) ds \\ \leq & \|\psi - \psi^{*}\|_{\rho} + \frac{\alpha M}{\Gamma(\alpha+1)} \int_{0}^{t} (t-s)^{\alpha-1} \lambda(s) ds \\ \leq & \|\psi - \psi^{*}\|_{\rho} + \frac{M}{\Gamma(\alpha)} \int_{0}^{t} (t-s)^{\alpha-1} \lambda(s) ds, \end{split}$$

which combined with the fact that $|z_p(s)|_{\mathcal{A}} \leq ||\psi - \psi^*||_{\rho}$ for $s \in [-\rho, 0]$ yields

$$\lambda(t) \le \|\psi - \psi^*\|_{\rho} + \frac{M}{\Gamma(\alpha)} \int_0^t (t-s)^{\alpha-1} \lambda(s) ds,$$

for $t \in [0, T]$ and p = 1, 2, ..., n.

Further, invoking Corollary 2 in [40], we infer that

$$\lambda(t) \le \|\psi - \psi^*\|_{\rho} E_{\alpha}(Mt^{\alpha}), \ t \in [0, T].$$

Since the function $E_{\alpha}(\theta)$ is nondecreasing for $\theta \in \mathbb{R}^+$, we conclude that

$$|z_p(t)|_{\mathcal{A}} \le \lambda(t) \le \|\psi - \psi^*\|_{\rho} E_{\alpha}(MT^{\alpha}), \quad t \in [0, T].$$

By (3.4), for all $t \in [0, T]$, one has

$$\|x(t) - x^*(t)\| < \varepsilon,$$

which gives the conclusion of the theorem. The proof is complete.

4. An example

Our example is as follows.

Example 4.1. In system (2.1), take n = m = 2, and for p = 1, 2, let

$$\begin{split} x_p(t) &= e_0 x_p^0(t) + e_1 x_p^1(t) + e_2 x_p^2(t) + e_{12} x_p^{12}(t) \in \mathcal{A}, \ a_1 = 12, \ a_2 = 8, \\ a_{11}(t) &= a_{12}(t) = e_0 \frac{1}{80} \cos t + e_2 \frac{1}{80} \sin t, \\ a_{21}(t) &= a_{22}(t) = e_0 \frac{3}{400} \sin t + e_1 \frac{1}{80} \cos t + e_2 \frac{1}{80} \sin t + e_{12} \frac{3}{400} \cos t, \\ b_{111}(t) &= b_{112}(t) = b_{121}(t) = b_{122}(t) = e_0 \frac{1}{100} + e_2 \frac{7}{400} \sin t + e_{12} \frac{7}{400} \cos t, \\ b_{211}(t) &= b_{212}(t) = b_{221}(t) = b_{222}(t) = e_0 \frac{1}{80} + e_1 \frac{3}{200} \cos \sqrt{2}t + e_2 \frac{3}{200} \sin \sqrt{2}t, \\ f_p(x) &= e_0 0.5 \sin x^0 + e_1 0.5 \sin x^1 + e_1 0.5 \sin x^2 + e_{12} 0.5 \sin x^{12}, \\ g_p(x) &= e_0 \frac{1}{3} |\sin x^0| + e_1 \frac{1}{4} |\sin x^1| + e_2 \frac{1}{5} |\sin x^2| + e_{12} \frac{1}{6} |\sin x^{12}|, \\ \tau_{pq}(t) &= 0.1 \sin 3t + 0.3, \ \sigma_{pql}(t) = 0.1 \sin^2 t, \ \nu_{pql}(t) = 0.3 \sin 3t + 0.5, \\ I_p(t) &= e_0 \sqrt{20.01} \cos \sqrt{5}t + e_{12} \sqrt{20.01} \sin \sqrt{5}t + \frac{1}{1+t^2}. \end{split}$$

It is easy to get that

$$\begin{split} L_q^f &= 0.5, \ L_q^g = \frac{1}{3}, \ M_q^f = 1, \ M_q^g = 0.5, \ \hat{a}_{11} = \hat{a}_{12} = 0.0125, \\ \hat{a}_{21} &= \hat{a}_{22} = 0.0146, \ \hat{b}_{111} = \hat{b}_{112} = \hat{b}_{121} = \hat{b}_{122} = 0.0216, \\ \hat{b}_{211} &= \hat{b}_{212} = \hat{b}_{221} = \hat{b}_{222} = 0.0195, \ \hat{I}_1 = \hat{I}_2 = 4.5837. \end{split}$$

Let r = 10, $\delta = 0.1$, $\varepsilon = 0.5$, $t_0 = 0$. Choose $\alpha = 0.3$, 0.5 and 0.7, T = 10, one has $\max_{1 \le p \le 2} \left\{ \frac{C_p}{a_p} + \frac{\hat{I}_p}{a_p} \right\} = 0.6075 < 10 = r$, $\xi = \max_{1 \le p \le 2} \left\{ \frac{D_p}{a_p} \right\} = 0.0051 < 1$ and $M = \max_{1 \le p \le 2} D_p = 0.0406$. Using the MATLAB program, to compute the Mittag-Leffler function, we obtain

$$E_{0.3}(MT^{0.3}) = E_{0.3}(0.081) \approx 1.0982,$$



Figure 1. Curves of $x_1^l(t)$ and $x_2^l(t)$ of system (2.1) for $\alpha = 0.3, l = 0, 1, 2, 12$.



Figure 2. Curves of $x_1^l(t)$ and $x_2^l(t)$ of system (2.1) for $\alpha = 0.5, l = 0, 1, 2, 12$.

$$E_{0.5}(MT^{0.5}) = E_{0.5}(0.1284) \approx 1.1631,$$

 $E_{0.7}(MT^{0.7}) = E_{0.7}(0.2035) \approx 1.2615,$

then $\delta E_{0.3}(MT^{0.3}) = 0.10982 < \varepsilon, \delta E_{0.5}(MT^{0.5}) = 0.11631 < \varepsilon$ and $\delta E_{0.7}(MT^{0.7}) = 0.12615 < \varepsilon$.

Therefore, according to Theorem 3.2, system (2.1) possesses a unique pseudo almost periodic mild solution, which is finite-time stable (see Figures 1-9).

5. Conclusions

In this paper, we have obtained the existence and finite-time stability of the pseudo almost periodic mild solutions of FCHHNN (2.1). This is the first article to investigate the almost periodic mild solutions of fractional-order Clifford-valued differential equations via direct approach. The results and methods of this paper are new. And the methods used in this paper can be applied to study the existence of almost periodic or almost automorphic mild solutions to other types of fractional-order neural networks. The research on almost periodic synchronization and almost automorphic synchronization of fractional-order complex neural network systems is our future direction.



Figure 3. Curves of $x_1^l(t)$ and $x_2^l(t)$ of system (2.1) for $\alpha = 0.7, l = 0, 1, 2, 12$.



Figure 4. Finite-time stability of $x_1^0(t)$, $x_1^1(t)$, $x_1^2(t)$ and $x_1^{12}(t)$ of system (2.1) for $\alpha = 0.3$.



Figure 5. Finite-time stability of $x_2^0(t)$, $x_2^1(t)$, $x_2^2(t)$ and $x_2^{12}(t)$ of system (2.1) for $\alpha = 0.3$.



Figure 6. Finite-time stability of $x_1^0(t)$, $x_1^1(t)$, $x_1^2(t)$ and $x_1^{12}(t)$ of system (2.1) for $\alpha = 0.5$.



Figure 7. Finite-time stability of $x_2^0(t)$, $x_2^1(t)$, $x_2^2(t)$ and $x_2^{12}(t)$ of system (2.1) for $\alpha = 0.5$.



Figure 8. Finite-time stability of $x_1^0(t)$, $x_1^1(t)$, $x_1^2(t)$ and $x_1^{12}(t)$ of system (2.1) for $\alpha = 0.7$.



Figure 9. Finite-time stability of $x_2^0(t)$, $x_2^1(t)$, $x_2^2(t)$ and $x_2^{12}(t)$ of system (2.1) for $\alpha = 0.7$.

Abbreviations

FCHHNN fractional-order Clifford-valued high-order Hopfield neural network

- CVNN Clifford valued neural network
- HHNN high-order Hopfield neural network

References

- M. S. Ali, G. Narayanan, V. Shekher, A. Alsaedi and B. Ahmad, Global Mittag-Leffler stability analysis of impulsive fractional-order complex-valued BAM neural networks with time varying delays, Commun. Nonlinear Sci. Numer. Simulat., 2020, 83, paper no. 105088.
- [2] C. Aouiti and F. Dridi, Weighted pseudo almost automorphic solutions for neutral type fuzzy cellular neural networks with mixed delays and D operator in Clifford algebra, Int. J. Syst. Sci., 2020, 51(10), 1759–1781.
- [3] C. Aouiti and F. Touati, Global dissipativity of Clifford-valued multidirectional associative memory neural networks with mixed delays, Comput. Appl. Math., 2020, 39, paper no. 310.
- [4] N. Boonsatit, G. Rajchakit, R. Sriraman, C. P. Lim and P. Agarwal, Finite-/fixed-time synchronization of delayed Clifford-valued recurrent neural networks, Adv. Differ. Equ., 2021, 2021(2021), paper no. 276.
- [5] S. Breuils, K. Tachibana and E. Hitzer, New Applications of Clifford's Geometric Algebra, Adv. Appl. Clifford Algebras, 2022, 32, paper no. 17.
- [6] S. Buchholz, A Theory of Neural Computation with Clifford Algebras, PhD thesis, University of Kiel, 2005.
- [7] S. Buchholz and G. Sommer, On Clifford neurons and Clifford multi-layer perceptrons, Neural Netw., 2008, 21(7), 925–935.
- [8] J. Chen, Z. Zeng and P. Jiang, Global Mittag-Leffler stability and synchronization of memristor-based fractional-order neural networks, Neural Netw., 2014, 51, 1–8.

- [9] Y. Chen, X. Zhang and Y. Xue, Global exponential synchronization of highorder quaternion Hopfield neural networks with unbounded distributed delays and time-varying discrete delays, Math. Comput. Simul., 2022, 193, 173–189.
- [10] T. Diagana, Almost Automorphic Type and Almost Periodic Type Functions in Abstract Spaces, Springer, New York, 2013.
- [11] L. Duan, L. Huang, Z. Guo and X. Fang, Periodic attractor for reactiondiffusion high-order Hopfield neural networks with time-varying delays, Comput. Math. Appl., 2017, 73(2), 233–245.
- [12] Z. He, C. Li, H. Li and Q. Zhang, Global exponential stability of high-order Hopfield neural networks with state-dependent impulses, Physica A, 2020, 542, paper no. 123434.
- [13] E. Hitzer, T. Nitta and Y. Kuroe, Applications of Clifford's geometric algebra, Adv. Appl. Clifford Algebras, 2013, 23(2), 377–404.
- [14] S. Huang, Y. Y. Qiao and G. C. Wen, Real and Complex Clifford Analysis, Springer, New York, 2006.
- [15] N. Huo and Y. Li, Finite-time Sp-almost periodic synchronization of fractionalorder octonion-valued Hopfield neural networks, Chaos Solitons Fractals, 2023, 173, paper no. 113721.
- [16] B. Li, Y. Cao and Y. Li, Almost periodic oscillation in distribution for octonionvalued neutral-type stochastic recurrent neural networks with D operator, Nonlinear Dyn., 2023, 111, 11371–11388.
- [17] B. Li and Y. Li, Existence and global exponential stability of pseudo almost periodic solution for Clifford-valued neutral high-order Hopfield neural networks with leakage delays, IEEE Access, 2019, 7, 150213–150225.
- [18] Y. Li, M. Huang and B. Li, Besicovitch almost periodic solutions for fractionalorder quaternion-valued neural networks with discrete and distributed delays, Math. Meth. Appl. Sci., 2022, 45(8), 4791–4808.
- [19] Y. Li and B. Li, Pseudo compact almost automorphy of neutral type Cliffordvalued neural networks with mixed delays, Discrete Contin. Dyn. Syst.-B, 2022, 27(9), 4703–4724.
- [20] Y. Li, C. Ruan and B. Li, Existence and finite-time stability of Besicovitch almost periodic solutions of fractional-order quaternion-valued neural networks with time-varying delays, Neural Process. Lett., 2022, 54, 2127–2141.
- [21] Y. Li and X. Wang, Almost periodic solutions in distribution of Clifford-valued stochastic recurrent neural networks with time-varying delays, Chaos Solitons Fractals, 2021, 153, 111536.
- [22] Y. Li, X. Wang and N. Huo, Weyl almost automorphic solutions in distribution sense of Clifford-valued stochastic neural networks with time-varying delays, Proc. Royal Soc. A, 2022, 478(2257), paper no. 20210719.
- [23] Y. Li, Y. Wang and B. Li, Existence and finite-time stability of a unique almost periodic positive solution for fractional-order Lasota-Wazewska red blood cell models, Int. J. Biomath., 2020, 13(2), paper no. 2050013.
- [24] Y. Liu, P. Xu, J. Lu and J. Liang, Global stability of Clifford-valued recurrent neural networks with time delays, Nonlinear Dyn., 2016, 84(2), 767–777.

- [25] D. Luo, Q. Jiang and Q. Wang, Anti-periodic solutions on Clifford-valued highorder Hopfield neural networks with multi-proportional delays, Neurocomputing, 2022, 472, 1–11.
- [26] W. Lv and B. Li, Existence and global attractivity of pseudo almost periodic solutions for Clifford-valued fuzzy neural networks with proportional delays, Math., 2021, 9(24), paper no. 3306.
- [27] I. Podlubny, Fractional Differential Equations, Mathematics in Science and Engineering, Academic Press, New York, 1999.
- [28] G. Rajchakit, P. Chanthorn, P. Kaewmesri, R. Sriraman and C. P. Lim, Global Mittag-Leffler stability and stabilization analysis of fractional-order quaternionvalued memristive neural networks, Math., 2020, 8(3), paper no. 422.
- [29] G. Rajchakit, P. Chanthorn, M. Niezabitowski, R. Raja, D. Baleanu and A. Pratap, *Impulsive effects on stability and passivity analysis of memristor-based fractional-order competitive neural networks*, Neurocomputing, 2020, 417, 290–301.
- [30] G. Rajchakit, A. Pratap, R. Raja, J. Cao, J. Alzabut and C. Huang, Hybrid control scheme for projective lag synchronization of Riemann-Liouville sense fractional order memristive BAM neural networks with mixed delays, Math., 2019, 7(8), paper no. 759.
- [31] G. Rajchakit, R. Sriraman, N. Boonsatit, P. Hammachukiattikul, C. P. Lim and P. Agarwal, Global exponential stability of Clifford-valued neural networks with time-varying delays and impulsive effects, Adv. Differ. Equ., 2021, 2021(2021), paper no. 208.
- [32] G. Rajchakit, R. Sriraman, C. P. Lim and B. Unyong, Existence, uniqueness and global stability of Clifford-valued neutral-type neural networks with time delays, Math. Comput. Simul., 2022, 201, 508–527.
- [33] G. Rajchakit, R. Sriraman, P. Vignesh and C. P. Lim, Impulsive effects on Clifford-valued neural networks with time-varying delays: An asymptotic stability analysis, Appl. Math. Comput., 2021, 407, paper no. 126309.
- [34] C. Xu, M. Liao, P. Li, L. Yao, Q. Qin and Y. Shang, Chaos control for a fractional-order jerk system via time delay feedback controller and mixed controller, Frac. Fract., 2021, 5(4), paper no. 257.
- [35] C. Xu, M. Liao, P. Li and S. Yuan, Impact of leakage delay on bifurcation in fractional-order complex-valued neural networks, Chaos Solitons Fractals, 2021, 142, paper no. 110535.
- [36] C. Xu, Z. Liu, M. Liao, P. Li, Q. Xiao and S. Yuan, Fractional-order bidirectional associate memory (BAM) neural networks with multiple delays: The case of Hopf bifurcation, Math. Comput. Simul., 2021, 182, 471–494.
- [37] Y. Xu, T. Lin, X. Liu and W. Li, Exponential bipartite synchronization of fractional-order multilayer signed networks via hybrid impulsive control, IEEE Trans. Cyber., 2023, 53, 3926–3938.
- [38] Y. Xu, F. Sun and W. Li, Exponential synchronization of fractional-order multilayer coupled neural networks with reaction-diffusion terms via intermittent control, Neural Comput. Appl., 2021, 33(23), 16019–16032.

- [39] Y. Xu, J. Yu, W. Li and J. Feng, Global asymptotic stability of fractional-order competitive neural networks with multiple time-varying-delay links, Appl. Math. Comput., 2021, 389, paper no. 125498.
- [40] H. Ye, J. Gao and Y. Ding, A generalized Gronwall inequality and its application to a fractional differential equation, J. Math. Anal. Appl., 2007, 328, 1075–1081.
- [41] J. Yu, C. Hu and H. Jiang, α-stability and α-synchronization for fractionalorder neural networks, Neural Netw., 2012, 35, 82–87.
- [42] F. Zhang, T. Huang, Q. Wu and Z. Zeng, Multistability of delayed fractionalorder competitive neural networks, Neural Netw., 2021, 140, 325–335.
- [43] S. Zhang, Y. Yu and J. Yu, LMI conditions for global stability of fractionalorder neural networks, IEEE Trans. Neural Netw. Learn. Syst., 2016, 28(10), 2423–2433.