

APPLICATIONS OF VARIATIONAL ITERATION METHOD TO A CLASS OF ORDINARY DIFFERENTIAL EQUATIONS

Lixin Liu^{1,†}

Abstract This paper proves the convergence of the variational iteration method for a class of n -th order ordinary differential equations with Lipschitz nonlinearity which can be regarded as a generalization of oscillation equations.

Keywords Variational iteration method, convergence analysis, oscillation equations.

MSC(2010) 49M30, 65K10.

1. Introduction

The variational iteration method was first proposed by He to approximately solve some nonlinear ordinary differential equations and nonlinear partial differential equations without linearization or small perturbation [4–6, 9]. This method was shown by many authors to be superior to many other analytic approximation methods, such as Adomian's decomposition method and perturbation methods. A key point is that a correction functional is constructed by a general Lagrange multiplier [10], which can be identified via variational methods.

In this paper we attempt to use the variational iteration method to get an exact solution of the following n -th order ordinary differential equation

$$\begin{cases} u^{(n)}(t) + a_{n-1}(t)u^{(n-1)}(t) + \cdots + a_1(t)u'(t) + F(t, u(t)) = 0, & t \in \mathbf{R}, \\ u^{(n-1)}(0) = u_0^{(n-1)}, \dots, u'(0) = u'_0, u(0) = u_0, \end{cases} \quad (1.1)$$

where $a_i \in C^{n-1}(\mathbf{R})$, $i = 1, 2, 3, \dots, n-1$, $F : \mathbf{R} \times \mathbf{R} \rightarrow \mathbf{R}$ is continuous function that is Lipschitz continuous with respect to the second variable, i.e., there exists constant $L > 0$ such that

$$|F(t, u_1) - F(t, u_2)| \leq L|u_1 - u_2|, \quad \forall t, u_1, u_2 \in \mathbf{R}.$$

The motivation for the study of the above equation originated from nonlinear oscillators. Several authors showed the validity of variational iteration method and got approximate solutions of oscillation equations, see for example, [7, 15]. See [1, 3, 11, 12, 16, 17] and the references therein for other new developments of variational iteration methods. The oscillation equations considered in this paper has more general form with Lipschitz nonlinear term. Our main theorem (Theorem

[†]The corresponding author.

¹Jilin Business and Technology College, Changchun 130507, China
Email: 17790097972@163.com(L. Liu)

3.1) shows that variational iteration sequence of (1.1) converges uniformly on any finite time interval to an exact solution.

2. Preliminary

In this section, we briefly recall the basic idea of the variational iteration method. Consider the following general nonlinear system

$$L(u(t)) + N(u(t)) = g(t), \quad (2.1)$$

where L is a linear operator, N is a nonlinear operator, and g is a given continuous function.

The essential technique of the variational iteration method is to construct a correction functional for system (2.1) as follows

$$u_{n+1}(t) = u_n(t) + \int_0^t \lambda(s; t) \left(Lu_n(s) + N\tilde{u}_n(s) - g(s) \right) ds, \quad (2.2)$$

where λ is a general Lagrange multiplier which can be determined by using the variational approach, and \tilde{u}_n denotes a restricted variation, i.e., $\delta\tilde{u}_n = 0$. See [2, 8, 13, 14, 18] for more details on iteration methods.

3. Main theorem

Theorem 3.1. *For any given $T > 0$, any initial data $v \in C^n([-T, T])$ satisfying the initial condition in (1.1), the variational iteration sequence of (1.1) converges uniformly on $[-T, T]$ to an exact solution.*

Proof. First we derive the variational iteration sequence of (1.1). Let $\lambda = \lambda(s; t)$ be a general Lagrange multiplier to be determined. Then the correction functional for (1.1) is

$$\begin{aligned} u_{k+1}(t) = & u_k(t) + \int_0^t \lambda(s; t) \left(u_k^{(n)}(s) + a_{n-1}(s)u_k^{(n-1)}(s) + \cdots \right. \\ & \left. + a_1(s)u_k'(s) + F(s, u_k(s)) \right) ds. \end{aligned}$$

Making the above correction functional stationary, notice that $\delta u_k(0) = 0$,

$$\begin{aligned} \delta u_{k+1}(t) = & \delta u_k(t) + \delta \int_0^t \lambda(s; t) \left(u_k^{(n)}(s) + a_{n-1}(s)(\tilde{u}_k)^{(n-1)}(s) + \cdots \right. \\ & \left. + a_1(s)(\tilde{u}_k)'(s) + F(s, \tilde{u}_k(s)) \right) ds \\ = & \delta u_k(t) + \int_0^t \lambda(s; t) (\delta u_k)^{(n)}(s) ds \\ = & \left(1 + (-1)^{n-1} \frac{\partial^{n-1} \lambda}{\partial s^{n-1}}(t; t) \right) \delta u_k(t) + \sum_{j=1}^{n-1} (-1)^{j-1} \frac{\partial^{j-1} \lambda}{\partial s^{j-1}}(t; t) (\delta u_k)^{(n-j)}(t) \\ & + \int_0^t (-1)^n \frac{\partial^n \lambda}{\partial s^n}(s; t) (\delta u_k)(s) ds \\ = & 0, \end{aligned}$$

where \tilde{u}_k denotes the restricted variation, i.e., $\delta\tilde{u}_k = 0$.

Thus the arbitrariness of δu_k yields

$$\begin{cases} \frac{\partial^n \lambda}{\partial s^n}(s; t) = 0, \forall s \in [0, t], \\ \frac{\partial^{j-1} \lambda}{\partial s^{j-1}}(t; t) = 0, \forall j = 1, 2, 3, \dots, n-1, \\ \frac{\partial^{n-1} \lambda}{\partial s^{n-1}}(t; t) = (-1)^n, \end{cases}$$

which can be readily solved to obtain $\lambda(s; t) = \frac{(-1)^n}{(n-1)!}(s-t)^{n-1}$. So the variational iteration sequence of (1.1) reads

$$\begin{aligned} u_{k+1}(t) = & u_k(t) + \int_0^t \frac{(-1)^n}{(n-1)!} (s-t)^{n-1} \left(u_k^{(n)}(s) + a_{n-1}(s)u_k^{(n-1)}(s) + \dots \right. \\ & \left. + a_1(s)u_k'(s) + F(s, u_k(s)) \right) ds. \end{aligned}$$

Now we define an operator $\mathbb{A}_T : C^0([-T, T]) \rightarrow C^0([-T, T])$ by

$$\begin{aligned} \mathbb{A}_T[u](t) := & \sum_{j=1}^{n-1} (-1)^j \int_0^t \left(\frac{(-1)^n (s-t)^{n-1}}{(n-1)!} a_j(s) \right)^{(j)} u(s) ds \\ & + \int_0^t \frac{(-1)^n (s-t)^{n-1}}{(n-1)!} F(s, u(s)) ds. \end{aligned}$$

It can be easily seen that \mathbb{A}_T is well-defined. Also defined $\tilde{\mathbb{A}}_T : C^n([-T, T]) \rightarrow C^n([-T, T])$ by

$$\begin{aligned} \tilde{\mathbb{A}}_T[u](t) := & u(t) + \int_0^t \frac{(-1)^n (s-t)^{n-1}}{(n-1)!} \left(u^{(n)}(s) + a_{n-1}(s)u^{(n-1)}(s) + \dots \right. \\ & \left. + a_1(s)u'(s) + F(s, u(s)) \right) ds. \end{aligned}$$

The classical Cauchy-Lipschitz theorem and extension theorem guarantee that the unique exact solution $\varphi(t)$ of (1.1) exists on $[-T, T]$. It is clear that φ is a fixed point of $\tilde{\mathbb{A}}_T$, i.e., $\tilde{\mathbb{A}}_T[\varphi] = \varphi$. Therefore for any $v \in C^n([-T, T])$ satisfying the initial condition in (1.1), we have

$$\begin{aligned} & |\tilde{\mathbb{A}}_T[v](t) - \varphi(t)| \\ = & |\tilde{\mathbb{A}}_T[v](t) - \tilde{\mathbb{A}}_T[\varphi](t)| \\ = & \left| (v - \varphi)(t) + \int_0^t \frac{(-1)^n (s-t)^{n-1}}{(n-1)!} \left((v^{(n)} - \varphi^{(n)})(s) + \sum_{j=1}^{n-1} a_j(s)(v^{(j)} - \varphi^{(j)})(s) \right. \right. \\ & \left. \left. + F(s, v(s)) - F(s, \varphi(s)) \right) ds \right| \\ = & \left| \sum_{j=1}^{n-1} (-1)^j \int_0^t \left(\frac{(-1)^j (s-t)^{n-1}}{(n-1)!} a_j(s) \right)^{(j)} (v - \varphi)(s) \right. \\ & \left. + \int_0^t \frac{(-1)^n (s-t)^{n-1}}{(n-1)!} \left(F(s, v(s)) - F(s, \varphi(s)) \right) ds \right| \end{aligned}$$

$$\begin{aligned}
&= |\mathbb{A}_T[v](t) - \mathbb{A}_T[\varphi](t)| \\
&\leq C(n, T, M) \left| \int_0^t |v - \varphi|(s) ds \right| \\
&\leq C(n, T, M) \|v - \varphi\|_{C^0([-T, T])} |t|, \quad \forall t \in [-T, T],
\end{aligned} \tag{3.1}$$

where $C(n, T, M)$ is a positive constant depending on $n, T, M := \max\{\|a_j\|_{C^{n-1}([-T, T])}; 1 \leq j \leq n-1\}$ and L .

One more step iteration gives

$$\begin{aligned}
|\tilde{\mathbb{A}}_T^2[v](t) - \varphi(t)| &= |\tilde{\mathbb{A}}_T(\tilde{\mathbb{A}}_T[v])(t) - \tilde{\mathbb{A}}_T(\tilde{\mathbb{A}}_T[\varphi])(t)| \\
&\leq C(n, T) \left| \int_0^t |\tilde{\mathbb{A}}_T[v](s) - \tilde{\mathbb{A}}_T[\varphi](s)| ds \right| \\
&= C(n, T) \left| \int_0^t |\mathbb{A}_T[v](s) - \mathbb{A}_T[\varphi](s)| ds \right| \\
&\leq C(n, T, M)^2 \left| \int_0^t \|v - \varphi\|_{C^0([-T, T])} s ds \right| \\
&= \frac{C(n, T, M)^2 \|v - \varphi\|_{C^0([-T, T])}}{2!} t^2, \quad \forall t \in [-T, T].
\end{aligned}$$

Inductively we have

$$|\tilde{\mathbb{A}}_T^N[v](t) - \varphi(t)| \leq \frac{C(n, T, M)^N \|v - \varphi\|_{C^0([-T, T])}}{N!} t^N, \quad \forall t \in [-T, T].$$

Thus

$$\|\tilde{\mathbb{A}}_T^N[v] - \varphi\|_{0, [-T, T]} \leq \frac{C(n, T, M)^N \|v - \varphi\|_{C^0([-T, T])}}{N!} (2T)^N \rightarrow 0, \quad N \rightarrow \infty.$$

The proof is complete. \square

Example 3.1. (Dissipative pendulum)

Consider the equation of dissipative pendulum with external force

$$\begin{cases} u''(t) + u'(t) + \sin(u(t)) = 0, & t \in \mathbf{R}, \\ u(0) = A > 0, u'(0) = 0. \end{cases} \tag{3.2}$$

Here $F(u) = \sin u$ is a Lipschitz function with Lipschitz constant 1. The variational iteration formula for this equation reads

$$u_{n+1}(t) = u_n(t) + \int_0^t (s-t) \left(u_n''(s) + u_n'(s) + \sin(u_n(s)) \right) ds.$$

Rewrite (3.2) into planar system

$$\begin{cases} \dot{u} = v, \\ \dot{v} = -v - \sin(u). \end{cases} \tag{3.3}$$

It can be easily seen that $(0, 0)$ is a saddle point, so for $A > 0$ small, the orbit of (3.3) will spiral towards the origin. For $A = 0.1$, initial approximation guess $v(t) = A \cos(t)$ and time interval $[0, 5]$. Figure 1 and Figure 2 show a comparison between the exact solutions versus 10-th and 15-th approximation solutions respectively. Graphically, the more we iterate, the closer, in an uniform way, these two solutions with each other.




Figure 1




Figure 2

Acknowledgements

I would like to thank the referee for his/her careful reading of this paper and for many valuable suggestions.

References

- [1] S. Cetinkaya and A. Demir, *On applications Shehu variational iteration method to time fractional initial boundary value problems*, Konuralp J. Math., 2024, 12(1), 13–20.
- [2] Y. Cui, S. Kang and H. Chen, *Iterative positive solutions to nonlinear q -fractional differential equations with integral boundary value conditions*, J. Nonl. Mod. Anal., 2023, 5, 446–455.
- [3] A. Entesar and O. Qasim, *Solve fractional differential equations via a hybrid method between variational iteration method and gray wolf optimization algorithm*, Asian-Eur. J. Math., 2021, 14(8), Paper No. 2150144, 9 pp.
- [4] J. He, *Approximate analytical solution for seepage flow with fractional derivatives in porous media*, Computer Methods in Applied Mechanics and Engineering, 1998, 1678, 57–68.

- [5] J. He, *Approximate solution of nonlinear differential equations with convolution product nonlinearities*, Computer Methods in Applied Mechanics and Engineering, 1998, 167, 69–73.
- [6] J. He, *Variational iteration method-a kind of non-linear analytical technique: Some examples*, International Journal of Non-linear Mechanics, 1999, 34, 699–708.
- [7] J. He, *Variational approach for nonlinear oscillators*, Chaos, Solitons & Fractals, 2007, 34, 1430–1439.
- [8] J. He, *A short remark on fractional variational iteration method*, Phys. Lett. A, 2011, 375(38), 3362–3364.
- [9] J. He and X. Wu, *Construction of solitary solution and compacton-like solution by variational iteration method*, Chaos, Solitons & Fractals, 2006, 29, 108–113.
- [10] M. Inokuti, H. Sekine and T. Mura, *General use of the Lagrange multiplier in nonlinear mathematical physics*, Variational Methods in the Mechanics of Solids, 1978, 33, 156–162.
- [11] D. Kumar, H. Jassim, J. Singh and M. Diykh, *Fractional Elzaki variational iteration method for solving nonlinear biological population model*, Lect. Notes Netw. Syst., 2024, 952, Springer, Cham, 368–376.
- [12] G. Monzón, *Fractional variational iteration method for higher-order fractional differential equations*, J. Fract. Calc. Appl., 2024, 15(1), Paper No. 4, 15 pp.
- [13] M. Nadeem and J. He, *He-Laplace variational iteration method for solving the nonlinear equations arising in chemical kinetics and population dynamics*, J. Math. Chem., 2021, 59(5), 1234–1245.
- [14] A. Prakash and M. Kumar, *He's variational iteration method for the solution of nonlinear Newell-Whitehead-Segel equation*, J. Appl. Anal. Comput., 2016, 6(3), 738–748.
- [15] M. Rafei, D. Ganji, H. Daniali and H. Pashaei, *The variational iteration method for nonlinear oscillators with discontinuities*, Journal of Sound and Vibration, 2007, 305, 614–620.
- [16] N. Sharma, G. Alhawael, P. Goswami and S. Joshi, *Variational iteration method for n-dimensional time-fractional Navier-Stokes equation*, Appl. Math. Sci. Eng., 2024, 32(1), Paper No. 2334387, 20 pp.
- [17] M. Shirazian, *A new acceleration of variational iteration method for initial value problems*, Math. Comput. Simulation, 2023, 214, 246–259.
- [18] J. Wen, C. Yue, Z. Liu and D. O'Regan, *A fractional Landweber iteration method for simultaneous inversion in a time-fractional diffusion equation*, J. Appl. Anal. Comput., 2023, 13(6), 3374–3402.