

# INERTIAL EXTRAPOLATION SCHEME FOR SYSTEM OF VARIATIONAL INCLUSIONS USING GENERALIZED YOSIDA AND CAYLEY OPERATORS

Imran Ali<sup>1</sup>, Mohd Ishtyak<sup>2</sup>, Javid Iqbal<sup>3</sup>, Yuanheng Wang<sup>4,†</sup>  
and Rais Ahmad<sup>5</sup>

**Abstract** The aim of this work is to study a system of variational inclusions involving generalized Yosida and Cayley operators through inertial extrapolation scheme in real Banach space. To obtain faster convergence of the sequences generated by algorithm, we use one inertial extrapolation scheme, although we have established some more iterative schemes. To achieve our goal, we prove an important Lemma ensuring the convergence of sum of two sequences. We provide a numerical example.

**Keywords** Cayley, Yosida, solution, convergence, scheme.

**MSC(2010)** 65J15, 47J25, 65K15.

## 1. Introduction

Variational inclusions are application oriented and can be treated as mathematical version of many problems of day-to-day life, such as economics, physics, engineering and space sciences, etc.. The system of variational inclusions extends the concept of variational inequalities. These systems have applications across various fields such as mathematical analysis, biological sciences, elasticity, image processing, biomedical sciences, and optimization. Furthermore, investigating variational inclusion systems provides novel methods for tackling analytical problems. For more literature on system of variational inclusions, one can see [1, 5, 7, 10–13, 22, 24, 27] and references therein.

---

<sup>†</sup>The corresponding author.

<sup>1</sup>Department of Engineering Mathematics, College of Engineering, Koneru Lakshmaiah Education Foundation, Vaddeswaram 522302, India

<sup>2</sup>Department of Applied Mathematics, Aligarh Muslim University, Aligarh 202002, India

<sup>3</sup>Department of Mathematics, Baba Ghulam Shah Badshah University, Rajouri 185234, India

<sup>4</sup>School of Mathematical Sciences, Zhejiang Normal University, Jinhua 321004, China

<sup>5</sup>Department of Mathematics, Aligarh Muslim University, Aligarh 202002, India

Email: [imranali@kluniversity.in](mailto:imranali@kluniversity.in)(I. Ali),  
[ishtyakalig@gmail.com](mailto:ishtyakalig@gmail.com)(M. Ishtyak), [javid2iqbal@gmail.com](mailto:javid2iqbal@gmail.com)(J. Iqbal),  
[yhwang@zjnu.cn](mailto:yhwang@zjnu.cn)(Y. Wang), [raisain.123@rediffmail.com](mailto:raisain.123@rediffmail.com)(R. Ahmad)

Yosida approximation operator are of great importance due to their applications. In the study of wave equations, heat equations and heat flow, etc., one can find the clear applications of Yosida approximation operators. For more details, we refer to [4, 14, 21].

The Cayley transform is a mapping that connects skew-symmetric matrices to special orthogonal matrices and is utilized in real, complex, and quaternionic analysis. In the context of Hilbert spaces, it serves as a mapping between linear operators. Considering the real projective line, the Cayley transform permutes the elements  $1, 0, -1, \infty$  in sequence and maps the positive real numbers to the interval  $[-1, 1]$ . Consequently, the Legendre polynomials can be applied to functions on the positive real numbers using the Cayley transform, resulting in Legendre rational functions.

On the Riemann sphere, the Cayley transform is given by

$$f(z) = \frac{z - i}{z + i}.$$

This transform maps the points  $\{\infty, 1, -1\}$  to  $\{1, -i, i\}$ . As a Möbius transformation, it permutes generalized circles in the complex plane, mapping the real line onto the unit circle. For more details, see [6, 8, 9, 15, 18, 19].

Various iterative algorithms appeared in the literature using proximal operators, resolvent operators, projection operators as well as sub-differential operator. In order to obtain faster convergence of the sequences generated by the considered algorithm, we have to choose such a scheme which expedite the speed of convergence. Several authors have used inertial extrapolation scheme using inertial extrapolation term  $\gamma(u_n - u_{n-1})$ , where  $\gamma$  is the extrapolation factor which accelerates the convergence rate of the method. While dealing with heavy ball method, Polyak [23], introduced inertial-type algorithm. There are two steps in the inertial-type algorithm, through these two steps consecutive iterations are gained by using former two terms, for reference see [20, 25, 26].

In view of the above mentioned facts, in this paper, we study a system of variational inclusions involving generalized Yosida and Cayley operators using inertial extrapolation scheme in real Banach space. Simultaneously, we have developed some more iterative schemes for our problem. The existence of a solution and the convergence of the sequences produced by our scheme are demonstrated. We construct a numerical example.

## 2. Preliminary tools and hypothesis

Let  $\widehat{\mathcal{X}}$  be a real Banach space with its topological dual  $\widehat{\mathcal{X}}^*$ . We denote the norm on  $\widehat{\mathcal{X}}$  by  $\|\cdot\|$  and duality pairing by  $\langle \cdot, \cdot \rangle$  between  $\widehat{\mathcal{X}}$  and  $\widehat{\mathcal{X}}^*$ . The class of all non-empty subsets of  $\widehat{\mathcal{X}}$  are denoted by  $2^{\widehat{\mathcal{X}}}$ .

**Definition 2.1.** The normalized duality mapping  $J : \widehat{\mathcal{X}} \rightarrow \widehat{\mathcal{X}}^*$  is defined by

$$J(\widehat{p}) = \{\widehat{q} \in \widehat{\mathcal{X}}^* : \langle \widehat{p}, \widehat{q} \rangle = \|\widehat{p}\|^2, \|\widehat{p}\| = \|\widehat{q}\|\}, \text{ for all } \widehat{p} \in \widehat{\mathcal{X}}.$$

**Definition 2.2.** The operator  $\widetilde{\mathcal{A}} : \widehat{\mathcal{X}} \rightarrow \widehat{\mathcal{X}}$  is said to be

(i) Accretive, if

$$\langle \widetilde{\mathcal{A}}(\widehat{p}) - \widetilde{\mathcal{A}}(\widehat{q}), J(\widehat{p} - \widehat{q}) \rangle \geq 0, \text{ for all } \widehat{p}, \widehat{q} \in \widehat{\mathcal{X}}.$$

(ii) Strongly accretive, if

$$\langle \tilde{\mathcal{A}}(\hat{p}) - \tilde{\mathcal{A}}(\hat{q}), J(\hat{p} - \hat{q}) \rangle \geq r_1 \|\hat{p} - \hat{q}\|^2, \text{ for all } \hat{p}, \hat{q} \in \hat{\mathcal{X}},$$

where  $r_1 > 0$  is a constant.

(iii) Lipschitz continuous, if

$$\|\tilde{\mathcal{A}}(\hat{p}) - \tilde{\mathcal{A}}(\hat{q})\| \leq \lambda_{\tilde{\mathcal{A}}} \|\hat{p} - \hat{q}\|, \text{ for all } \hat{p}, \hat{q} \in \hat{\mathcal{X}},$$

where  $\lambda_{\tilde{\mathcal{A}}} > 0$  is a constant.

**Definition 2.3.** A multi-valued mapping  $\mathcal{M} : \hat{\mathcal{X}} \rightarrow 2^{\hat{\mathcal{X}}}$  is said to be accretive, if for all  $\hat{p}, \hat{q} \in \hat{\mathcal{X}}$

$$\langle u - v, J(\hat{p} - \hat{q}) \rangle \geq 0, \text{ for all } u \in \mathcal{M}(\hat{p}), v \in \mathcal{M}(\hat{q}).$$

**Definition 2.4.** [16, 28] Let  $\tilde{\mathcal{A}} : \hat{\mathcal{X}} \rightarrow \hat{\mathcal{X}}$  be a mapping. A multi-valued mapping  $\mathcal{M} : \hat{\mathcal{X}} \rightarrow 2^{\hat{\mathcal{X}}}$  is said to be  $\tilde{\mathcal{A}}$ -accretive if  $\mathcal{M}$  is accretive and

$$[\tilde{\mathcal{A}} + \rho\mathcal{M}](\hat{\mathcal{X}}) = \hat{\mathcal{X}}, \text{ where } \rho > 0 \text{ is a constant.}$$

**Definition 2.5.** [16] Let  $\tilde{\mathcal{A}} : \hat{\mathcal{X}} \rightarrow \hat{\mathcal{X}}$  be a mapping and  $\mathcal{M} : \hat{\mathcal{X}} \rightarrow 2^{\hat{\mathcal{X}}}$  be  $\tilde{\mathcal{A}}$ -accretive multi-valued mapping. The generalized resolvent operator  $R_{\tilde{\mathcal{A}}, \rho}^{\mathcal{M}} : \hat{\mathcal{X}} \rightarrow \hat{\mathcal{X}}$  associated with  $\tilde{\mathcal{A}}$ , is defined as:

$$R_{\tilde{\mathcal{A}}, \rho}^{\mathcal{M}}(\hat{p}) = [\tilde{\mathcal{A}} + \rho\mathcal{M}]^{-1}(\hat{p}), \text{ for all } \hat{p} \in \hat{\mathcal{X}} \text{ and } \rho > 0 \text{ is a constant.}$$

**Theorem 2.1.** [2] Let  $\tilde{\mathcal{A}} : \hat{\mathcal{X}} \rightarrow \hat{\mathcal{X}}$  be strongly accretive operator with constant  $r_1$  and  $\mathcal{M} : \hat{\mathcal{X}} \rightarrow 2^{\hat{\mathcal{X}}}$  be  $\tilde{\mathcal{A}}$ -accretive multi-valued mapping. Then

$$\|R_{\tilde{\mathcal{A}}, \rho}^{\mathcal{M}}(\hat{p}) - R_{\tilde{\mathcal{A}}, \rho}^{\mathcal{M}}(\hat{q})\| \leq \frac{1}{r_1} \|\hat{p} - \hat{q}\|, \text{ for all } \hat{p}, \hat{q} \in \hat{\mathcal{X}}.$$

That is, the generalized resolvent operator  $R_{\tilde{\mathcal{A}}, \rho}^{\mathcal{M}}$  is Lipschitz continuous.

**Definition 2.6.** [2] Let  $\tilde{\mathcal{B}} : \hat{\mathcal{X}} \rightarrow \hat{\mathcal{X}}$  be a mapping and  $R_{\tilde{\mathcal{B}}, \gamma}^{\mathcal{N}} : \hat{\mathcal{X}} \rightarrow \hat{\mathcal{X}}$  is the generalized resolvent operator associated with  $\tilde{\mathcal{B}}$ . The generalized Cayley operator  $C_{\tilde{\mathcal{B}}, \gamma}^{\mathcal{N}} : \hat{\mathcal{X}} \rightarrow \hat{\mathcal{X}}$  is defined as

$$C_{\tilde{\mathcal{B}}, \gamma}^{\mathcal{N}}(\hat{p}) = [2R_{\tilde{\mathcal{B}}, \gamma}^{\mathcal{N}} - \tilde{\mathcal{B}}](\hat{p}), \text{ for all } \hat{p} \in \hat{\mathcal{X}} \text{ and } \gamma > 0 \text{ is a constant.}$$

**Proposition 2.1.** [2, 17] The generalized Cayley operator  $C_{\tilde{\mathcal{B}}, \gamma}^{\mathcal{N}} : \hat{\mathcal{X}} \rightarrow \hat{\mathcal{X}}$  is Lipschitz continuous with constant  $\lambda_C$ , that is

$$\|C_{\tilde{\mathcal{B}}, \gamma}^{\mathcal{N}}(\hat{p}) - C_{\tilde{\mathcal{B}}, \gamma}^{\mathcal{N}}(\hat{q})\| \leq \lambda_C \|\hat{p} - \hat{q}\|, \text{ for all } \hat{p}, \hat{q} \in \hat{\mathcal{X}},$$

where  $\lambda_C = \frac{2 + \lambda_{\tilde{\mathcal{B}}} r_2}{r_2}$  and the generalized resolvent operator  $R_{\tilde{\mathcal{B}}, \gamma}^{\mathcal{N}} : \hat{\mathcal{X}} \rightarrow \hat{\mathcal{X}}$  is  $\frac{1}{r_2}$ -Lipschitz continuous.

**Definition 2.7.** [3] The generalized Yosida approximation operator  $Y_{\tilde{\mathcal{A}},\rho}^{\mathcal{M}} : \hat{\mathcal{X}} \rightarrow \hat{\mathcal{X}}$  is defined as

$$Y_{\tilde{\mathcal{A}},\rho}^{\mathcal{M}}(\hat{p}) = \frac{1}{\rho} \left[ \tilde{\mathcal{A}} - R_{\tilde{\mathcal{A}},\rho}^{\mathcal{M}} \right] (\hat{p}), \text{ for all } \hat{p} \in \hat{\mathcal{X}} \text{ and } \rho > 0 \text{ is a constant.}$$

**Proposition 2.2.** [3] The generalized Yosida approximation operator  $Y_{\tilde{\mathcal{A}},\rho}^{\mathcal{M}} : \hat{\mathcal{X}} \rightarrow \hat{\mathcal{X}}$  is Lipschitz continuous with constant  $\lambda_Y$ , that is

$$\left\| Y_{\tilde{\mathcal{A}},\rho}^{\mathcal{M}}(\hat{p}) - Y_{\tilde{\mathcal{A}},\rho}^{\mathcal{M}}(\hat{q}) \right\| \leq \lambda_Y \|\hat{p} - \hat{q}\|, \text{ for all } \hat{p}, \hat{q} \in \hat{\mathcal{X}},$$

where  $\lambda_Y = \frac{\lambda_{\tilde{\mathcal{A}}} r_1 + 1}{\rho r_1}$  and the generalized resolvent operator  $R_{\tilde{\mathcal{A}},\rho}^{\mathcal{M}} : \hat{\mathcal{X}} \rightarrow \hat{\mathcal{X}}$  is  $\frac{1}{r_1}$ -Lipschitz continuous.

**Lemma 2.1.** Let  $\{s_n\}$  and  $\{t_n\}$  be sequences of non-negative real numbers such that

$$\begin{aligned} s_{n+1} &\leq (1 - a_n)s_n + a_n\hat{\alpha}_n + \xi_n, \\ \text{and } t_{n+1} &\leq (1 - a_n)t_n + a_n\hat{\beta}_n + \delta_n, \text{ for all } n \geq 1, \end{aligned}$$

where

- (i)  $\{a_n\} \subset [0, 1]$ ,  $\sum_{n=1}^{\infty} a_n = \infty$  or equivalently  $\prod_{n=1}^{\infty} (1 - a_n) = 0$ ,
- (ii)  $\limsup(\hat{\alpha}_n + \hat{\beta}_n) \leq 0$ ,
- (iii)  $\xi_n \geq 0$ ,  $\delta_n \geq 0$ ,  $\sum_{n=1}^{\infty} \xi_n < \infty$ ,  $\sum_{n=1}^{\infty} \delta_n < \infty$ .

Then

$$s_n + t_n \rightarrow 0, \text{ as } n \rightarrow \infty.$$

**Proof.** For any  $\epsilon > 0$ , let  $N$  be an integer such that

$$\alpha_n < \frac{\epsilon}{2}, \beta_n < \frac{\epsilon}{2}, \sum_{n=N}^{\infty} \xi_n < \frac{\epsilon}{2} \text{ and } \sum_{n=N}^{\infty} \delta_n < \frac{\epsilon}{2}, \text{ } n \geq N.$$

Given,

$$s_{n+1} \leq (1 - a_n)s_n + a_n\hat{\alpha}_n + \xi_n, \quad (2.1)$$

so, we have

$$\begin{aligned} s_1 &\leq (1 - a_0)s_0 + a_0\hat{\alpha}_0 + \xi_0, \\ s_2 &\leq (1 - a_1)s_1 + a_1\hat{\alpha}_1 + \xi_1, \\ &\vdots \\ s_N &\leq (1 - a_{N-1})s_{N-1} + a_{N-1}\hat{\alpha}_{N-1} + \xi_{N-1}, \\ &\vdots \\ s_n &\leq (1 - a_{n-1})s_{n-1} + a_{n-1}\hat{\alpha}_{n-1} + \xi_{n-1}. \end{aligned} \quad (2.2)$$

Combining (2.1) and (2.2), we obtain

$$s_{n+1} \leq (1 - a_n) \{ (1 - a_{n-1})s_{n-1} + a_{n-1}\hat{\alpha}_{n-1} + \xi_{n-1} \} + a_n\hat{\alpha}_n + \xi_n. \quad (2.3)$$

Rearranging the terms and using all the previous inequalities with (2.3), we have

$$s_{n+1} \leq (\Pi_{k=N}^n(1 - a_k)) s_N + (1 - \Pi_{k=N}^n(1 - a_k)) \frac{\epsilon}{2} + \sum_{k=N}^n \xi_k. \quad (2.4)$$

Similarly, we can write

$$t_{n+1} \leq (\Pi_{k=N}^n(1 - a_k)) t_N + (1 - \Pi_{k=N}^n(1 - a_k)) \frac{\epsilon}{2} + \sum_{k=N}^n \delta_k. \quad (2.5)$$

Adding (2.4) and (2.5), we obtain

$$(s_{n+1} + t_{n+1}) \leq (\Pi_{k=N}^n(1 - a_k)) (s_N + t_N) + (1 - \Pi_{k=N}^n(1 - a_k)) \epsilon + \sum_{k=N}^n \xi_k + \sum_{k=N}^n \delta_k. \quad (2.6)$$

Using conditions (ii) and (iii), we obtain

$$s_n + t_n \rightarrow 0, \quad \text{as } n \rightarrow \infty.$$

□

### 3. Phrasing of problem and iterative schemes

Let  $\tilde{\mathcal{A}}, \tilde{\mathcal{B}} : \hat{\mathcal{X}} \rightarrow \hat{\mathcal{X}}$  be single-valued mappings. Suppose that  $\mathcal{M}, \mathcal{N} : \hat{\mathcal{X}} \rightarrow CB(\hat{\mathcal{X}})$  are multi-valued mappings,  $Y_{\tilde{\mathcal{A}}, \rho}^{\mathcal{M}} : \hat{\mathcal{X}} \rightarrow \hat{\mathcal{X}}$  and  $C_{\tilde{\mathcal{B}}, \gamma}^{\mathcal{N}} : \hat{\mathcal{X}} \rightarrow \hat{\mathcal{X}}$  are generalized Yosida approximation operator and generalized Cayley operator, respectively. We will study the following system of variational inclusions involving generalized Yosida and Cayley operators.

Find  $\hat{p}, \hat{q} \in \hat{\mathcal{X}}$  such that

$$\begin{aligned} 0 &\in Y_{\tilde{\mathcal{A}}, \rho}^{\mathcal{M}}(\hat{p}) + \mathcal{M}(\hat{q}), \\ 0 &\in C_{\tilde{\mathcal{B}}, \gamma}^{\mathcal{N}}(\hat{q}) + \mathcal{N}(\hat{p}). \end{aligned} \quad (3.1)$$

If  $Y_{\tilde{\mathcal{A}}, \rho}^{\mathcal{M}}(\hat{p}) = 0 = C_{\tilde{\mathcal{B}}, \gamma}^{\mathcal{N}}(\hat{q})$ , then the problem (3.1) reduces to the system of variational inclusions, that is, find  $\hat{p}, \hat{q} \in \hat{\mathcal{X}}$  such that

$$\begin{aligned} 0 &\in \mathcal{M}(\hat{q}), \\ 0 &\in \mathcal{N}(\hat{p}). \end{aligned} \quad (3.2)$$

One can obtain many previously studied systems of variational inclusions from system (3.1).

The fixed point formulation of system (3.1) is given below.

**Lemma 3.1.** *The system of variational inclusions involving generalized Yosida and Cayley operators (3.1) has a solution  $\hat{p}, \hat{q} \in \hat{\mathcal{X}}$  if and only if the following system of fixed point equations is satisfied:*

$$\hat{q} = R_{\tilde{\mathcal{A}}, \rho}^{\mathcal{M}} \left[ \tilde{\mathcal{A}}(\hat{q}) - \rho Y_{\tilde{\mathcal{A}}, \rho}^{\mathcal{M}}(\hat{p}) \right], \quad (3.3)$$

$$\hat{p} = R_{\tilde{\mathcal{B}}, \gamma}^{\mathcal{N}} \left[ \tilde{\mathcal{B}}(\hat{p}) - \gamma C_{\tilde{\mathcal{B}}, \gamma}^{\mathcal{N}}(\hat{q}) \right]. \quad (3.4)$$

**Proof.** Proof is easy and hence omitted.  $\square$

Using Lemma 3.1, we suggest the following iterative scheme for solving system (3.1).

**Iterative Scheme 3.1.** For any  $\hat{p}_0, \hat{q}_0 \in \hat{\mathcal{X}}$ , compute sequences  $\{\hat{p}_n\}$  and  $\{\hat{q}_n\}$  by the following scheme:

$$\hat{q}_{n+1} = (1 - \alpha_n)\hat{q}_n + \alpha_n R_{\tilde{\mathcal{A}}, \rho}^{\mathcal{M}} \left[ A(\hat{q}_n) - \rho Y_{\tilde{\mathcal{A}}, \rho}^{\mathcal{M}}(\hat{p}_n) \right], \quad (3.5)$$

$$\hat{p}_{n+1} = (1 - \beta_n)\hat{p}_n + \beta_n R_{\tilde{\mathcal{B}}, \gamma}^{\mathcal{N}} \left[ \tilde{\mathcal{B}}(\hat{p}_n) - \gamma C_{\tilde{\mathcal{B}}, \gamma}^{\mathcal{N}}(\hat{q}_n) \right], \quad (3.6)$$

where  $n = 0, 1, 2, \dots$ ,  $\alpha_n, \beta_n \in [0, 1]$ ,  $\rho > 0$  and  $\gamma > 0$  are constants.

Equations (3.3) and (3.4) can be rewritten as

$$\hat{q} = R_{\tilde{\mathcal{A}}, \rho}^{\mathcal{M}} \left[ \frac{\tilde{\mathcal{A}}(\hat{q}) + \tilde{\mathcal{A}}(\hat{q})}{2} - \rho Y_{\tilde{\mathcal{A}}, \rho}^{\mathcal{M}}(\hat{p}) \right], \quad (3.7)$$

$$\text{and } \hat{p} = R_{\tilde{\mathcal{B}}, \gamma}^{\mathcal{N}} \left[ \frac{\tilde{\mathcal{B}}(\hat{p}) + \tilde{\mathcal{B}}(\hat{p})}{2} - \gamma C_{\tilde{\mathcal{B}}, \gamma}^{\mathcal{N}}(\hat{q}) \right]. \quad (3.8)$$

Based on (3.7) and (3.8), we suggest the following iterative scheme to solve the system (3.1).

**Iterative Scheme 3.2.** For any  $\hat{p}_0, \hat{q}_0 \in \hat{\mathcal{X}}$ , compute the sequences  $\{\hat{p}_{n+1}\}$  and  $\{\hat{q}_{n+1}\}$  by the recurrence relations:

$$\hat{q}_{n+1} = (1 - \alpha_n)\hat{q}_n + \alpha_n R_{\tilde{\mathcal{A}}, \rho}^{\mathcal{M}} \left[ \frac{\tilde{\mathcal{A}}(\hat{q}_n) + \tilde{\mathcal{A}}(\hat{q}_{n+1})}{2} - \rho Y_{\tilde{\mathcal{A}}, \rho}^{\mathcal{M}}(\hat{p}_{n+1}) \right], \quad (3.9)$$

$$\hat{p}_{n+1} = (1 - \beta_n)\hat{p}_n + \beta_n R_{\tilde{\mathcal{B}}, \gamma}^{\mathcal{N}} \left[ \frac{\tilde{\mathcal{B}}(\hat{p}_n) + \tilde{\mathcal{B}}(\hat{p}_{n+1})}{2} - \gamma C_{\tilde{\mathcal{B}}, \gamma}^{\mathcal{N}}(\hat{q}_{n+1}) \right]. \quad (3.10)$$

Where  $n = 0, 1, 2, \dots$ ,  $\alpha_n, \beta_n \in [0, 1]$ ,  $\rho > 0$  and  $\gamma > 0$  are constants.

We established the following inertial extrapolation scheme.

**Iterative Scheme 3.3.** For any  $\hat{p}_0, \hat{q}_0 \in \hat{\mathcal{X}}$ , compute the sequences  $\{\hat{p}_{n+1}\}$  and  $\{\hat{q}_{n+1}\}$  by the recurrence relations:

$$\hat{w}_n = \hat{q}_n + \gamma'_n(\hat{p}_n - \hat{p}_{n-1}), \quad (3.11)$$

$$\hat{q}_{n+1} = (1 - \alpha_n)\hat{q}_n + \alpha_n R_{\tilde{\mathcal{A}}, \rho}^{\mathcal{M}} \left[ \frac{\tilde{\mathcal{A}}(\hat{q}_n) + \tilde{\mathcal{A}}(\hat{w}_n)}{2} - \rho Y_{\tilde{\mathcal{A}}, \rho}^{\mathcal{M}}(\hat{w}_n) \right], \quad (3.12)$$

$$\hat{v}_n = \hat{p}_n + \gamma''_n(\hat{q}_n - \hat{q}_{n-1}), \quad (3.13)$$

$$\hat{p}_{n+1} = (1 - \beta_n)\hat{p}_n + \beta_n R_{\tilde{\mathcal{B}}, \gamma}^{\mathcal{N}} \left[ \frac{\tilde{\mathcal{B}}(\hat{p}_n) + \tilde{\mathcal{B}}(\hat{v}_n)}{2} - \gamma C_{\tilde{\mathcal{B}}, \gamma}^{\mathcal{N}}(\hat{v}_n) \right]. \quad (3.14)$$

Where  $\alpha_n, \beta_n, \gamma'_n, \gamma''_n \in [0, 1]$ ,  $\gamma'_n$  and  $\gamma''_n$  are the extrapolating terms for  $n \geq 1$ ,  $\rho > 0$  and  $\gamma > 0$  are constants.

## 4. Existence and convergence results

Existence and convergence results for the system (3.1) discussed below.

**Theorem 4.1.** *Let  $\hat{\mathcal{X}}$  be a real Banach space. Let  $\tilde{\mathcal{A}} : \hat{\mathcal{X}} \rightarrow \hat{\mathcal{X}}$  be the single-valued mappings such that  $\tilde{\mathcal{A}}$  is  $\lambda_{\tilde{\mathcal{A}}}$ -Lipschitz continuous, strongly accretive with constant  $r_1$ ;  $\tilde{\mathcal{B}}$  is  $\lambda_{\tilde{\mathcal{B}}}$ -Lipschitz continuous and strongly accretive with constant  $r_2$ . Let  $\mathcal{M}, \mathcal{N} : \hat{\mathcal{X}} \rightarrow 2^{\hat{\mathcal{X}}}$  be the multi-valued mappings such that  $\mathcal{M}$  is  $\tilde{\mathcal{A}}$ -accretive and  $\mathcal{N}$  is  $\tilde{\mathcal{B}}$ -accretive. Let  $R_{\tilde{\mathcal{A}},\rho}^{\mathcal{M}}, R_{\tilde{\mathcal{B}},\gamma}^{\mathcal{N}} : \hat{\mathcal{X}} \rightarrow \hat{\mathcal{X}}$  be the generalized resolvent operators such that  $R_{\tilde{\mathcal{A}},\rho}^{\mathcal{M}}$  is  $\frac{1}{r_1}$ -Lipschitz continuous and  $R_{\tilde{\mathcal{B}},\gamma}^{\mathcal{N}}$  is  $\frac{1}{r_2}$ -Lipschitz continuous. Let  $Y_{\tilde{\mathcal{A}},\rho}^{\mathcal{M}} : \hat{\mathcal{X}} \rightarrow \hat{\mathcal{X}}$  be the generalized Yosida approximation operator and  $C_{\tilde{\mathcal{B}},\gamma}^{\mathcal{N}} : \hat{\mathcal{X}} \rightarrow \hat{\mathcal{X}}$  be the generalized Cayley operator such that  $Y_{\tilde{\mathcal{A}},\rho}^{\mathcal{M}}$  is  $\lambda_Y$ -Lipschitz continuous and  $C_{\tilde{\mathcal{B}},\gamma}^{\mathcal{N}}$  is  $\lambda_C$ -Lipschitz continuous. Suppose that the following conditions are satisfied for  $\alpha_n, \beta_n, \gamma'_n, \gamma''_n \in [0, 1]$ , for all  $n \geq 1$  such that*

$$\frac{r_1 + \lambda_{\tilde{\mathcal{A}}}}{r_1} > 1, \quad \frac{r_2 + \lambda_{\tilde{\mathcal{B}}}}{r_2} > 1, \quad \sum_{n=1}^{\infty} \alpha_n = \infty, \quad \sum_{n=1}^{\infty} \beta_n = \infty, \quad (4.1)$$

$$\left. \begin{aligned} \sum_{n=1}^{\infty} \gamma'_n \left[ \xi(\theta_1) \|\hat{p}_n - \hat{p}_{n-1}\| \right] &< \infty \\ \sum_{n=1}^{\infty} \gamma''_n \left[ \xi(\theta_2) \|\hat{q}_n - \hat{q}_{n-1}\| \right] &< \infty \end{aligned} \right\}, \quad (4.2)$$

$$\limsup[\alpha_n \theta \rho \lambda_Y] \leq 0, \quad \limsup[\beta_n \theta' \gamma \lambda_C] \leq 0, \quad (4.3)$$

where  $\theta = \frac{1}{r_1}$ ,  $\theta' = \frac{1}{r_2}$ ,  $\lambda_Y = \frac{\lambda_{\tilde{\mathcal{A}}} r_1 + 1}{\rho r_1}$ ,  $\lambda_C = \frac{2 + \lambda_{\tilde{\mathcal{B}}} r_2}{r_2}$ , all the constants are positive and  $\gamma'_n, \gamma''_n$  are the extrapolating terms.

Then, the sequences  $\{\hat{p}_n\}$  and  $\{\hat{q}_n\}$  produced by scheme 3.1 converge strongly to the solution of system (3.1).

**Proof.** Let  $\hat{p}, \hat{q} \in \hat{\mathcal{X}}$  be the solution of system of variational inclusions involving generalized Yosida and Cayley operators (3.1). Using (3.7) and (3.8), we have

$$\hat{q}^* = (1 - \alpha_n) \hat{q}^* + \alpha_n R_{\tilde{\mathcal{A}},\rho}^{\mathcal{M}} \left[ \frac{\tilde{\mathcal{A}}(\hat{q}^*) + \tilde{\mathcal{A}}(\hat{q}^*)}{2} - \rho Y_{\tilde{\mathcal{A}},\rho}^{\mathcal{M}}(\hat{p}^*) \right], \quad (4.4)$$

$$\hat{p}^* = (1 - \beta_n) \hat{p}^* + \beta_n R_{\tilde{\mathcal{B}},\gamma}^{\mathcal{N}} \left[ \frac{\tilde{\mathcal{B}}(\hat{p}^*) + \tilde{\mathcal{B}}(\hat{p}^*)}{2} - \gamma C_{\tilde{\mathcal{B}},\gamma}^{\mathcal{N}}(\hat{q}^*) \right], \quad (4.5)$$

where  $\alpha_n, \beta_n \in [0, 1]$ , for all  $n \geq 1$ . Using (3.12), (4.4) and (3.11) and Lipschitz continuity of the generalized resolvent operator  $R_{\tilde{\mathcal{A}},\rho}^{\mathcal{M}}$ , we evaluate

$$\begin{aligned} \|\hat{q}_{n+1} - \hat{q}^*\| &= \left\| \left\{ (1 - \alpha_n) \hat{q}_n + \alpha_n R_{\tilde{\mathcal{A}},\rho}^{\mathcal{M}} \left[ \frac{\tilde{\mathcal{A}}(\hat{q}_n) + \tilde{\mathcal{A}}(\hat{w}_n)}{2} - \rho Y_{\tilde{\mathcal{A}},\rho}^{\mathcal{M}}(\hat{w}_n) \right] \right\} \right. \\ &\quad \left. - \left\{ (1 - \alpha_n) \hat{q}^* + \alpha_n R_{\tilde{\mathcal{A}},\rho}^{\mathcal{M}} \left[ \frac{\tilde{\mathcal{A}}(\hat{q}^*) + \tilde{\mathcal{A}}(\hat{q}^*)}{2} - \rho Y_{\tilde{\mathcal{A}},\rho}^{\mathcal{M}}(\hat{p}^*) \right] \right\} \right\| \end{aligned}$$

$$\begin{aligned}
&\leq (1 - \alpha_n) \|\hat{q}_n - \hat{q}^*\| + \alpha_n \theta \left\| \left[ \frac{\tilde{\mathcal{A}}(\hat{q}_n) + \tilde{\mathcal{A}}(\hat{w}_n)}{2} - \rho Y_{\tilde{\mathcal{A}}, \rho}^{\mathcal{M}}(\hat{w}_n) \right] \right. \\
&\quad \left. - \left[ \frac{\tilde{\mathcal{A}}(\hat{q}^*) + \tilde{\mathcal{A}}(\hat{p}^*)}{2} - \rho Y_{\tilde{\mathcal{A}}, \rho}^{\mathcal{M}}(\hat{p}^*) \right] \right\| \\
&\leq (1 - \alpha_n) \|\hat{q}_n - \hat{q}^*\| + \frac{\alpha_n \theta}{2} \|\tilde{\mathcal{A}}(\hat{q}_n) - \tilde{\mathcal{A}}(\hat{q}^*)\| \\
&\quad + \frac{\alpha_n \theta}{2} \|\tilde{\mathcal{A}}(\hat{w}_n) - \tilde{\mathcal{A}}(\hat{q}^*)\| + \alpha_n \theta \rho \|Y_{\tilde{\mathcal{A}}, \rho}^{\mathcal{M}}(\hat{w}_n) - Y_{\tilde{\mathcal{A}}, \rho}^{\mathcal{M}}(\hat{p}^*)\|. \quad (4.6)
\end{aligned}$$

Using the Lipschitz continuity of the mapping  $\tilde{\mathcal{A}}$  and generalized Yosida approximation operator  $Y_{\tilde{\mathcal{A}}, \rho}^{\mathcal{M}}$ , from (4.6), we obtain

$$\begin{aligned}
\|\hat{q}_{n+1} - \hat{q}^*\| &\leq (1 - \alpha_n) \|\hat{q}_n - \hat{q}^*\| + \frac{\alpha_n \theta}{2} \lambda_{\tilde{\mathcal{A}}} \|\hat{q}_n - \hat{q}^*\| \\
&\quad + \frac{\alpha_n \theta}{2} \lambda_{\tilde{\mathcal{A}}} \|\hat{w}_n - \hat{q}^*\| + \alpha_n \theta \rho \lambda_Y \|\hat{w}_n - \hat{p}^*\|. \quad (4.7)
\end{aligned}$$

Applying (3.11), we can write

$$\begin{aligned}
\|\hat{w}_n - \hat{q}^*\| &= \|\hat{q}_n + \gamma'_n(\hat{p}_n - \hat{p}_{n-1}) - \hat{q}^*\| \\
&\leq \|\hat{q}_n - \hat{q}^*\| + \gamma'_n \|\hat{p}_n - \hat{p}_{n-1}\|, \quad (4.8)
\end{aligned}$$

and

$$\|\hat{w}_n - \hat{p}^*\| \leq \|\hat{q}_n - \hat{p}^*\| + \gamma'_n \|\hat{p}_n - \hat{p}_{n-1}\|. \quad (4.9)$$

Making use of (4.8) and (4.9), (4.7) becomes

$$\begin{aligned}
\|\hat{q}_{n+1} - \hat{q}^*\| &\leq (1 - \alpha_n) \|\hat{q}_n - \hat{q}^*\| + \frac{\alpha_n \theta}{2} \lambda_{\tilde{\mathcal{A}}} \|\hat{q}_n - \hat{q}^*\| + \frac{\alpha_n \theta}{2} \lambda_{\tilde{\mathcal{A}}} \left[ \|\hat{q}_n - \hat{q}^*\| \right. \\
&\quad \left. + \gamma'_n \|\hat{p}_n - \hat{p}_{n-1}\| \right] + \alpha_n \theta \rho \lambda_Y \left[ \|\hat{q}_n - \hat{p}^*\| + \gamma'_n \|\hat{p}_n - \hat{p}_{n-1}\| \right] \\
&\leq \left[ (1 - \alpha_n) + \frac{\alpha_n \theta}{2} \lambda_{\tilde{\mathcal{A}}} + \frac{\alpha_n \theta}{2} \lambda_{\tilde{\mathcal{A}}} \right] \|\hat{q}_n - \hat{q}^*\| \\
&\quad + \frac{\alpha_n \theta}{2} \lambda_{\tilde{\mathcal{A}}} \gamma'_n \|\hat{p}_n - \hat{p}_{n-1}\| + \alpha_n \theta \rho \lambda_Y \|\hat{q}_n - \hat{p}^*\| \\
&\quad + \alpha_n \theta \rho \lambda_Y \gamma'_n \|\hat{p}_n - \hat{p}_{n-1}\| \\
&\leq \left[ (1 - \alpha_n) + \alpha_n \theta \lambda_{\tilde{\mathcal{A}}} \right] \|\hat{q}_n - \hat{q}^*\| \\
&\quad + \left[ \left( \frac{\alpha_n \theta}{2} \lambda_{\tilde{\mathcal{A}}} + \alpha_n \theta \rho \lambda_Y \right) \gamma'_n \right] \|\hat{p}_n - \hat{p}_{n-1}\| \\
&\quad + \alpha_n \theta \rho \lambda_Y \|\hat{q}_n - \hat{p}^*\|. \quad (4.10)
\end{aligned}$$

Using (3.14), (4.5) and using the Lipschitz continuity of generalized resolvent operator  $R_{\tilde{\mathcal{B}}, \gamma}^{\mathcal{M}}$ , we evaluate

$$\begin{aligned}
\|\hat{p}_{n+1} - \hat{p}^*\| &= \left\| \left\{ (1 - \beta_n) \hat{p}_n + \beta_n R_{\tilde{\mathcal{B}}, \gamma}^{\mathcal{N}} \left[ \frac{\tilde{\mathcal{B}}(\hat{p}_n) + \tilde{\mathcal{B}}(\hat{v}_n)}{2} - \gamma C_{\tilde{\mathcal{B}}, \gamma}^{\mathcal{N}}(\hat{v}_n) \right] \right\} \right. \\
&\quad \left. - \left\{ (1 - \beta_n) \hat{p}^* + \beta_n R_{\tilde{\mathcal{B}}, \gamma}^{\mathcal{N}} \left[ \frac{\tilde{\mathcal{B}}(\hat{p}^*) + \tilde{\mathcal{B}}(\hat{q}^*)}{2} - \gamma C_{\tilde{\mathcal{B}}, \gamma}^{\mathcal{N}}(\hat{q}^*) \right] \right\} \right\|
\end{aligned}$$



$$\begin{aligned}
&\leq (1 - \beta_n) \|\hat{p}_n - \hat{p}^*\| + \beta_n \theta' \left\| \left[ \frac{\tilde{\mathcal{B}}(\hat{p}_n) + \tilde{\mathcal{B}}(\hat{v}_n)}{2} - \gamma C_{\tilde{\mathcal{B}}, \gamma}^{\mathcal{N}}(\hat{v}_n) \right] \right. \\
&\quad \left. - \left[ \frac{\tilde{\mathcal{B}}(\hat{p}^*) + \tilde{\mathcal{B}}(\hat{q}^*)}{2} - \gamma C_{\tilde{\mathcal{B}}, \gamma}^{\mathcal{N}}(\hat{q}^*) \right] \right\| \\
&\leq (1 - \beta_n) \|\hat{p}_n - \hat{p}^*\| + \frac{\beta_n \theta'}{2} \|\tilde{\mathcal{B}}(\hat{p}_n) - \tilde{\mathcal{B}}(\hat{p}^*)\| \\
&\quad + \frac{\beta_n \theta'}{2} \|\tilde{\mathcal{B}}(\hat{v}_n) - \tilde{\mathcal{B}}(\hat{p}^*)\| + \beta_n \theta' \gamma \left\| C_{\tilde{\mathcal{B}}, \gamma}^{\mathcal{N}}(\hat{v}_n) - C_{\tilde{\mathcal{B}}, \gamma}^{\mathcal{N}}(\hat{q}^*) \right\|.
\end{aligned} \tag{4.11}$$

Using the Lipschitz continuity of the mapping  $\tilde{\mathcal{B}}$  and generalized Cayley operator  $C_{\tilde{\mathcal{B}}, \gamma}^{\mathcal{N}}$ , from (4.11), we obtain

$$\begin{aligned}
\|\hat{p}_{n+1} - \hat{p}^*\| &\leq (1 - \beta_n) \|\hat{p}_n - \hat{p}^*\| + \frac{\beta_n \theta'}{2} \lambda_{\tilde{\mathcal{B}}} \|\hat{p}_n - \hat{p}^*\| \\
&\quad + \frac{\beta_n \theta'}{2} \lambda_{\tilde{\mathcal{B}}} \|\hat{v}_n - \hat{p}^*\| + \beta_n \theta' \gamma \lambda_C \|\hat{v}_n - \hat{q}^*\|.
\end{aligned} \tag{4.12}$$

Applying (3.13), we can write

$$\begin{aligned}
\|\hat{v}_n - \hat{p}^*\| &= \|\hat{p}_n + \gamma_n''(\hat{q}_n - \hat{q}_{n-1}) - \hat{p}^*\| \\
&\leq \|\hat{p}_n - \hat{p}^*\| + \gamma_n'' \|\hat{q}_n - \hat{q}_{n-1}\|,
\end{aligned} \tag{4.13}$$

and

$$\|\hat{v}_n - \hat{q}^*\| \leq \|\hat{p}_n - \hat{q}^*\| + \gamma_n'' \|\hat{q}_n - \hat{q}_{n-1}\|. \tag{4.14}$$

Making use of (4.13) and (4.14), (4.12) becomes

$$\begin{aligned}
\|\hat{p}_{n+1} - \hat{p}^*\| &\leq (1 - \beta_n) \|\hat{p}_n - \hat{p}^*\| + \frac{\beta_n \theta'}{2} \lambda_{\tilde{\mathcal{B}}} \|\hat{p}_n - \hat{p}^*\| + \frac{\beta_n \theta'}{2} \lambda_{\tilde{\mathcal{B}}} [\|\hat{p}_n - \hat{p}^*\| \\
&\quad + \gamma_n'' \|\hat{q}_n - \hat{q}_{n-1}\|] + \beta_n \theta' \gamma \lambda_C [\|\hat{p}_n - \hat{q}^*\| + \gamma_n'' \|\hat{q}_n - \hat{q}_{n-1}\|] \\
&\leq [(1 - \beta_n) + \beta_n \theta' \lambda_{\tilde{\mathcal{B}}}] \|\hat{p}_n - \hat{p}^*\| \\
&\quad + \left[ \left( \frac{\beta_n \theta'}{2} \lambda_{\tilde{\mathcal{B}}} + \beta_n \theta' \gamma \lambda_C \right) \gamma_n'' \right] \|\hat{q}_n - \hat{q}_{n-1}\| \\
&\quad + \beta_n \theta' \gamma \lambda_C \|\hat{p}_n - \hat{q}^*\|.
\end{aligned} \tag{4.15}$$

Adding (4.10) and (4.15), we obtain

$$\begin{aligned}
\|\hat{q}_{n+1} - \hat{q}^*\| + \|\hat{p}_{n+1} - \hat{p}^*\| &\leq [(1 - \alpha_n) + \alpha_n \theta \lambda_{\tilde{\mathcal{B}}}] \|\hat{q}_n - \hat{q}^*\| \\
&\quad + \left[ \left( \frac{\alpha_n \theta}{2} \lambda_{\tilde{\mathcal{B}}} + \alpha_n \theta \rho \lambda_Y \right) \gamma_n' \right] \|\hat{p}_n - \hat{p}_{n-1}\| \\
&\quad + \alpha_n \theta \rho \lambda_Y \|\hat{q}_n - \hat{p}^*\| \\
&\quad + [(1 - \beta_n) + \beta_n \theta' \lambda_B] \|\hat{p}_n - \hat{p}^*\| \\
&\quad + \left[ \left( \frac{\beta_n \theta'}{2} \lambda_{\tilde{\mathcal{B}}} + \beta_n \theta' \gamma \lambda_C \right) \gamma_n'' \right] \|\hat{q}_n - \hat{q}_{n-1}\|
\end{aligned}$$

$$\begin{aligned}
& +\beta_n\theta'\gamma\lambda_C\|\widehat{p}_n-\widehat{q}^*\| \\
& = \left[ (1-\alpha_n(1-\theta\lambda_{\widetilde{\mathcal{A}}}))\|\widehat{q}_n-\widehat{q}^*\| \right. \\
& \quad + \left[ (1-\beta_n(1-\theta'\lambda_{\widetilde{\mathcal{B}}}))\|\widehat{p}_n-\widehat{p}^*\| \right. \\
& \quad + \xi(\theta_1)\gamma'_n\|\widehat{p}_n-\widehat{p}_{n-1}\| + \xi(\theta_2)\gamma''_n\|\widehat{q}_n-\widehat{q}_{n-1}\| \\
& \quad + \alpha_n\theta\rho\lambda_Y\|\widehat{q}_n-\widehat{p}^*\| + \beta_n\theta'\gamma\lambda_C\|\widehat{p}_n-\widehat{q}^*\| \\
& = \xi(\widehat{\theta})[\|\widehat{q}_n-\widehat{q}^*\| + \|\widehat{p}_n-\widehat{p}^*\|] \\
& \quad + \xi(\theta_1)\gamma'_n\|\widehat{p}_n-\widehat{p}_{n-1}\| + \xi(\theta_2)\gamma''_n\|\widehat{q}_n-\widehat{q}_{n-1}\| \\
& \quad + \alpha_n\theta\rho\lambda_Y\|\widehat{q}_n-\widehat{p}^*\| + \beta_n\theta'\gamma\lambda_C\|\widehat{p}_n-\widehat{q}^*\|,
\end{aligned} \tag{4.16}$$

where

$$\begin{aligned}
\xi(\widehat{\theta}) &= \text{map}\{[1-\alpha_n(1-\theta\lambda_{\widetilde{\mathcal{A}}})], [1-\beta_n(1-\theta'\lambda_{\widetilde{\mathcal{B}}})]\}, \\
\xi(\theta_1) &= \frac{\alpha_n\theta}{2}\lambda_{\widetilde{\mathcal{A}}} + \alpha_n\theta\rho\lambda_Y, \quad \xi(\theta_2) = \frac{\beta_n\theta}{2}\lambda_{\widetilde{\mathcal{B}}} + \beta_n\theta'\gamma\lambda_C.
\end{aligned}$$

By condition (4.1),

$$1-\theta\lambda_{\widetilde{\mathcal{A}}} < 1, \quad 1-\theta'\lambda_{\widetilde{\mathcal{B}}} < 1, \quad \sum_{n=1}^{\infty}\alpha_n = \infty \quad \text{and} \quad \sum_{n=1}^{\infty}\beta_n = \infty.$$

By condition (4.2),

$$\left. \begin{aligned} \sum_{n=1}^{\infty}\gamma'_n\left[\xi(\theta_1)\|\widehat{p}_n-\widehat{p}_{n-1}\|\right] &< \infty \\ \sum_{n=1}^{\infty}\gamma''_n\left[\xi(\theta_2)\|\widehat{q}_n-\widehat{q}_{n-1}\|\right] &< \infty \end{aligned} \right\}.$$

Also, applying condition (4.3), we have

$$\limsup[\alpha_n\theta\rho\lambda_Y] \leq 0, \quad \limsup[\beta_n\theta'\gamma\lambda_C] \leq 0.$$

Applying Lemma 2.1, we conclude that  $\widehat{p}_n \rightarrow \widehat{p}$  and  $\widehat{q}_n \rightarrow \widehat{q}$ , as  $n \rightarrow \infty$ . This completes the proof.  $\square$

The following numerical example is constructed showing that all the conditions of Theorem 4.1 are satisfied. We also establish the convergence graph and computational table for illustration.

**Example 4.1.** Let  $\widehat{\mathcal{X}} = \mathbb{R}$  with usual inner product and norm. Let  $\widetilde{\mathcal{A}}, \widetilde{\mathcal{B}} : \widehat{\mathcal{X}} \rightarrow \widehat{\mathcal{X}}$  be the mappings such that  $\widetilde{\mathcal{A}}(\widehat{p}) = (\frac{3\widehat{p}}{2})$  and  $\widetilde{\mathcal{B}}(\widehat{p}) = (\frac{\widehat{p}}{6})$  and the multi-valued mappings  $\mathcal{M}, \mathcal{N} : \widehat{\mathcal{X}} \rightarrow CB(\widehat{\mathcal{X}})$  defined by  $\mathcal{M}(\widehat{p}) = \{\frac{2\widehat{p}}{5}\}$  and  $\mathcal{N}(\widehat{p}) = \{\frac{3\widehat{p}}{7}\}$ .

(i)  $\widetilde{\mathcal{A}}$  is  $\lambda_{\widetilde{\mathcal{A}}}$ -Lipschitz and  $r_1$ -strongly accretive

$$\|\widetilde{\mathcal{A}}(\widehat{p}) - \widetilde{\mathcal{A}}(\widehat{q})\| = \left\| \frac{3\widehat{p}}{2} - \frac{3\widehat{q}}{2} \right\| = \frac{3}{2}\|\widehat{p} - \widehat{q}\| \leq 2\|\widehat{p} - \widehat{q}\|,$$

that is,  $\tilde{\mathcal{A}}$  is  $\lambda_{\tilde{\mathcal{A}}} = 2$ -Lipschitz continuous.

$\tilde{\mathcal{A}}$  is  $r_1$ -strongly accretive.

$$\langle \tilde{\mathcal{A}}(\hat{p}) - \tilde{\mathcal{A}}(\hat{q}), \hat{p} - \hat{q} \rangle = \langle \frac{3\hat{p}}{2} - \frac{3\hat{q}}{2}, \hat{p} - \hat{q} \rangle = \frac{3}{2} \|\hat{p} - \hat{q}\|^2 \geq \frac{2}{3} \|\hat{p} - \hat{q}\|^2,$$

that is,  $\tilde{\mathcal{A}}$  is  $r_1 = \frac{2}{3}$  strongly accretive.

(ii) Similarly, one can prove that  $\tilde{\mathcal{B}}$  is  $\lambda_{\tilde{\mathcal{B}}} = \frac{8}{6}$ -Lipschitz continuous and  $r_2 = \frac{3}{4}$ -strongly accretive.

(iii) For  $\rho = \gamma = 1$ , it is easy to show that  $\mathcal{M}$  is  $\tilde{\mathcal{A}}$ -accretive mapping and  $\mathcal{N}$  is  $\tilde{\mathcal{B}}$ -accretive mapping.

(iv) For  $\rho = \gamma = 1$ , we calculate

$$R_{\tilde{\mathcal{A}},\rho}^{\mathcal{M}}(\hat{p}) = [\tilde{\mathcal{A}} + \rho\mathcal{M}]^{-1}(\hat{p}) = \left(\frac{10\hat{p}}{19}\right),$$

$$R_{\tilde{\mathcal{B}},\gamma}^{\mathcal{N}}(\hat{p}) = [\tilde{\mathcal{B}} + \gamma\mathcal{N}]^{-1}(\hat{p}) = \left(\frac{42\hat{p}}{67}\right).$$

The Lipschitz continuity of  $R_{\tilde{\mathcal{A}},\rho}^{\mathcal{M}}$  and  $R_{\tilde{\mathcal{B}},\gamma}^{\mathcal{N}}$  is calculated below:

$$\|R_{\tilde{\mathcal{A}},\rho}^{\mathcal{M}}(\hat{p}) - R_{\tilde{\mathcal{A}},\rho}^{\mathcal{M}}(\hat{q})\| = \left\| \frac{10\hat{p}}{19} - \frac{10\hat{q}}{19} \right\| \leq \frac{3}{2} \|\hat{p} - \hat{q}\|.$$

Similarly,

$$\|R_{\tilde{\mathcal{B}},\gamma}^{\mathcal{N}}(\hat{p}) - R_{\tilde{\mathcal{B}},\gamma}^{\mathcal{N}}(\hat{q})\| = \left\| \frac{42\hat{p}}{67} - \frac{42\hat{q}}{67} \right\| \leq \frac{4}{3} \|\hat{p} - \hat{q}\|,$$

that is,  $R_{\tilde{\mathcal{A}},\rho}^{\mathcal{M}}$  is  $\frac{1}{(2/3)}$ -Lipschitz continuous and  $R_{\tilde{\mathcal{B}},\gamma}^{\mathcal{N}}$  is  $\frac{1}{(3/4)}$ -Lipschitz continuous.

(v) The generalized Yosida approximation operator and generalized Cayley operator are calculated as:

$$Y_{\tilde{\mathcal{A}},\rho}^{\mathcal{M}}(\hat{p}) = \frac{1}{\rho} [\tilde{\mathcal{A}} - R_{\tilde{\mathcal{A}},\rho}^{\mathcal{M}}](\hat{p}) = \left(\frac{37\hat{p}}{38}\right),$$

$$C_{\tilde{\mathcal{B}},\gamma}^{\mathcal{N}}(\hat{p}) = [2R_{\tilde{\mathcal{B}},\gamma}^{\mathcal{N}} - \tilde{\mathcal{B}}](\hat{p}) = \left(\frac{35\hat{p}}{402}\right).$$

$$\text{Also, } \|Y_{\tilde{\mathcal{A}},\rho}^{\mathcal{M}}(\hat{p}) - Y_{\tilde{\mathcal{A}},\rho}^{\mathcal{M}}(\hat{q})\| = \left\| \frac{37\hat{p}}{38} - \frac{37\hat{q}}{38} \right\|$$

$$\leq \frac{75}{76} \|\hat{p} - \hat{q}\|,$$

that is,  $Y_{\tilde{\mathcal{A}},\rho}^{\mathcal{M}}$  is  $\lambda_Y = \frac{75}{76}$ -Lipschitz continuous. And

$$\|C_{\tilde{\mathcal{B}},\gamma}^{\mathcal{N}}(\hat{p}) - C_{\tilde{\mathcal{B}},\gamma}^{\mathcal{N}}(\hat{q})\| = \left\| \frac{35\hat{p}}{402} - \frac{35\hat{q}}{402} \right\| \leq \frac{13}{134} \|\hat{p} - \hat{q}\|,$$

that is,  $C_{\tilde{\mathcal{B}},\gamma}^{\mathcal{N}}$  is  $\lambda_C = \frac{13}{134}$ -Lipschitz continuous.

- (vi) For  $\alpha_n = 1 - \frac{1}{n}$ ,  $\beta_n = 1 - \frac{1}{n+2}$ , we compute the sequence  $\hat{p}_n$  and  $\hat{q}_n$  by the Iterative scheme 3.1, in the following way:

$$\begin{aligned}\hat{q}_{n+1} &= \left(\frac{1}{n}\right)\hat{q}_n + \frac{10}{19}\left(\frac{n-1}{n}\right)\left(\frac{3\hat{q}_n}{2} - \frac{37\hat{p}_n}{38}\right), \\ \hat{p}_{n+1} &= \left(\frac{1}{n+2}\right)\hat{p}_n + \frac{42(n+1)}{67(n+2)}\left(\frac{7\hat{p}_n}{6} - \frac{35\hat{q}_n}{402}\right).\end{aligned}$$

- (vii) It is easy to check that condition (4.1), (4.2) and (4.3) of Theorem 4.1 are satisfied. Thus, system (3.1) admits a solution.




Figure 1. Convergence Graph for different initial values of  $\hat{p}$  and  $\hat{q}$ .

## 5. Conclusions

Due to applications of Yosida approximation operator and Cayley operator in contemporary science, this paper is centered on solving a system of variational inclusions involving the generalized Yosida and the Cayley operators in real Banach space. The solution to our problem has been achieved by developing an inertial extrapolation scheme, although several other schemes have also been developed. It is well-known that the inertial extrapolation scheme provides a faster rate of convergence.

We remark that scientists of other discipline may use our results for practical and applications purposes.

**Table 1.** Computational Table showing the output for different initial values of  $\hat{p}$  and  $\hat{q}$ .

No. of Iteration	For $\hat{p}_0 = 1$ $\hat{p}_n$	For $\hat{q}_0 = -1$ . $\hat{q}_n$	For $\hat{p}_0 = -2$ $\hat{p}_n$	For $\hat{q}_0 = 2$ $\hat{q}_n$
n=1	1	-1	-2	2
n=2	0.56867229	-2.3285738	-1.1438657	4.6506265
n=3	0.25954374	-2.2475433	-0.5439893	4.4687547
n=4	0.10520368	-1.39417701	-0.229106768	2.7665369
n=5	0.03907625	-0.64115197	-0.085211068	1.2730781
n=10	6.3088058e-05	-0.0008671193	-0.0001278830	0.0085015
n=15	7.8936310e-09	-7.57314208e-08	-1.58118674e-08	1.5119973e-07
n=20	1.5590855e-13	-1.1832846e-12	-3.116451e-13	2.363213e-12
n=25	8.029934e-19	-5.2264155e-18	-1.60430115e-18	1.0439134e-17
n=30	1.447128e-24	-8.479071e-24	-2.8908184e-24	1.693656e-23
n=35	1.1019388e-30	-5.990450e-30	-2.2011664e-30	1.1965819e-29
n=40	4.0400406e-37	-2.07791406e-36	-8.07003053e-37	4.1506175e-36
n=45	7.848087e-44	-3.8693066e-43	-1.567657e-43	7.7289247e-43
n=50	8.6912112e-51	-4.1449943e-50	-1.736069e-50	8.27961e-50

## Acknowledgements

All authors are thankful to anonymous referees and editor of Journal of Applied Analysis and Computation for their valuable comments which improve this paper a lot.

## References

- [1] R. Ahmad, I. Ali, X. B. Li and M. Ishtyak, *System of multi-valued mixed variational inclusions with XOR-operation in real ordered uniformly smooth Banach spaces*, Mathematics, 2019, 7(1027).
- [2] R. Ahmad, I. Ali, M. Rahaman and M. Ishtyak, *Cayley inclusion problem with its corresponding generalized resolvent equation problem in uniformly smooth Banach spaces*, Appl. Anal., 2022, 101, 1354–1368.
- [3] R. Ahmad, M. Ishtyak, A. K. Rajpoot and Y. Wang, *Solving system of mixed variational inclusions involving generalized Cayley operator and generalized Yosida approximation operator with error terms in  $q$ -uniformly smooth space*, Mathematics, 2022, 10, 4131.
- [4] I. Ahmad, J. C. Yao, R. Ahmad and M. Ishtyak, *System of Yosida inclusions involving XOR-operation*, J. Nonlinear Convex Anal., 2017, 18, 831–845.
- [5] B. Ali, A. A. Adam and A. Adamu, *An accelerated algorithm involving quasi- $\phi$ -nonexpansive operators for solving split problems*, Journal of Nonlinear Modeling and Analysis, 2023, 5(1), 54–72.
- [6] S. Althubiti and A. Mennouni, *A novel-projection method for Cayley-type systems of singular integro-differential equations*, Mathematics, 2022, 10, 2694.

- [7] Q. H. Ansari and J. C. Yao, *A fixed point theorem and its applications to system of variational inequalities*, Bull. Austral. Math. Soc., 1999, 59, 433–442.
- [8] C. L. Byrne, *A unified treatment of some iterative algorithm in signal processing and image reconstruction*, Inverse Problem, 2004, 20, 103–120.
- [9] A. Cayley, *Sur quelques propriétés des déterminants gauches*, J. Reine Angew. Math., 1846, 1846(32), 119–123.
- [10] L. C. Ceng, S. M. Guu and J. C. Yao, *Hybrid viscosity CQ method for finding a common solution of a variational inequality, a general system of variational inequalities and a fixed point theorem*, Fixed Point Theory Appl., 2013, 313, 1–25.
- [11] L. C. Ceng, A. Latif and J. C. Yao, *On solutions of a system of variational inequalities and fixed point problems in Banach spaces*, Fixed Point Theory Appl., 2013, 176, 1–34.
- [12] L. C. Ceng, S. Plubtring, M. M. Wong and J. C. Yao, *System of variational inequalities with constraints of mixed equilibria, variational inequalities, convex minimization and fixed point problems*, J. Nonlinear Convex Anal., 2015, 16, 385–421.
- [13] L. C. Ceng, C. Y. Wang and J. C. Yao, *Strong convergence theorems by a relaxed extra gradient method for a general system of variational inequalities*, Math. Methods Oper. Res., 2008, 67, 375–390.
- [14] A. De, *Hille-Yosida Theorem and Some Applications*, Ph.D. Thesis, Department of Mathematics and its Applications, Central European University, Budapest, Hungary, 2015.
- [15] X. Fang and M. Huang, *The monotonicity of the linear complementarity problem*, Journal of Nonlinear Modeling and Analysis, 2022, 4(2), 161–170.
- [16] Y. P. Fang and N. J. Huang, *H-accretive operators and resolvent operator technique for solving variational inclusions in Banach spaces*, Appl. Math. Lett., 2004, 17, 647–653.
- [17] J. Feng, W. Li, H. Chen and Y. Chen, *A class of differential inverse variational inequalities in finite dimensional spaces*, Journal of Applied Analysis and Computation, 2018, 8(6), 1664–1678.
- [18] G. Helmberg, *Introduction to Spectral Theory in Hilbert Space: The Cayley Transform*, North-Holland Series in Applied Mathematics and Mechanics, Applied Mathematics and Mechanics # 6, North-Holland Publishing Company: Amsterdam, The Netherlands, 1969.
- [19] J. Iqbal, A. K. Rajpoot, M. Islam, R. Ahmad and Y. Wang, *System of generalized variational inclusions involving Cayley operators and XOR-operation in  $q$ -uniformly smooth Banach spaces*, Mathematics, 2022, 10, 2837.
- [20] S. Jabeen, M. A. Noor and K. I. Noor, *Inertial iterative methods for general quasi-variational inequalities and dynamical systems*, J. Math. Anal., 2020, 11, 14–29.
- [21] C. Li and S. Wu, *A two-step matrix splitting method for the mixed linear complementarity problem*, Journal of Applied Analysis and Computation, 2024, 14(5), 2750–2761.

- [22] J. S. Pang, *Asymmetric variational inequality problems over product of sets: Applications and iterative methods*, Math. Program, 1985, 31, 206–219.
- [23] B. T. Polyak, *Some methods of speeding up the convergence of iterative methods*, USSR Comput. Math. Phys., 1964, 4, 1–17.
- [24] X. Qin, S. S. Chang, Y. J. Cho and S. M. Kang, *Approximation of solutions to a system of variational inclusions in Banach spaces*, J. Inequal. Appl., 2010, 2010, 916806.
- [25] B. Tan and S. Y. Cho, *An inertial Mann-like algorithm for fixed points of non-expansive mappings in Hilbert spaces*, J. Appl. Numer. Optim., 2020, 2, 335–351.
- [26] B. Tan and S. Li, *Strong convergence of inertial Mann algorithm for solving hierarchical fixed point problems*, J. Nonlinear Var. Anal., 2020, 4, 337–355.
- [27] W. Y. Yan, Y. P. Fang and N. J. Huang, *A new system of set-valued variational inclusions with  $H$ -monotone operators*, Math. Inequal. Appl., 2005, 8, 537–546.
- [28] C. Zhu, J. Chen, J. Chen, C. Chen and H. Huang, *A new generalization of  $F$ -metric spaces and some fixed point theorems and an application*, J. Appl. Anal. Comput., 2021, 11, 2649–2663.