BIFURCATIONS OF LIMIT CYCLES IN A CLASS OF QUARTIC PLANAR VECTOR FIELDS

Yuan Chang¹ and Liqin Zhao^{1,†}

Abstract This paper studies the number and the distribution of limit cycles of a class of planar quartic vector fields

$$\dot{x} = -y(ay^2 - 1) + \varepsilon p(x, y), \quad \dot{y} = x(ax^2 - \sqrt{ax} - 1) + \varepsilon q(x, y),$$

where a > 0, $0 < \varepsilon \ll 1$, p(x, y) and q(x, y) are polynomials in (x, y) of the degree 4. By the bifurcation theory and qualitative analysis, we obtain four new configurations of limit cycles, two of which can have at least 12 limit cycles.

Keywords Quartic system, distribution of limit cycles, Hopf bifurcation, homoclinic bifurcation, heteroclinic bifurcation.

MSC(2010) 34C05, 34C07.

1. Introduction and the main results

The second part of well-known Hilbert's 16th problem is to find an upper bound for the number of the limit cycles and their distributions of the planar polynomial system

$$\dot{x} = P(x, y), \quad \dot{y} = Q(x, y) \tag{1.1}$$

of the degree n. The maximum number of limit cycles of system (1.1) is denoted by H(n), which is known as the Hilbert number.

People usually study H(n) by bifurcation theory. They usually find limit cycles by Hopf bifurcation, quadruple transformation, detection functions, Poincaré bifurcation [1, 3-5, 7, 8, 15-20]. They also discuss the limit cycles bifurcating from polycycles such as homoclinic loop, double-homoclinic loop, heteroclinic loop, and so on [6, 8-10, 12-14, 18, 21-23]. The authors in [11] find more limit cycles increasing with n by investigating a property T(n) (see Definition 2.1 in [11]) on the rectangle $x_0 \leq x \leq x_{n+1}, y_0 \leq y \leq y_{n+1}$, and obtain the least increasing rate of H(n) with respect n.

The method of finding limit cycles near double-homoclinic loops or heteroclinic loops given in [9] and [14] contains three main steps:

(i) finding discriminate values to determine the existence and stability of homoclinic loops, double-homoclinic loops;

[†]The corresponding author.

 $^{^1\}mathrm{School}$ of Mathematical Sciences, Beijing Normal University, Beijing 100875, China

Email: yuanchang@mail.bnu.edu.cn(Y. Chang),

zhaoliqin@bnu.edu.cn(L. Zhao)

(ii) varying parameters to change the stability of these loops to create limit cycles near the loops;

(iii) breaking the homoclinic loops to find medium limit cycles.

Let us pay attentions to the distribution of limit cycles for quartic systems.

(i) The configuration of 13 limit cycles: [21] obtained 4 different configurations of 13 limit cycles near three figure-eight loops, and [1] obtain 13 small amplitude limit cycles.

(ii) The configuration of 15 limit cycles: [23] considered a cubic Hamiltonian system under quartic perturbations and obtained 4 different configurations of 15 limit cycles with a large limit cycle enclosing 14 limit cycles near two figure-eight loops of unperturbed system.

(iii) The configuration of 20 limit cycles and above: [12] considered a fourthorder near-Hamiltonian system and obtained three categories of configurations of limit cycles with 20, 21 and 23 limit cycles respectively, where 23 limit cycles has a distribution with 2 large limit cycles enclosing 7 medium limit cycles and 14 limit cycles obtained by homoclinic or heteroclinic bifurcation.

(iv) The configuration of 21 small amplitude limit cycles: [7] studied the local cyclicity of holomorphic quartic centers, and it is proved that 21 limit cycles of small amplitude bifurcate from a unique singular point.

Motivated by [21,22] and [23], in this paper, we study the number and distribution of limit cycles of a class of planar quartic system

$$\begin{cases} \dot{x} = H_y(x, y) + \varepsilon p(x, y), \\ \dot{y} = -H_x(x, y) + \varepsilon q(x, y), \end{cases}$$
(1.2)

where $0 < \varepsilon \ll 1$,

$$H(x,y) = \frac{x^2}{2} + \frac{y^2}{2} + \frac{\sqrt{ax^3}}{3} - \frac{a}{4}(x^4 + y^4), \quad a > 0,$$
(1.3)

p(x, y) and q(x, y) are polynomials of degree four. Let $x = \frac{1}{\sqrt{a}}x$, $y = \frac{1}{\sqrt{a}}y$, then system (1.2) can be transformed to

$$\begin{cases} \dot{x} = -y(y^2 - 1) + \varepsilon \sum_{i+j=0}^{4} a_{ij} x^i y^j \equiv f(x, y), \\ \dot{y} = x(x^2 - x - 1) + \varepsilon \sum_{i+j=0}^{4} b_{ij} x^i y^j \equiv g(x, y), \end{cases}$$
(1.4)

where a_{ij} , b_{ij} are arbitrary constants. The unperturbed system of (1.4) has a first integral

$$H(x,y) = \frac{x^2}{2} + \frac{y^2}{2} + \frac{x^3}{3} - \frac{x^4}{4} - \frac{y^4}{4}.$$
 (1.5)

Let

$$\begin{split} \delta_1 &= b_{01} + a_{10}, \ \delta_2 = b_{02} + \frac{1}{2}a_{11}, \ \delta_3 = b_{11} + 2a_{20}, \ \delta_4 = b_{03} + \frac{1}{3}a_{12}, \\ \delta_5 &= b_{12} + a_{21}, \ \delta_6 = b_{21} + 3a_{30}, \ \delta_7 = b_{04} + \frac{1}{4}a_{13}, \ \delta_8 = b_{13} + \frac{2}{3}a_{22}, \\ \delta_9 &= b_{22} + \frac{3}{2}a_{31}, \ \delta_{10} = b_{31} + 4a_{40}, \ \delta_{11} = 2a_{00} + \frac{2}{3}a_{02} + \frac{4}{5}a_{04}, \end{split}$$

$$\delta = (\delta_1, \delta_2, \dots, \delta_{10}, \delta_{11}), \quad \delta^* = (\delta_1^*, \delta_2^*, \dots, \delta_{10}^*, \delta_{11}^*)$$

Our main results are as follows.

Theorem 1.1. For $0 < \varepsilon \ll 1$,

- (i) there exist δ^* with $\delta_{10}^* < -3$ and $\delta_7^* = 5\delta_{11}^* = \delta_{10}^*$ such that (1.4) has at least 12 limit cycles, two of which are small amplitude limit cycles. The configuration of these limit cycles are shown in Figure 1(a).
- (ii) there exist δ^* with $\delta_8^* > 0$ and $\delta_7^* = \delta_8^*$ such that (1.4) has at least 12 or 11 limit cycles. The configuration of these limit cycles are shown in Figure 1(b) and Figure 1(d).
- (iii) there exist δ^* with $\delta_8^* > 0$ and $\delta_2^* = \frac{1}{2}\delta_{11}^* = \delta_8^*$ such that (1.4) has at least 9 limit cycles. The configuration of these limit cycles are shown in Figure 1(c).



Figure 1. (a), (b): The configuration of 12 limit cycles, (c) The configuration of 9 limit cycles, (d) The configuration of 11 limit cycles.

Remark 1.1. In Theorem 1.1, two new configurations of 12 limit cycles for the quartic system (1.4) are obtained in Figure 1(a) and Figure 1(b).

This paper is organized as follows. In Section 2, we will give the expression of displacement function for some orbits of system $(1.4)_{\varepsilon=0}$ and introduce some useful results in [9] and [10]. In Section 3, we will give the proof of Theorem

and

1.1. For simplicity, some results obtained by Maple and Mathematica are listed in Appendix.

2. Preliminaries

2.1. The expression of displacement function

When $\varepsilon = 0$, system (1.4) is reduced to

$$\dot{x} = -y(y^2 - 1), \quad \dot{y} = x(x^2 - x - 1).$$
 (2.1)

For i = 1, 2, system (2.1) has nine singular points:

• five elementary centers O(0,0), $C_i(\frac{1+\sqrt{5}}{2},(-1)^{i+1})$, $C_{i+2}(\frac{1-\sqrt{5}}{2},(-1)^{i+1})$,

• four hyperbolic saddles $S_i(\frac{1+(-1)^{i+1}\sqrt{5}}{2}, 0), S_{i+2}(0, (-1)^{i+1}).$ Let $h_1 = \frac{13+5\sqrt{5}}{24}, h_2 = \frac{13-5\sqrt{5}}{24}, h_3 = \frac{1}{4}, h_4 = \frac{19+5\sqrt{5}}{24}, h_5 = \frac{19-5\sqrt{5}}{24}$, then from (1.5) we have $H(O) = 0, \quad H(S_1) = h_1, \quad H(S_2) = h_2.$

$$H(O) = 0, \quad H(S_1) = h_1, \quad H(S_2) = h_2,$$

 $H(S_i) = h_3, \quad H(C_j) = h_4, \quad H(C_k) = h_5,$

where i, k = 3, 4 and j = 1, 2. The closed orbits of (2.1) can be described as follows (see Figure 2):



Figure 2. The phase portrait of system (2.1).

(1) Γ_h^1 , $h \in (-\infty, h_2)$: the family of closed orbits which surrounding all nine singular points.

- (2) $\Gamma_{h_2} = L_3 \cup L_4 \cup \{S_2\}$: the double-homoclinic loop connecting S_2 .
- (3) Γ_h^2 , $h \in (0, h_2)$: the family of closed orbits which surrounding O.

(4) Γ_h^3 , $h \in (h_2, h_3)$: the family of closed orbits which surrounding S_i , C_j , i = 1, 3, 4, j = 1, 2, 3, 4.

(5) $\Gamma_{h_3} = L_5 \cup L_6 \cup L_7 \cup L_8 \cup \{S_3, S_4\}$: two homoclinic loops L_5, L_6 and a heteroclinic loop $L_7 \cup L_8$, where L_{i+2} surrounding C_i , $i = 3, 4, L_7 \cup L_8$ surrounding C_1 C_2 and S_1 .

(6) $\Gamma_{\underline{h}}^4$, $h \in (h_3, h_5)$: the family of closed orbits which surrounding C_3 or C_4 .

(7) Γ_h^5 , $h \in (h_3, h_1)$: the family of closed orbits which surrounding C_1 , C_2 and S_1 . (8) $\Gamma_{h_1} = L_1 \cup L_2 \cup \{S_1\}$: the double-homoclinic loop connecting S_1 .

(9) Γ_h^6 , $h \in (h_1, h_4)$: the family of closed orbits which surrounding C_1 or C_2 .

Let $L_9 = \Gamma_{\frac{1}{2}}^5$ and $L_{10} = \Gamma_{-\frac{1}{4}}^1$. Recall that the displacement function of $L_i(i = 1...10)$ can be expressed as

$$d_i(\varepsilon, \delta) = \varepsilon N_i M_i(\delta) + O(\varepsilon^2),$$

where $N_i > 0$ and

$$M_{i}(\delta) = \int_{L_{i}} g_{0}(x, y) dx - f_{0}(x, y) dy$$

=
$$\sum_{j+k=0}^{4} a_{jk} A_{ijk} + b_{jk} B_{ijk}$$
 (2.2)

with $A_{ijk} = -\int_{L_i} x^j y^k dy$, $B_{ijk} = \int_{L_i} x^j y^k dx$. If $d_i = 0$, then there exist a homoclinic loop (resp. a heteroclinic orbit) L_i^* near L_i for $i = 1, \ldots, 6$ (resp. i = 7, 8). By (2.2) and the integration by parts, we have the following lemma.

Lemma 2.1. For L_i , i = 1, ..., 10, we have

$$M_{i}(\delta) = \begin{cases} \delta_{1}B_{i01} + \delta_{2}B_{i02} + \delta_{3}B_{i11} + \delta_{4}B_{i03} + \delta_{5}B_{i12} + \delta_{6}B_{i21} + \delta_{7}B_{i04} \\ + \delta_{8}B_{i13} + \delta_{9}B_{i22} + \delta_{10}B_{i31}, \quad i = 1, \dots, 6, 9, 10, \\ \delta_{1}B_{i01} + \delta_{2}B_{i02} + \delta_{3}B_{i11} + \delta_{4}B_{i03} + \delta_{5}B_{i12} + \delta_{6}B_{i21} + \delta_{7}B_{j04} \\ + \delta_{8}B_{i13} + \delta_{9}B_{i22} + \delta_{10}B_{i31} + (-1)^{i+1}\delta_{11}, \quad i = 7, 8. \end{cases}$$

By using Mathematica and Maple, the explicit expressions of the coefficients B_{ijk} are obtained, as given in Appendix. Then for $i = 1, 2, 5, 6, 1 \le j + k \le 4$, we have

$$B_{2jk} = (-1)^{k+1} B_{1jk}, \ B_{6jk} = (-1)^{k+1} B_{5jk}.$$

2.2. The approach to study homoclinic loop and double homoclinic loop

For the homoclinic loop L^* (resp. the double-homoclinic loop $L_a^* \cup L_b^*$) of system (1.4), let σ_0 is the divergence of saddle point,

$$\sigma_1 = \oint_L (f_{0x} + g_{0x}) dt \quad (\text{resp. } \sigma_{1i} = \oint_{L_i} (f_{0x} + g_{0x}) dt, \quad \text{for } i = a, b),$$

and R is the first saddle value at the saddle point. Then from [9] and [10], the stability of homoclinic loops and double-homoclinic loops can be discriminated by the following lemmas.

Lemma 2.2 ([9,10]). For $\varepsilon > 0$ small, the homoclinic loop L^* is inner stable (resp. unstable) if $\sigma_0 < 0$ (resp. > 0) or $\sigma_0 \equiv 0$, $\sigma_1 < 0$ (resp. > 0) or $\sigma_0 = \sigma_1 \equiv 0$, R < 0 (resp. > 0).

Lemma 2.3 ([9,10]). For $\varepsilon > 0$ small, the double-homoclinic loop $L_a^* \cup L_b^*$ is outer stable (resp. unstable) if $\sigma_0 < 0$ (resp. > 0) or $\sigma_0 \equiv 0, \sigma_{11} + \sigma_{12} < 0$ (resp. > 0) or $\sigma_0 = \sigma_{11} + \sigma_{12} \equiv 0, R > 0$ (resp. < 0).

3. Proof of the main results

Let $S_{j\varepsilon} = (x_j^s, y_j^s)$ is the saddle point of (1.4) near S_j , $C_{k\varepsilon} = (x_k^c, y_k^c)$ is the focus point of (1.4) near C_k , σ_{0j} is the divergence of $S_{j\varepsilon}$ and R_j is the first saddle value at $S_{j\varepsilon}$, where j, k = 1, 2, 3, 4. Now we consider the configuration of limit cycles in Figure 1(a) first, let $\delta_7 = \delta_{10}$ for (1.4), then by the implicit function theorem, we have the following lemma.

Lemma 3.1. For $\varepsilon > 0$ small, $d_1 = 0$ and $d_1 = d_2 = 0$ implies that there exist two functions

$$\begin{split} \phi_1 &= -\frac{1}{B_{101}} (B_{102}\delta_2 + B_{103}\delta_4 + B_{111}\delta_3 + B_{112}\delta_5 + B_{113}\delta_8 + B_{121}\delta_6 + B_{122} \ \delta_9 \\ &+ (B_{104} + B_{131}) \ \delta_{10}) + O(\varepsilon), \\ \phi_2 &= -\frac{B_{104}\delta_{10} + B_{112}\delta_5 + B_{122}\delta_9}{B_{102}} + O(\varepsilon) \end{split}$$

satisfying

• $d_1 \ge (<)0 \Leftrightarrow \delta_1 \le (>)\phi_1$,

•
$$d_1 = 0, \ d_2 \ge (<)0 \Leftrightarrow \delta_2 \ge (<)\phi_2$$

Thus, there exist a homoclinic loop L_1^* near L_1 as $d_1 = 0$ and a double-homoclinic loop $L_1^* \cup L_2^*$ near $L_1 \cup L_2$ as $d_1 = d_2 = 0$. Then we consider the stability of the double-homoclinic loop $L_1^* \cup L_2^*$. Using the implicit function theorem, we have

Lemma 3.2. If $\delta_i = \phi_i$ i = 1, 2, the divergence of $S_{1\varepsilon}$ can be expressed as

$$\sigma_{01} = \left(\left(\frac{1+\sqrt{5}}{2} - \frac{B_{111}}{B_{101}}\right)\delta_3 - \frac{B_{103}}{B_{101}}\delta_4 + \left(\left(\frac{1+\sqrt{5}}{2}\right)^2 - \frac{B_{121}}{B_{101}}\right)\delta_6 - \frac{B_{113}}{B_{101}}\delta_8 + \left(\left(\frac{1+\sqrt{5}}{2}\right)^3 - \frac{B_{131}}{B_{101}}\right)\delta_{10}\right)\varepsilon + O(\varepsilon^2).$$

If $\sigma_{01} = 0$, there exist a function

$$\phi_3 = -\frac{1}{(1+\sqrt{5})B_{101} - 2B_{111}} (2B_{103}\delta_4 + ((3+\sqrt{5})B_{101} - 2B_{121})\delta_6 + 2B_{113}\delta_8 + ((4-2\sqrt{5})B_{101} - 2B_{131})\delta_{10}) + O(\varepsilon)$$

satisfying $\sigma_{01} \ge (<)0 \Leftrightarrow \delta_3 \ge (<)\phi_3$ for $\varepsilon > 0$ small.

Under $\delta_i = \phi_i$, i = 1, 2, 3, then $\sigma_{01} = 0$, we consider the integral

$$\oint_{L_i^*} (f_{0x} + g_{0y}) dt \equiv \sigma_{1i}(\varepsilon, \delta), \quad i = 1, 2,$$

which converges finitely to $\sigma_{1i}(0,\delta) = \oint_{L_i} (f_{0x} + g_{0y}) dt$ from [8,9]. Then we have

Lemma 3.3. If $\delta_i = \phi_i$, i = 1, 2, 3, we obtain

 $\sigma_{11} = -\ 9.1505100310861\delta_4 - 0.20773663099846\delta_5 + 0.00333547529614\delta_6$

$$-14.7488764676445\delta_8 - 0.4636116024714\delta_9 + 0.00333542440294\delta_{10} + O(\varepsilon)$$

- $\sigma_{12} = -\ 9.1505100310861\delta_4 + 0.20773663099846\delta_5 + 0.00333547529614\delta_6$
 - $-14.7488764676445\delta_8 + 0.4636116024714\delta_9 + 0.00333542440294\delta_{10} + O(\varepsilon).$

For $\varepsilon > 0$ small, there exists two functions

$$\begin{split} \phi_4 &= 0.000364506933669710 \delta_{10} + 0.000364512500787624 \delta_6 - 0.0227021914946979 \delta_5 \\ &- 0.0506651105671599 \delta_9 - 1.61180922347967 \delta_8 + O(\varepsilon), \end{split}$$

 $\phi_5 = -2.23172774220490\delta_9 + O(\varepsilon)$

satisfying

• $\sigma_{11} \ge (<)0 \Leftrightarrow \delta_4 \le (>)\phi_4,$ • $\sigma_{11} = 0, \ \sigma_{12} \ge (<)0 \Leftrightarrow \delta_5 \ge (<)\phi_5.$

Proof. We know that

$$\sigma_{1i}(0,\delta) = \oint_{L_i} (f_{0x} + g_{0y})dt = \oint_{L_i} F(x,y)dt,$$

where

$$F(x,y) = \frac{1}{y - y^3} (\delta_1 + 2\delta_2 y + \delta_3 x + 3\delta_4 y^2 + 2\delta_5 xy + \delta_6 x^2 + 3\delta_8 xy^2 + 2\delta_9 x^2 y + (x^3 + 4y^3)\delta_{10}).$$

From Lemma 3.1 and 3.2, the results can be obtained by straightforward computation and the implicit function theorem. $\hfill \Box$

When $\sigma_{11} = \sigma_{12} = 0$, we consider the first saddle value of $S_{1\varepsilon}$. Hence we have the following lemma.

Lemma 3.4. For system (1.4), we obtain

(i) The first saddle value of $S_{1\varepsilon}$ denoted by R_1 as follows:

$$R_{1} = \frac{10^{6}}{\left(\sqrt{5}+3\right)^{2} \left(25+11\sqrt{5}\right)^{2} \left(5+\sqrt{5}\right)^{14}} \left(1024 \left(10336421\sqrt{5}\right)^{2} \left(25+11\sqrt{5}\right)^{2} \left(25+11\sqrt{5}\right)^{2} \left(5+\sqrt{5}\right)^{14} \delta_{4}\right)^{2} + 23112940 \delta_{1} + \frac{3}{2} \times 10^{-6} \left(\sqrt{5}+3\right)^{2} \left(25+11\sqrt{5}\right)^{2} \left(5+\sqrt{5}\right)^{14} \delta_{4} + 512 \left(38127971\sqrt{5}+85256735\right) \delta_{3} + 76800 \left(514229\sqrt{5}\right)^{2} + 1149851 \delta_{8} + 512 \left(64870463\sqrt{5}\right)^{2} + 145054765 \delta_{6} + 102400000 \left(51499217\sqrt{5}\right)^{2} + 115155750 \delta_{10} \varepsilon + O(\varepsilon^{2}).$$

(ii) For $\varepsilon > 0$ small, under $\delta_i = \phi_i$, $i = 1, \ldots, 5$, we have

$$R_1 = (-0.324163041746872\delta_8 + 0.00875795054111555\delta_6 + 0.00875788799581975\delta_{10})\varepsilon + O(\varepsilon^2)$$

and

~

$$\begin{split} \phi_6 &= -0.999992858742074\delta_{10} + 37.0135730147729\delta_8 + O(\varepsilon) \\ satisfying \ R_1 \geq (<)0 \Leftrightarrow \delta_6 \geq (<)\phi_6. \end{split}$$

The proof of Lemma 3.4 is given in Appendix. Let $I_{kpq} = \oint_{\Gamma_h^{6(k)}} x^p y^q dx$, and $\Gamma_h^{6(k)}$ is the family of closed orbits surrounding C_k for k = 1, 2. Now we consider the Hopf bifurcation of $C_{1\varepsilon}$ and $C_{2\varepsilon}$, which is equivalent to discussing the number of zeros of

$$M_k(h) = \delta_1 I_{k01} + \delta_2 I_{k02} + \delta_3 I_{k11} + \delta_4 I_{k03} + \delta_5 I_{k12} + \delta_6 I_{k21} + \delta_7 I_{k04} + \delta_8 I_{k13} + \delta_9 I_{k22} + \delta_{10} I_{k31}, \quad k = 1, 2$$
(3.1)

near $h = h_4$ for system (1.4). Then for $k = 1, 2, C_{k\varepsilon}$ has the same stability as the $\tilde{C}_{k\varepsilon}$ of system

$$\begin{aligned} \dot{x} &= -y(y^2 - 1) \\ &\equiv f_1(x, y), \\ \dot{y} &= x(x^2 - x - 1) + \varepsilon(\delta_1 y + \delta_2 y^2 + \delta_3 x y + \delta_4 y^3 + \delta_5 x y^2 + \delta_6 x^2 y \\ &\quad + \delta_7 y^4 + \delta_8 x y^3 + \delta_9 x^2 y^2 + \delta_{10} x^3 y) \\ &\equiv g_1(x, y). \end{aligned}$$
(3.2)

For $\tilde{C}_{1\varepsilon}$ and $\tilde{C}_{2\varepsilon}$ of system (3.2), we have the following lemma.

Lemma 3.5. If $\delta_i = \phi_i$, i = 1, ..., 6, the divergence of $\tilde{C}_{k\varepsilon}$, k = 1, 2 can be expressed as

$$div(\tilde{C}_{1\varepsilon}) = (0.00252924305133200\delta_9 - 0.000322469361956088\delta_{10} + 0.0591420565428257\delta_8)\varepsilon + O(\varepsilon^2),$$

$$div(\tilde{C}_{2\varepsilon}) = (0.0000147035582569595\delta_{10} - 0.00291043144903150\delta_9 + 0.0591613858149482\delta_8)\varepsilon + O(\varepsilon^2).$$

For $\varepsilon > 0$ small, there exists two functions

$$\begin{split} \phi_9 &= 0.127496391375381\delta_{10} - 23.3833029655569\delta_8 + O(\varepsilon), \\ \phi_8 &= 0.00280124722141849\delta_{10} + O(\varepsilon) \end{split}$$

satisfying

• div $(\tilde{C}_{1\varepsilon}) \ge (<)0 \Leftrightarrow \delta_9 \ge (<)\phi_9,$ • div $(\tilde{C}_{1\varepsilon}) = 0$, div $(\tilde{C}_{2\varepsilon}) \ge (<)0 \Leftrightarrow \delta_8 \ge (<)\phi_8.$

If $\operatorname{div}(\tilde{C}_{1\varepsilon}) = \operatorname{div}(\tilde{C}_{2\varepsilon}) = 0$, then we consider the first order focus value at $\tilde{C}_{k\varepsilon}$ for k = 1, 2. Hence, we have the following lemma.

Lemma 3.6. If $\delta_i = \phi_i$, i = 1, ..., 6, 8, 9, for k = 1, 2, the first order focus value F_k at $\tilde{C}_{k\varepsilon}$ can be expressed as

$$F_1 = -3.43865692299175(\delta_{10})^2 + 2.39687192915894\delta_{10},$$

$$F_2 = -0.823617871320597(\delta_{10})^2 - 1.66116528264483\delta_{10}.$$
(3.3)

For simplicity, the proof of Lemma 3.6 is given in Appendix. Next, we will give the proof of Theorem 1.1.

Proof of Theorem 1.1.

(i) The configuration of 12 limit cycles in Figure 1(a).

Under $\delta_7 = \delta_{10}$ and $\delta_i = \phi_i$, $i = 1, \dots, 6, 8, 9$, let $\delta_{10} < -3$, $\delta_{11} = \frac{1}{5}\delta_{10}$, then $F_k < 0$ in (3.3), and $C_{k\varepsilon}$ is a stable weak focus for k = 1, 2. Meanwhile, we obtain

$M_3 _{\varepsilon=0} = 0.0755839462405386\delta_{10},$	$M_4 _{\varepsilon=0} = -0.770557959539282\delta_{10},$
$M_5 _{\varepsilon=0} = -0.191195050488594\delta_{10},$	$M_6 _{\varepsilon=0} = 0.014677374520996\delta_{10},$
$M_7 _{\varepsilon=0} = -0.014188113535115\delta_{10},$	$M_8 _{\varepsilon=0} = -0.023440887449928\delta_{10},$
$M_9 _{\varepsilon=0} = -0.00568643542107560\delta_{10},$	$M_{10} _{\varepsilon=0} = -1.23218313343672\delta_{10},$
	(3.4)

and

$$div(C_{3\varepsilon}) = 0.625083775142779\delta_{10}\varepsilon + O(\varepsilon^{2}), div(O_{\varepsilon}) = 0.114878305554985\delta_{10}\varepsilon + O(\varepsilon^{2}), div(C_{4\varepsilon}) = -0.058223521406800\delta_{10}\varepsilon + O(\varepsilon^{2}).$$
(3.5)

By Lemma 3.5, keep δ_{10} fixed, let $0 < \delta_8 - \phi_8 \ll \varepsilon$ for $\varepsilon > 0$ small, $C_{2\varepsilon}$ change its stability from stable to unstable, then a stable limit cycle Γ_1 appears near $C_{2\varepsilon}$. Keeping δ_8 fixed and letting δ_9 satisfy $0 < \delta_9 - \phi_9 \ll \delta_8 - \phi_8 \ll \varepsilon$ change the stability of $C_{1\varepsilon}$ from stable to unstable, then there exist a stable limit cycle Γ_2 near $C_{1\varepsilon}$, see Figure 3(a).

Then we consider the homocilnic and double-homoclinic bifurcation of system (1.4) by Lemma 2.2 and 2.3. From (3.4), we know that $d_9 > 0$. Keep δ_9 fixed and let δ_6 satisfy $0 < \phi_6 - \delta_6 \ll \delta_9 - \phi_9 \ll \delta_8 - \phi_8 \ll \varepsilon$ such that the double-homoclinic loop $L_1^* \cup L_2^*$ is outer unstable and two homoclinic loop L_1^* and L_2^* are inner stable, which implies that there exist two unstable limit cycles Γ_3 and Γ_4 , and a stable limit cycle Γ_5 satisfying $\Gamma_2 \subset \Gamma_4 \subset L_1^*$, $\Gamma_1 \subset \Gamma_3 \subset L_2^*$ and $L_1^* \cup L_2^* \subset \Gamma_5$, see Figure 3(b). Keeping δ_6 fixed and letting δ_5 satisfy $0 < \delta_5 - \phi_5 \ll \phi_6 - \delta_6 \ll$ $\delta_9 - \phi_9 \ll \delta_8 - \phi_8 \ll \varepsilon$ make L_2^* to change its stability from stable to unstable and generate a stable limit cycle Γ_6 satisfying $\Gamma_3 \subset \Gamma_6 \subset L_2^*$. Keep δ_5 fixed, let δ_4 satisfy $0 < \phi_4 - \delta_4 \ll \delta_5 - \phi_5 \ll \phi_6 - \delta_6 \ll \delta_9 - \phi_9 \ll \delta_8 - \phi_8 \ll \varepsilon$, then the stability of L_1^* is inner unstable, which generates a stable limit cycle Γ_7 satisfying $\Gamma_4 \subset \Gamma_7 \subset L_1^*$, see Figure 3(c). Keeping δ_4 fixed and letting δ_3 satisfy $0 < \phi_3 - \delta_3 \ll \phi_4 - \delta_4 \ll \delta_5 - \phi_5 \ll \phi_6 - \delta_6 \ll \delta_9 - \phi_9 \ll \delta_8 - \phi_8 \ll \varepsilon \text{ force}$ the double-homoclinic loop $L_1^* \cup L_2^*$ to change its stability from outer unstable to outer stable, and two homoclinic loop L_1^* and L_2^* change their stability from inner unstable to inner stable. Then there exist three unstable limit cycles Γ_8 , Γ_9 and Γ_{10} satisfying $\Gamma_6 \subset \Gamma_8 \subset L_2^*$, $\Gamma_7 \subset \Gamma_9 \subset L_1^*$ and $L_1^* \cup L_2^* \subset \Gamma_{10} \subset \Gamma_5$, see Figure 3(d).

Keep δ_3 fixed, let δ_2 satisfy $0 < \delta_2 - \phi_2 \ll \phi_3 - \delta_3 \ll \phi_4 - \delta_4 \ll \delta_5 - \phi_5 \ll \phi_6 - \delta_6 \ll \delta_9 - \phi_9 \ll \delta_8 - \phi_8 \ll \varepsilon$ then L_2^* has broken. A stable limit cycle Γ_{11} is created with $\Gamma_8 \subset \Gamma_{11}$. Now keeping δ_2 fixed and letting δ_1 satisfy $0 < \phi_1 - \delta_1 \ll \delta_2 - \phi_2 \ll \phi_3 - \delta_3 \ll \phi_4 - \delta_4 \ll \delta_5 - \phi_5 \ll \phi_6 - \delta_6 \ll \delta_9 - \phi_9 \ll \delta_8 - \phi_8 \ll \varepsilon$, a stable limit cycle Γ_{12} is born out by breaking L_1^* , and $\Gamma_9 \subset \Gamma_{12}$.

By (3.4), (3.5) and Poincaré-Bendixson theorem, there are no more limit cycles can be found. Hence, system (1.4) can have a configuration of 12 limit cycles given



Figure 3. The configuration of 12 limit cycles in Theorem 1.1(i).

in Figure 3(e) or Figure 1(a).

(ii) The configuration of 12 or 11 limit cycles in Figure 1(b) and Figure 1(d).

Let $(c_1, c_2, c_3, c_4, c_5, c_6, c_7, c_8, c_9, c_{10}, c_{11}) = (\delta_1, \delta_2, \delta_3, \delta_5, \delta_4, \delta_9, \delta_6, \delta_{11}, \delta_{10}, \delta_8, \delta_7)$, then we consider $d_5 = 0$ and $d_5 = d_6 = 0$, which implies

$$\begin{split} \psi_1 &= 0.206731072574653c_9 - 1.94237745771629c_2 + 0.521848068796729c_3 \\ &- 2.88671366858555c_5 + 1.01099255361162c_4 - 0.315116996222073c_7 \\ &- 3.88475491543259c_{11} + 1.50128108534460c_{10} \\ &- 0.610308916135520c_6 + O(\varepsilon), \end{split}$$

 $\psi_2 = 0.520492322228799c_4 - 2c_{11} - 0.314207165919789c_6 + O(\varepsilon)$

satisfying

• $d_5 \ge (<)0 \Leftrightarrow c_1 \le (>)\psi_1,$ • $d_5 = 0, d_6 \ge (<)0 \Leftrightarrow c_2 \ge (<)\psi_2,$

for $\varepsilon > 0$ small. Thus, there exist a homoclinic loop L_5^* (resp. L_6^*) near L_5 (resp. L_6) as $d_5 = 0$ (resp. $d_5 = d_6 = 0$). Then we consider the stability of two homoclinic loops L_5^* and L_6^* . The divergence of $S_{3\varepsilon}$ and $S_{4\varepsilon}$ are expressed as follows:

$$\sigma_{03} = (4c_{11} + 3c_5 + 2c_2 + c_1)\varepsilon + O(\varepsilon^2),$$

$$\sigma_{04} = (-4c_{11} + 3c_5 - 2c_2 + c_1)\varepsilon + O(\varepsilon^2).$$

Under $c_i = \psi_i$, i = 1, 2, for $\varepsilon > 0$ small, we have

$$\begin{split} \psi_3 &= -\ 0.396151839847431 c_9 - 0.217086807805309 c_5 - 1.99480405639497 c_4 \\ &+ \ 0.603848160152564 c_7 - 2.87685473054684 c_{10} \\ &+ \ 1.20420936555072 c_6 + O(\varepsilon), \end{split}$$

 $\psi_4 = 0.603673008228676c_6 + O(\varepsilon)$

satisfying

- $\sigma_{03} \ge (<)0 \Leftrightarrow c_3 \ge (<)\psi_3$,
- $\sigma_{03} = 0, \ \sigma_{04} \ge (<)0 \Leftrightarrow c_4 \le (>)\psi_4.$

Under $c_i = \phi_i$, $i = 1, \ldots, 4$, we know that the integral

$$\oint_{L_i^*} (f_{0x} + g_{0y}) dt \equiv \sigma_{1i}(\varepsilon, \delta)$$

is converges finitely to $\sigma_{1i}(0,\delta) = \oint_{L_i} (f_{0x} + g_{0y}) dt$ for i = 5, 6. Then we have

$$\begin{split} \sigma_{15} &= -\ 0.648880199073c_5 - 0.04352635293150c_7 + 0.32126094789594c_{10} \\ &- 0.08653274020959c_6 - 0.043526234558150c_9 + O(\varepsilon), \\ \sigma_{16} &= -\ 0.648880199073c_5 - 0.04352635293150c_7 + 0.32126094789594c_{10} \\ &+ 0.08653274020959c_6 - 0.043526234558150c_9 + O(\varepsilon). \end{split}$$

For $\varepsilon > 0$ small, the implicit function theorem implies that

$$\begin{split} \psi_5 &= -\ 0.0670789995138281 c_9 - 0.0670791819409518 c_7 + 0.495100556859195 c_{10} \\ &- 0.133357036219043 c_6 + O(\varepsilon), \end{split}$$

$$\psi_6 = O(\varepsilon)$$

satisfying

•
$$\sigma_{15} \ge (<)0 \Leftrightarrow c_5 \le (>)\psi_5$$
,

•
$$\sigma_{15} = 0, \ \sigma_{16} \ge (<)0 \Leftrightarrow c_6 \ge (<)\psi_6.$$

If $\sigma_{15} = \sigma_{16} = 0$, then we consider the first saddle value R_j at $S_{j\varepsilon}$, j = 3, 4. Under $c_i = \psi_i$, i = 1, ..., 6, a straightforward computation shows that

$$R_{3} = (0.090176437715357c_{9} + 0.090176144273414c_{7} + 0.73481810027781c_{10})\varepsilon + O(\varepsilon^{2}),$$

$$R_{4} = (0.090176437715357c_{9} + 0.090176144273414c_{7} + 0.73481810027781c_{10})\varepsilon + O(\varepsilon^{2}).$$

Noting that the dominant part of R_3 and R_4 are identical, then for $\varepsilon > 0$ small, the implicit function theorem implies that there exist functions

$$\psi_7 = -1.00000325409725c_9 - 8.14869726576291c_{10} + O(\varepsilon)$$

$$\equiv \psi_7^* + O(\varepsilon)$$

and $\tilde{\psi}_7 = \psi_7^* + O(\varepsilon)$ satisfying $R_3 = 0 \Leftrightarrow c_7 = \psi_7$ and $R_4 = 0 \Leftrightarrow c_7 = \tilde{\psi}_7$. Then for R_3 and R_4 , we have the following two cases: (A) If $R_3 = R_4$, then $R_j \ge (<)0 \Leftrightarrow c_7 \ge (<)\psi_7$ for j = 3, 4.

(B) If $R_3 = 0$, then $R_4 > (<)0 \Leftrightarrow \psi_7 > (<)\tilde{\psi}_7$. For case (A), if $c_i = \psi_i$, i = 1, ..., 6 and $R_3 = R_4 = 0$, then there exist two functions

$$\psi_8 = 0.266668205545192c_9 + 4.47190585729362c_{10} + O(\varepsilon)$$

$$\psi_9 = -2529591.83803435c_{10} + O(\varepsilon)$$

such that

- $d_7 \ge (<)0 \Leftrightarrow c_8 \ge (<)\psi_8$,
- $d_7 = 0, d_8 \ge (<)0 \Leftrightarrow c_9 \ge (<)\psi_9,$

for $\varepsilon > 0$ small. Under $c_i = \psi_i$, $i = 1, \ldots, 9$, let $c_{11} = c_{10}$ and $c_{10} > 0$, then we have



Figure 4. The configuration of 12 or 11 limit cycles in Theorem 1.1(ii).

 $M_1|_{\varepsilon=0} = 0.352449325754430c_{10},$ $M_3|_{\varepsilon=0} = 0.06503013303720c_{10},$

 $M_2|_{\varepsilon=0} = 0.35244932575439c_{10},$ $M_4|_{\varepsilon=0} = -1.661787862219c_{10},$

$$M_9|_{\varepsilon=0} = 1.117130072425c_{10}, \qquad M_{10}|_{\varepsilon=0} = -4.687696341429c_{10}, \qquad (3.6)$$

and

$$div(C_{1\varepsilon}) = (0.487209419999990c_{10})\varepsilon + O(\varepsilon^2),$$

$$div(C_{2\varepsilon}) = (0.487209419999990c_{10})\varepsilon + O(\varepsilon^2),$$

$$div(C_{3\varepsilon}) = (-0.071614570000099c_{10})\varepsilon + O(\varepsilon^2),$$

$$div(C_{4\varepsilon}) = (-0.071614570000099c_{10})\varepsilon + O(\varepsilon^2),$$

$$div(O_{\varepsilon}) = (-0.08423254793588c_{10})\varepsilon + O(\varepsilon^2).$$

$$(3.7)$$

From (3.6) and (3.7), we know that $d_i > 0$ and $\operatorname{div}(C_{k\varepsilon}) > 0$ for i, k = 1, 2, then there exist two stable limit cycles $\tilde{\Gamma}_1$ and $\tilde{\Gamma}_2$ with $\tilde{\Gamma}_k$ surrounding $C_{k\varepsilon}$ for k = 1, 2. $d_3 > 0$ and $\operatorname{div}(O_{\varepsilon}) < 0$ create an unstable limit cycle $\tilde{\Gamma}_3$ surrounding O_{ε} , see Figure 4(a). Keeping c_{10} fixed and letting c_8 and c_9 satisfy $0 < \psi_8 - c_8 \ll \psi_9 - c_9 \ll \varepsilon$, then L_7^* and L_8^* are broken. From (3.6), $d_9 > 0$ implies that there exist an unstable limit cycle $\tilde{\Gamma}_4$ surrounding $\tilde{\Gamma}_1$ and $\tilde{\Gamma}_2$, see Figure 4(b).

Using the same arguments as the proof in (i), if $0 < \psi_1 - c_1 \ll c_2 - \psi_2 \ll \psi_3 - c_3 \ll c_4 - \psi_4 \ll \psi_5 - c_5 \ll c_6 - \psi_6 \ll \psi_7 - c_7 \ll \psi_8 - c_8 \ll \psi_9 - c_9 \ll \varepsilon$, eight limit cycles denoted by $\tilde{\Gamma}_k$, $k = 5, \ldots, 12$ are born out by homoclinic bifurcation, where $\tilde{\Gamma}_5 \subset \tilde{\Gamma}_7 \subset \tilde{\Gamma}_9 \subset \tilde{\Gamma}_{11}$ and $\tilde{\Gamma}_6 \subset \tilde{\Gamma}_8 \subset \tilde{\Gamma}_{10} \subset \tilde{\Gamma}_{12}$.

By (3.6), (3.7) and Poincaré-Bendixson theorem, there are no more limit cycles can be found. Thus, system (1.4) can have a configuration of 12 limit cycles given in Figure 4(c) or Figure 1(b).

For case (**B**), if $\psi_7 < \psi_7$, we have $R_4 < 0$, then there still have 12 limit cycles by the above discussion. If $\psi_7 > \tilde{\psi}_7$, then $R_4 > 0$. For $0 < \psi_1 - c_1 \ll \psi_2 - c_2 \ll$ $\psi_3 - c_3 \ll \psi_4 - c_4 \ll \psi_5 - c_5 \ll \psi_6 - c_6 \ll \psi_7 - c_7 \ll \psi_8 - c_8 \ll \psi_9 - c_9 \ll \varepsilon$, system (1.4) can have a configuration of 11 limit cycles given in Figure 4(d) or Figure 1(d).

(iii) The configuration of 9 limit cycles in Figure 1(c).

Let $(e_1, e_2, e_3, e_4, e_5, e_6, e_7, e_8, e_9, e_{10}, e_{11}) = (\delta_1, \delta_3, \delta_2, \delta_6, \delta_{10}, \delta_5, \delta_7, \delta_9, \delta_8, \delta_2, \delta_{11})$, then we consider $d_3 = 0$ and $d_3 = d_4 = 0$, which implies

$$\xi_1 = 0.0133493269231805e_5 + 0.0699742865225566e_2 - 0.115599134404137e_3$$

 $-0.0566249595993767e_4 + 0.00412670152129421e_9 + O(\varepsilon),$

 $\xi_2 = -2.65065878384217e_5 - 2.98859713453560e_3 - 1.65065878384213e_4 - 2.67707411234879e_9 + O(\varepsilon)$

satisfying

- $d_3 \ge (<)0 \Leftrightarrow e_1 \ge (<)\xi_1,$
- $d_3 = 0, d_4 \ge (<)0 \Leftrightarrow e_2 \le (>)\xi_2,$

for $\varepsilon > 0$ small. Thus, there exist a homoclinic loop L_3^* (resp. L_4^*) near L_3 (resp. L_4) as $d_3 = 0$ (resp. $d_4 = 0$). Then we consider the stability of two homoclinic loops L_3^* and L_4^* and the double-homoclinic loop $L_3^* \cup L_4^*$. The divergence of $S_{2\varepsilon}$ can be expressed as

$$\sigma_{02} = \left(e_1 + \left(\frac{1-\sqrt{5}}{2}\right)e_2 + \left(\frac{1-\sqrt{5}}{2}\right)^2 e_4 + \left(\frac{1-\sqrt{5}}{2}\right)^3 e_5\right)\varepsilon + O(\varepsilon^2).$$

Under $e_i = \xi_i$, i = 1, 2, we have

$$\xi_3 = -0.807972116469746e_5 - 0.807972116469731e_4 - 0.966493886747985e_9 + O(\varepsilon)$$

satisfying $\sigma_{02} \ge (<)0 \Leftrightarrow e_3 \ge (<)\xi_3$ for $\varepsilon > 0$ small.

Under $e_i = \xi_i$, i = 1, 2, 3, we know that the integral

$$\oint_{L_i^*} (f_{0x} + g_{0y}) dt \equiv \sigma_{1i}(\varepsilon, \delta)$$

is converges finitely to $\sigma_{1i}(0,\delta) = \oint_{L_i} (f_{0x} + g_{0y}) dt$ for i = 3, 4. Then we have

$$\begin{split} \sigma_{13} &= -8.28835327774732e_4 - 11.9902272944765e_9 + 11.4671586220581e_5 + O(\varepsilon) \\ \sigma_{14} &= 13.9333141863372e_4 - 17.9535962832705e_9 - 13.3690282910674e_5 + O(\varepsilon). \end{split}$$

For $\varepsilon > 0$ small, the implicit function theorem implies

$$\begin{aligned} \xi_4 &= -1.44663564554711e_9 + 1.38352676795827e_5 + O(\varepsilon), \\ \xi_5 &= 6.45048712058850e_9 + O(\varepsilon) \end{aligned}$$

satisfying

- $\sigma_{13} \ge (<)0 \Leftrightarrow e_4 \le (>)\xi_4,$
- $\sigma_{13} = 0, \ \sigma_{14} \ge (<)0 \Leftrightarrow e_5 \ge (<)\xi_5.$

If $d_5 = 0$, then there exist a homoclinic loop L_5^* near L_5 . Under $e_i = \xi_i$, $i = 1, \ldots, 5$, the function $d_5 = 0$ implies

$$\begin{split} \xi_6 &= -\ 36.2658668936540 e_9 + 1.92125792695251 e_{10} + 3.84251585390503 e_7 \\ &+ \ 0.603673008228676 e_8 + O(\varepsilon) \end{split}$$

satisfying $d_5 \ge (<)0 \Leftrightarrow e_6 \ge (<)\xi_6$ for $\varepsilon > 0$ small. Under $e_6 = \xi_6$, we consider the stability of L_5^* . The divergence of $S_{3\varepsilon}$ can be expressed as

$$\sigma_{03} = (4e_7 + 3e_3 + 2e_{10} + e_1)\varepsilon + O(\varepsilon^2).$$

Under $e_i = \xi_i$, i = 1, ..., 6, the implicit function theorem implies

$$\xi_7 = 8.81823189932078e_9 - 0.5e_{10} + O(\varepsilon)$$

with $\sigma_{03} \ge (<)0 \Leftrightarrow e_7 \ge (<)\xi_7$ for $\varepsilon > 0$ small.

If $d_1 = 0$, then there exist a homoclinic loop L_1^* near L_1 . Under $e_i = \xi_i$, $i = 1, \ldots, 7$, the function $d_1 = 0$ implies

$$\xi_8 = -8.84756452029888e_9 + O(\varepsilon)$$

with $d_1 \ge (<)0 \Leftrightarrow e_8 \le (>)\xi_8$ for $\varepsilon > 0$ small. Under $e_i = \xi_i$, i = 1, ..., 8, let $e_9 > 0$ and $e_{10} = \frac{1}{2}e_{11} = e_9$, then we have

$$\begin{split} M_2|_{\varepsilon=0} &= -40.6328910646561e_9, & M_6|_{\varepsilon=0} &= 25.3774299377964e_9, \\ M_7|_{\varepsilon=0} &= -1.31633151616553e_9, & M_8|_{\varepsilon=0} &= -80.0131548318940e_9, \\ M_9|_{\varepsilon=0} &= -90.3252285537313e_9, & M_{10}|_{\varepsilon=0} &= 56.6383768634101e_9, \end{split}$$

and

$$div(O_{\varepsilon}) = (1.38752288302499e_9)\varepsilon + O(\varepsilon^2), div(C_{1\varepsilon}) = (-12.4380043676037e_9)\varepsilon + O(\varepsilon^2),$$

$$div(C_{2\varepsilon}) = (59.6514480049598e_9)\varepsilon + O(\varepsilon^2), div(C_{3\varepsilon}) = (-0.45477200064143e_9)\varepsilon + O(\varepsilon^2), div(S_{1\varepsilon}) = (55.4130703327366e_9)\varepsilon + O(\varepsilon^2), div(C_{4\varepsilon}) = (-76.5743328924742e_9)\varepsilon + O(\varepsilon^2).$$
(3.9)

Keep e_9 fixed, let $0 < e_8 - \xi_8 \ll \varepsilon$, then L_1^* is broken, and an unstable limit cycle $\hat{\Gamma}_1$ is born out with div $(C_{1\varepsilon}) < 0$ from (3.9). Then we consider the stability of L_5^* . Keeping e_8 fixed and letting e_7 satisfy $0 < \xi_7 - e_7 \ll e_8 - \xi_8 \ll \varepsilon$ make L_5^* inner stable, which creates an unstable limit cycle $\hat{\Gamma}_2$ by div $(C_{3\varepsilon}) < 0$ from (3.9). Keep e_7 fixed, let e_6 satisfy $0 < e_6 - \xi_6 \ll \xi_7 - e_7 \ll e_8 - \xi_8 \ll \varepsilon$, then L_5^* is broken such that a stable limit cycle $\hat{\Gamma}_3$ is born out with $\hat{\Gamma}_2 \subset \hat{\Gamma}_3$, see Figure 5(a).



Figure 5. The configuration of 9 limit cycles in Theorem 1.1(iii).

From (3.8) and (3.9), we know that $d_{10} > 0$ and $\operatorname{div}(O_{\varepsilon}) > 0$. Using the same arguments as the proof in (i), if $0 < \xi_1 - e_1 \ll e_2 - \xi_2 \ll \xi_3 - e_3 \ll \xi_4 - e_4 \ll e_5 - \xi_5 \ll e_6 - \xi_6 \ll \xi_7 - e_7 \ll e_8 - \xi_8 \ll \varepsilon$, six limit cycles denoted by $\hat{\Gamma}_i, i = 4, \ldots, 9$ are born out by homoclinic and double-homoclinic bifurcation, where $\hat{\Gamma}_4$, $\hat{\Gamma}_6$, $\hat{\Gamma}_8$ are three large limit cycle satisfying $\hat{\Gamma}_8 \subset \hat{\Gamma}_6 \subset \hat{\Gamma}_4$, and $\hat{\Gamma}_5$, $\hat{\Gamma}_7$, $\hat{\Gamma}_9$ are three limit cycle satisfying $\hat{\Gamma}_5 \subset \hat{\Gamma}_7 \subset \hat{\Gamma}_9$.

By (3.8), (3.9) and Poincaré-Bendixson theorem, there are no more limit cycles can be found. Thus, system (1.4) can have a configuration of 9 limit cycles given in Figure 5(b) or Figure 1(c). \Box

Appendix

(i) For i = 1,

The coefficients of B_{ijk} .

$B_{101} = -0.71904466554725,$	$B_{102} = -1.20091460208964,$
$B_{111} = -1.13729463105521,$	$B_{103} = -1.73071612964796,$
$B_{112} = -1.89532210354832,$	$B_{121} = -1.82324946574506,$
$B_{104} = -2.40182920417926,$	$B_{113} = -2.73035698260822,$
$B_{122} = -3.03578632023233,$	$B_{131} = -2.96054409680028.$

(ii) For $i = 3$, $B_{302} = B_{312} = B_{304} = B_{322} = 0$ and		
$\begin{split} B_{301} &= 0.551994572607374, \\ B_{313} &= -0.00227791684252500, \\ B_{303} &= 0.0638100947891942, \end{split}$	$\begin{split} B_{311} &= -0.0386254263825246, \\ B_{321} &= 0.0312566703729678, \\ B_{331} &= -0.00736875600955716. \end{split}$	
(iii) For $i = 4$, $B_{402} = B_{412} = B_{404} = B_{422} = 0$ and		
$B_{401} = -9.81473228309552,$ $B_{403} = -24.4489399024426,$ $B_{413} = -20.8436385173344,$	$\begin{split} B_{411} &= -7.11432788344956, \\ B_{421} &= -13.4327242365103, \\ B_{431} &= -20.5470521199600. \end{split}$	
(iv) For $i = 5$,		
$\begin{split} B_{501} &= -0.346076110948626, \\ B_{511} &= 0.180599150155223, \\ B_{521} &= -0.109054464546348, \\ B_{504} &= -1.34442087312147, \\ B_{513} &= 0.519557519456792, \end{split}$	$\begin{split} B_{502} &= -0.672210436560734, \\ B_{503} &= -0.999022639846328, \\ B_{512} &= 0.349880371151931, \\ B_{522} &= -0.211213336173452, \\ B_{531} &= 0.071544685608874. \end{split}$	
(v) For $i = 7$, $B_{702} = B_{712} = B_{704} = B_{722} = 0$ and		
$B_{701} = 0.860340022251728,$ $B_{703} = 0.523816320319384,$ $B_{711} = 0.220440474306400,$	$B_{713} = 0.0964066908842402,$ $B_{721} = 0.0805755727012404,$ $B_{731} = 0.0343493803409738.$	
(vi) For $i = 8$, $B_{802} = B_{812} = B_{804} = B_{822} = 0$ and		
$B_{801} = -6.62001300968878,$ $B_{803} = -15.9462292017875,$ $B_{813} = -20.5549267098162,$	$B_{811} = -7.75865784091594,$ $B_{821} = -11.5342330095348,$ $B_{831} = -19.0262241837840.$	
(vii) For $i = 9$, $B_{902} = B_{912} = B_{904} = B_{922} = 0$ and		
$B_{901} = -4.19725312629488,$ $B_{903} = -10.5746251421787,$ $B_{913} = -15.4276285714229,$	$B_{911} = -6.09755329224136,$ $B_{921} = -9.48569314042712,$ $B_{931} = -15.5832464326685.$	
(viii) For $i = 10$, $B_{10,0,2} = B_{10,1,2} = B_{10,0,4} = B_{10,2,2} = 0$ and		
$B_{10,0,1} = -11.6691715767516,$ $B_{10,0,3} = -32.7737807547298,$ $B_{10,1,3} = -23.7135257919004,$	$B_{10,1,1} = -7.16862898645332,$ $B_{10,2,1} = -16.2371008419409,$ $B_{10,3,1} = -23.4057298283944.$	

The proof of Lemma 3.4 (i). Let

$$T = \begin{pmatrix} 1 & b \\ c & bc+1 \end{pmatrix}, \quad J = \frac{\partial(f,g)}{\partial(x,y)}(S_{1\varepsilon})$$

such that $TJT^{-1} = \text{diag}(\lambda_{11}, \lambda_{12})$, where $S_{1\varepsilon} = (x_1^s, y_1^s)$, λ_{11} and λ_{12} are the eigenvalues of J. A straightforward computation shows that

$$\begin{aligned} x_1^s &= \frac{1+\sqrt{5}}{2} - \frac{1}{5+\sqrt{5}} \left((7+3\sqrt{5})b_{40} + (4+2\sqrt{5})b_{30} + (3+\sqrt{5})b_{20} + (1+\sqrt{5})b_{10} \right. \\ &+ 2b_{00})\,\varepsilon + O(\varepsilon^2), \\ y_1^s &= \frac{1}{2} \left(-\left(1+\sqrt{5}\right)a_{10} - \left(3-\sqrt{5}\right)a_{20} - 2\left(2+\sqrt{5}\right)a_{30} - \left(7+3\sqrt{5}\right)a_{40} \right. \\ &- 2a_{00})\,\varepsilon + O(\varepsilon^2). \end{aligned}$$

Then we have

$$\begin{split} f_x(S_{1\varepsilon}) &= \left(4 \, a_{40} \left(\frac{1+\sqrt{5}}{2} \right)^3 + 3 \, a_{30} \left(\frac{1+\sqrt{5}}{2} \right)^2 + 2 \, a_{20} \left(\frac{1+\sqrt{5}}{2} \right) + a_{10} \right) \varepsilon \\ &+ O(\varepsilon^2), \\ f_y(S_{1\varepsilon}) &= 1 + \left(a_{31} \left(\frac{1+\sqrt{5}}{2} \right)^3 + a_{21} \left(\frac{1+\sqrt{5}}{2} \right)^2 + a_{11} \left(\frac{1+\sqrt{5}}{2} \right) + a_{01} \right) \varepsilon \\ &+ O(\varepsilon^2), \\ g_x(S_{1\varepsilon}) &= \frac{5+\sqrt{5}}{2} - \frac{1}{5+\sqrt{5}} (6 \sqrt{5} b_{00} + 3 \, b_{10} \sqrt{5} + 4 \, b_{20} \sqrt{5} + 2 \, b_{30} \sqrt{5} - 4 \, b_{40} \sqrt{5} \\ &+ 2 \, b_{00} + 11 \, b_{10} + 8 \, b_{20} + 4 \, b_{30} - 8 \, b_{40}) \varepsilon + O(\varepsilon^2), \\ g_y(S_{1\varepsilon}) &= \left(b_{31} \left(\frac{1+\sqrt{5}}{2} \right)^3 + b_{21} \left(\frac{1+\sqrt{5}}{2} \right)^2 + b_{11} \left(\frac{1+\sqrt{5}}{2} \right) + b_{01} \right) \varepsilon \\ &+ O(\varepsilon^2), \end{split}$$

 $\quad \text{and} \quad$

$$\begin{split} \lambda_{11} &= \frac{\sqrt{10+2\sqrt{5}}}{2} + \frac{1}{20\sqrt{10+2\sqrt{5}}} \left(40\sqrt{5}a_{21} + 70\sqrt{5}a_{31} + 10\,a_{10}\sqrt{10+2\sqrt{5}} \right. \\ &+ 10\,a_{20}\sqrt{10+2\sqrt{5}} + 45\,a_{30}\sqrt{10+2\sqrt{5}} + 80\,a_{40}\sqrt{10+2\sqrt{5}} \\ &+ 5\,b_{11}\sqrt{10+2\sqrt{5}} + 15\,b_{21}\sqrt{10+2\sqrt{5}} + 20\,b_{31}\sqrt{10+2\sqrt{5}} \\ &+ 10\,a_{20}\sqrt{5}\sqrt{10+2\sqrt{5}} + 15\,a_{30}\sqrt{5}\sqrt{10+2\sqrt{5}} + 10\,b_{31}\sqrt{5}\sqrt{10+2\sqrt{5}} \\ &+ 5\,b_{11}\sqrt{5}\sqrt{10+2\sqrt{5}} + 40\,a_{40}\sqrt{5}\sqrt{10+2\sqrt{5}} + 10\,\sqrt{5}a_{01} + 30\sqrt{5}a_{11} \\ &- 10\,b_{30} - 20\,b_{20} - 40\,b_{10} + 20\,b_{00} - 28\sqrt{5}b_{00} - 4\,b_{10}\sqrt{5} - 12\,b_{20}\sqrt{5} \\ &+ 12\,b_{40}\sqrt{5} + 50\,a_{01} + 50\,a_{11} + 100\,a_{21} + 150\,a_{31} + 5\,b_{21}\sqrt{5}\sqrt{10+2\sqrt{5}} \\ &+ 10\,b_{01}\sqrt{10+2\sqrt{5}} - 6\,b_{30}\sqrt{5} + 20\,b_{40} \right)\varepsilon + O(\varepsilon^2), \\ \lambda_{12} &= -\frac{\sqrt{10+2\sqrt{5}}}{2} + \frac{1}{20\sqrt{10+2\sqrt{5}}}\left(5\,b_{11}\sqrt{5}\sqrt{10+2\sqrt{5}} + 5\,b_{21}\sqrt{5}\right) \end{split}$$

$$\begin{array}{l} \times \sqrt{10+2\sqrt{5}}+15\,a_{30}\,\sqrt{5}\sqrt{10+2\sqrt{5}}+40\,a_{40}\,\sqrt{5}\sqrt{10+2\sqrt{5}}\\ +\,10\,a_{20}\,\sqrt{5}\sqrt{10+2\,\sqrt{5}}+20\,b_{31}\,\sqrt{10+2\,\sqrt{5}}-10\,\sqrt{5}a_{01}-30\,\sqrt{5}a_{11}\\ +\,10\,a_{10}\,\sqrt{10+2\,\sqrt{5}}+10\,a_{20}\,\sqrt{10+2\,\sqrt{5}}+45\,a_{30}\,\sqrt{10+2\,\sqrt{5}}\\ +\,80\,a_{40}\,\sqrt{10+2\,\sqrt{5}}+10\,b_{01}\,\sqrt{10+2\,\sqrt{5}}+5\,b_{11}\,\sqrt{10+2\,\sqrt{5}}\\ +\,15\,b_{21}\,\sqrt{10+2\,\sqrt{5}}+28\,\sqrt{5}b_{00}+4\,b_{10}\,\sqrt{5}+12\,b_{20}\,\sqrt{5}+6\,b_{30}\,\sqrt{5}\\ -\,12\,b_{40}\,\sqrt{5}-20\,b_{40}+40\,b_{10}-20\,b_{00}-50\,a_{01}-50\,a_{11}-100\,a_{21}\\ -\,150\,a_{31}+10\,b_{31}\,\sqrt{5}\sqrt{10+2\,\sqrt{5}}-40\,\sqrt{5}a_{21}-70\,\sqrt{5}a_{31}\\ +10\,b_{30}+20\,b_{20})\,\varepsilon+O(\varepsilon^2). \end{array}$$

Thus, for matrix T, we have

$$\begin{split} b &= \sqrt{\frac{2}{5+\sqrt{5}}} + \frac{\sqrt{2}}{\left(5+\sqrt{5}\right)^3} \left(6\sqrt{5+\sqrt{5}}\sqrt{5}b_{00} + 3\sqrt{5+\sqrt{5}}\sqrt{5}b_{10} + 4\sqrt{5+\sqrt{5}} \right. \\ &\times \sqrt{5}b_{20} + 2\sqrt{5+\sqrt{5}}\sqrt{5}b_{30} - 4\sqrt{5+\sqrt{5}}\sqrt{5}b_{40} - 20\sqrt{5}\sqrt{2}a_{20} - 45\sqrt{5}\sqrt{2}a_{30} \\ &+ 10\sqrt{5+\sqrt{5}}\sqrt{5}a_{11} + 15\sqrt{5+\sqrt{5}}\sqrt{5}a_{21} + 25\sqrt{5+\sqrt{5}}\sqrt{5}a_{31} + 2\sqrt{5+\sqrt{5}}b_{00} \\ &+ 11\sqrt{5+\sqrt{5}}b_{10} + 8\sqrt{5+\sqrt{5}}b_{20} + 4\sqrt{5+\sqrt{5}}b_{30} - 8\sqrt{5+\sqrt{5}}b_{40} - 15\sqrt{2}a_{10} \\ &+ 10\sqrt{5}\sqrt{2}b_{11} + 15\sqrt{5}\sqrt{2}b_{21} + 25\sqrt{5}\sqrt{2}b_{31} - 5\sqrt{5}\sqrt{2}a_{10} + 5\sqrt{5}\sqrt{2}b_{01} \\ &- 105\sqrt{2}a_{30} - 220\sqrt{2}a_{40} + 15\sqrt{2}b_{01} + 20\sqrt{2}b_{11} + 35\sqrt{2}b_{21} + 55\sqrt{2}b_{31} \\ &+ 20\sqrt{5+\sqrt{5}}a_{11} + 35\sqrt{5+\sqrt{5}}a_{21} + 55\sqrt{5}+\sqrt{5}a_{31} - 40\sqrt{2}a_{20} \\ &+ 5\sqrt{5+\sqrt{5}}\sqrt{5}a_{01} + 15\sqrt{5+\sqrt{5}}a_{01} - 100\sqrt{5}\sqrt{2}a_{40} \right)\varepsilon + O(\varepsilon^2), \\ c &= -\frac{\sqrt{10+2\sqrt{5}}}{4} + \left(\frac{\sqrt{2}\sqrt{5}+\sqrt{5}}{40(25+11\sqrt{5})}\left(55\sqrt{5}a_{01} + 90\sqrt{5}a_{11} + 145\sqrt{5}a_{21} \\ &+ 34\sqrt{5}b_{00} + 37b_{10}\sqrt{5} + 36b_{20}\sqrt{5} + 18b_{30}\sqrt{5} - 36b_{40}\sqrt{5} + 125a_{01} \\ &+ 525a_{31} + 70b_{00} + 85b_{10} + 80b_{20} + 40b_{30} - 80b_{40} + 325a_{21} \\ &+ 235\sqrt{5}a_{31} + 200a_{11}\right)\varepsilon + O(\varepsilon^2). \end{split}$$

Then system (1.4) can be reduced to

$$\begin{cases} \dot{u} = \lambda_{11} \left(u + \sum_{k=2}^{3} \sum_{j+l=k} m_{jl} u^{j} v^{l} + O(|(u,v)|^{4}) \right), \\ \dot{v} = -\lambda_{12} \left(-v + \sum_{k=2}^{3} \sum_{j+l=k} n_{jl} u^{j} v^{l} + O(|(u,v)|^{4}) \right), \end{cases}$$

via the linear transformation

$$\begin{pmatrix} u \\ v \end{pmatrix} = T \begin{pmatrix} x - x_1^s \\ y - y_1^s \end{pmatrix}.$$

By the formula in [10], the first saddle value at $S_{1\varepsilon}$ is

$$R_1 = m_{21} + n_{12} - m_{20}m_{11} + n_{02}n_{11},$$

where $m_{11}, m_{20}, m_{21}, n_{11}, n_{02}$ and n_{12} are given as follows:

$$\begin{split} m_{21} &= -\frac{3\left(5\sqrt{5}+17\right)\sqrt{2}}{4\left(5+\sqrt{5}\right)^{3/2}} + \frac{4000}{\left(5+\sqrt{5}\right)^{17/2}\left(25+11\sqrt{5}\right)} \left(16860\sqrt{5}+\sqrt{5}\sqrt{5}a_{22}\right) \\ &+ 20583\sqrt{5}+\sqrt{5}\sqrt{5}b_{21}+28644\sqrt{5}+\sqrt{5}\sqrt{5}b_{31}+5210\sqrt{2}\sqrt{5}b_{12} \\ &+ 12834\sqrt{2}\sqrt{5}b_{00}+211210\sqrt{2}\sqrt{5}a_{21}+8412\sqrt{5}+\sqrt{5}\sqrt{5}b_{01} \\ &+ 14742\sqrt{2}\sqrt{5}b_{10}+25290\sqrt{5}+\sqrt{5}\sqrt{5}b_{13}+27222\sqrt{5}+\sqrt{5}\sqrt{5}a_{20} \\ &+ 56550\sqrt{2}\sqrt{5}a_{03}+5210\sqrt{5}+\sqrt{5}\sqrt{5}a_{12}+14106\sqrt{2}\sqrt{5}b_{20} \\ &+ 82665\sqrt{2}a_{01}\sqrt{5}+133755\sqrt{2}a_{11}\sqrt{5}+13611\sqrt{5}+\sqrt{5}\sqrt{5}b_{11} \\ &+ 15630\sqrt{5}+\sqrt{5}\sqrt{5}b_{03}+91500a_{13}\sqrt{2}\sqrt{5}+8412\sqrt{5}+\sqrt{5}\sqrt{5}a_{10} \\ &+ 30435\sqrt{5}+\sqrt{5}\sqrt{5}b_{03}+91500a_{13}\sqrt{2}\sqrt{5}+8412\sqrt{5}+\sqrt{5}\sqrt{5}a_{10} \\ &+ 30435\sqrt{5}+\sqrt{5}b_{11}+138075\sqrt{5}+\sqrt{5}a_{30}+64050\sqrt{5}+\sqrt{5}b_{31} \\ &+ 46025\sqrt{5}+\sqrt{5}b_{21}+11650\sqrt{2}b_{12}+472280\sqrt{2}a_{21}+184845\sqrt{2}a_{01} \\ &+ 726465\sqrt{2}a_{31}+28698\sqrt{2}b_{00}+32964\sqrt{2}b_{10}+31542\sqrt{2}b_{20}+6111\sqrt{2}b_{30} \\ &+ 37700\sqrt{2}b_{22}+204600a_{13}\sqrt{2}+126450\sqrt{2}a_{03}+56550\sqrt{5}+\sqrt{5}b_{13} \\ &+ 37700\sqrt{5}+\sqrt{5}a_{22}+11650\sqrt{5}+\sqrt{5}a_{40}+34950\sqrt{5}+\sqrt{5}b_{03} \\ &- 94062\sqrt{2}b_{40}+60870\sqrt{5}+\sqrt{5}a_{20}+299085\sqrt{2}a_{11}+256200\sqrt{5}+\sqrt{5}a_{40} \\ &+ 324885\sqrt{2}a_{31}\sqrt{5}-42066\sqrt{2}\sqrt{5}b_{40}+61749\sqrt{5}+\sqrt{5}\sqrt{5}a_{30} \\ &+ 16860\sqrt{2}b_{22}\sqrt{5}+2733\sqrt{2}\sqrt{5}b_{30}\right)\varepsilon + O(\varepsilon^2), \\ n_{11} &= -\frac{3\sqrt{5}+1}{2(5+\sqrt{5})} - \frac{3200}{(5+\sqrt{5})^{15/2}(25+11\sqrt{5})(\sqrt{5}+3)} \left(7370\sqrt{5+\sqrt{5}}b_{10} \\ &+ 89275\sqrt{2}\sqrt{5}a_{22}+290890\sqrt{2}\sqrt{5}a_{20}+3296\sqrt{5}+\sqrt{5}\sqrt{5}b_{10} \\ &+ 89275\sqrt{2}\sqrt{5}a_{22}+290890\sqrt{2}\sqrt{5}a_{20}+34100\sqrt{2}\sqrt{5}a_{02} \\ &- 24675\sqrt{5}+\sqrt{5}\sqrt{5}b_{22}+55175\sqrt{2}\sqrt{5}a_{12}+466455\sqrt{2}\sqrt{5}b_{30} \\ \end{array}$$

$$\begin{split} &+175565 \sqrt{2}\sqrt{5}a_{10} + 19617 \sqrt{5} + \sqrt{5}\sqrt{5}b_{40} - 21075 \sqrt{5} + \sqrt{5}b_{02} \\ &+ 7710 \sqrt{5} + \sqrt{5}b_{20} + 15080 \sqrt{5} + \sqrt{5}b_{30} + 43865 \sqrt{5} + \sqrt{5}b_{40} \\ &+ 1043025 \sqrt{2}a_{30} + 1617225 \sqrt{2}a_{40} - 55175 \sqrt{5} + \sqrt{5}b_{22} - 34100 \sqrt{5} + \sqrt{5}b_{12} \\ &+ 76250 \sqrt{2}a_{02} + 123375 \sqrt{2}a_{12} + 199625 \sqrt{2}a_{22} - 15250 \sqrt{5} + \sqrt{5}\sqrt{5}b_{12} \\ &+ 650450 \sqrt{2}a_{20} + 228750 \sqrt{2}a_{00} + 3752 \sqrt{5} + \sqrt{5}\sqrt{5}b_{00} + 392575 \sqrt{2}a_{10} \\ &+ 8390 \sqrt{5} + \sqrt{5}b_{00} - 9425 \sqrt{5} + \sqrt{5}b_{02} \sqrt{5} + 3448 \sqrt{5} + \sqrt{5}\sqrt{5}b_{20} \\ &+ 6744 \sqrt{5} + \sqrt{5}\sqrt{5}b_{30} \right) \varepsilon + O(\varepsilon^2), \\ n_{12} &= \frac{3 (5 \sqrt{5} + 17) \sqrt{2}}{4 (5 + \sqrt{5})^{3/2}} + \frac{4000}{(5 + \sqrt{5})^{17/2} (25 + 11 \sqrt{5})} \left(16860 \sqrt{5} + \sqrt{5}\sqrt{5}b_{32} \right) \\ &+ 20583 \sqrt{5} + \sqrt{5}\sqrt{5}b_{21} + 28644 \sqrt{5} + \sqrt{5}\sqrt{5}b_{31} - 5210 \sqrt{2}\sqrt{5}b_{12} \\ &- 12834 \sqrt{2}\sqrt{5}b_{00} - 211210 \sqrt{2}\sqrt{5}a_{21} + 8412 \sqrt{5} + \sqrt{5}\sqrt{5}b_{13} \\ &- 14742 \sqrt{2}\sqrt{5}b_{10} - 2733 \sqrt{2}\sqrt{5}b_{30} + 25290 \sqrt{5} + \sqrt{5}\sqrt{5}b_{13} \\ &- 56550 \sqrt{2}\sqrt{5}a_{03} + 91500 a_{13} \sqrt{2}\sqrt{5} + 8412 \sqrt{5} + \sqrt{5}\sqrt{5}a_{10} \\ &+ 46025 \sqrt{5} + \sqrt{5}b_{21} - 11650 \sqrt{2}b_{12} - 472280 \sqrt{2}a_{21} - 184845 \sqrt{2}a_{01} \\ &+ 37700 \sqrt{5} + \sqrt{5}a_{22} + 11650 \sqrt{5} + \sqrt{5}a_{12} + 18810 \sqrt{5} + \sqrt{5}a_{10} \\ &- 726465 \sqrt{2}a_{31} - 28698 \sqrt{2}b_{00} - 32964 \sqrt{2}b_{10} - 31542 \sqrt{2}b_{20} - 6111 \sqrt{2}b_{30} \\ &- 37700 \sqrt{2}b_{22} - 204000 a_{13} \sqrt{2} - 126450 \sqrt{2}a_{33} + 56550 \sqrt{5} + \sqrt{5}b_{13} \\ &+ 18810 \sqrt{5} + \sqrt{5}b_{11} + 138075 \sqrt{5} + \sqrt{5}a_{30} + 64050 \sqrt{5} + \sqrt{5}b_{31} \\ &+ 18810 \sqrt{5} + \sqrt{5}b_{11} + 138075 \sqrt{5} + \sqrt{5}a_{40} + 60870 \sqrt{5} + \sqrt{5}a_{20} \\ &- 299085 \sqrt{2}a_{11} + 94062 \sqrt{2}b_{40} - 324885 \sqrt{2}a_{31} \sqrt{5} + 27222 \sqrt{5} + \sqrt{5}\sqrt{5}a_{20} \\ &+ 42066 \sqrt{2}\sqrt{5}b_{40} - 16860 \sqrt{2}b_{22} \sqrt{5} \right) \varepsilon + O(\varepsilon^2), \\ m_{20} &= \frac{3\sqrt{5} + 1}{4(5 + \sqrt{5})} + \frac{1600}{(5 + \sqrt{5})^{15/2} (25 + 11\sqrt{5}) (\sqrt{5} + 3)} \left(7370 \sqrt{5} + \sqrt{5}b_{10} \\ &+ 15080 \sqrt{5} + \sqrt{5}b_{30} + 43865 \sqrt{5} + \sqrt{5}b_{40} + 370125 \sqrt{2}a_{10} + 619950 \sqrt{2}a_{20} \right)$$

$$\begin{split} &+ 1898925 \sqrt{2}a_{40} + 55175 \sqrt{5} + \sqrt{5}b_{22} + 34100 \sqrt{5} + \sqrt{5}b_{12} + 228750 \sqrt{2}a_{00} \\ &+ 123375 \sqrt{2}a_{12} + 199625 \sqrt{2}a_{22} + 68200 \sqrt{5} + \sqrt{5}a_{21} + 165525 \sqrt{5} + \sqrt{5}a_{31} \\ &+ 15250 \sqrt{5} + \sqrt{5}\sqrt{5}b_{12} + 3752 \sqrt{5} + \sqrt{5}\sqrt{5}b_{00} + 30500 \sqrt{5} + \sqrt{5}a_{21} \sqrt{5} \\ &+ 89275 \sqrt{2}\sqrt{5}a_{22} + 277250 \sqrt{2}\sqrt{5}a_{20} + 34100 \sqrt{2}\sqrt{5}a_{02} \\ &+ 3448 \sqrt{5} + \sqrt{5}\sqrt{5}b_{20} - 6820 \sqrt{2}\sqrt{5}b_{11} + 24675 \sqrt{5} + \sqrt{5}\sqrt{5}b_{22} \\ &+ 55175 \sqrt{2}\sqrt{5}a_{12} - 10040 \sqrt{2}\sqrt{5}b_{01} + 19617 \sqrt{5} + \sqrt{5}\sqrt{5}b_{40} \\ &+ 21075 \sqrt{5} + \sqrt{5}b_{02} - 22450 \sqrt{2}b_{01} - 15250 \sqrt{2}b_{11} \\ &+ 70425 \sqrt{2}b_{31} + 8390 \sqrt{5} + \sqrt{5}b_{00} + 21075 \sqrt{5} + \sqrt{5}a_{11} + 4215 \sqrt{2}\sqrt{5}b_{21} \\ &+ 479100 \sqrt{2}\sqrt{5}a_{30} + 849225 \sqrt{2}\sqrt{5}a_{40} + 1071300 \sqrt{2}a_{30} + 76250 \sqrt{2}a_{02} \\ &+ 9425 \sqrt{5} + \sqrt{5}a_{11} \sqrt{5} + 102300 \sqrt{2}\sqrt{5}a_{00} + 3296 \sqrt{5} + \sqrt{5}\sqrt{5}b_{10} \\ &+ 31495 \sqrt{2}\sqrt{5}b_{31} + 9425 \sqrt{5} + \sqrt{5}b_{50} \sqrt{5} + 74025 \sqrt{5} + \sqrt{5}a_{31} \sqrt{5} \\ &+ 7710 \sqrt{5} + \sqrt{5}b_{20} + 6744 \sqrt{5} + \sqrt{5}\sqrt{5}b_{30} + 9425 \sqrt{2}b_{21} \\ &+ 165525 \sqrt{2}\sqrt{5}a_{10} \right) \varepsilon + O(\varepsilon^2), \\ m_{11} = \frac{(-3\sqrt{5} - 1) \sqrt{2}}{(5 + \sqrt{5})^{3/2}} + \frac{1600}{(\sqrt{5} + 3) (25 + 111 \sqrt{5}) (5 + \sqrt{5})^{13/2}} \left(13640 \sqrt{5} + \sqrt{5} \\ &\times \sqrt{5}a_{22} - 3699 \sqrt{2}\sqrt{5}b_{10} - 4626 \sqrt{2}\sqrt{5}b_{30} + 44444 \sqrt{5} + \sqrt{5}\sqrt{5}a_{20} \\ &+ 11650 \sqrt{5} + \sqrt{5}a_{12} + 59980 \sqrt{5} + \sqrt{5}a_{10} + 99380 \sqrt{5} + \sqrt{5}a_{20} \\ &+ 18850 \sqrt{5} + \sqrt{5}a_{12} + 59980 \sqrt{5} + \sqrt{5}a_{30} + 159360 \sqrt{5} + \sqrt{5}a_{30} \\ &+ 18850 \sqrt{2}b_{12} - 16245 \sqrt{2}a_{21} - 6205 \sqrt{2}a_{01} - 10040 \sqrt{2}a_{11} - 26285 \sqrt{2}a_{31} \\ &- 8271 \sqrt{2}b_{10} - 8278 \sqrt{2}b_{20} - 10344 \sqrt{2}b_{30} - 20232 \sqrt{2}b_{40} + 30500 \sqrt{2}b_{22} \\ &+ 8430 \sqrt{2}\sqrt{5}b_{12} + 13640 \sqrt{2}b_{22} \sqrt{5} - 3708 \sqrt{2}5b_{00} - 7265 \sqrt{2}\sqrt{3}a_{41} \\ &+ 30500 \sqrt{5} + \sqrt{5}a_{12} - 8292 \sqrt{2}b_{00} + 5210 \sqrt{5}\sqrt{2}b_{40} \\ &+ 8430 \sqrt{5} + \sqrt{5}a_{12} - 8292 \sqrt{2}b_{00} + 5210 \sqrt{5}\sqrt{2}b_{40} \\ &+ 8430 \sqrt{5} + \sqrt{5}a_{12} - 8292 \sqrt{2}b_{00} + 5210 \sqrt{5}\sqrt{2}b_{20} \right) \varepsilon + O(\varepsilon^2), \\ n_{02} = \frac{(3\sqrt{5} + 1)\sqrt{2}}{2(5 + \sqrt{5})^{3/2}} + \frac{800}{(\sqrt{5} + 3) (25 + 11\sqrt{5}) (5 + \sqrt{5})^{13$$

$$\begin{split} &\times \sqrt{5}a_{22} - 644 \sqrt{5} + \sqrt{5} \sqrt{5}b_{21} - 4812 \sqrt{5} + \sqrt{5} \sqrt{5}b_{31} + 8430 \sqrt{2} \sqrt{5}b_{12} \\ &+ 24125 \sqrt{2} \sqrt{5}a_{21} + 1534 \sqrt{5} + \sqrt{5} \sqrt{5}b_{01} + 9048 \sqrt{2} \sqrt{5}b_{40} + 8271 \sqrt{2}b_{10} \\ &- 42360 \sqrt{5} + \sqrt{5} \sqrt{5}a_{20} - 8430 \sqrt{5} + \sqrt{5} \sqrt{5}a_{12} + 3702 \sqrt{2} \sqrt{5}b_{20} \\ &+ 2775 \sqrt{2}a_{01} \sqrt{5} + 9700 \sqrt{2}a_{11} \sqrt{5} + 1042 \sqrt{5} + \sqrt{5} \sqrt{5}b_{11} + 8278 \sqrt{2}b_{20} \\ &- 10760 \sqrt{5} + \sqrt{5}b_{31} - 290130 \sqrt{5} + \sqrt{5}a_{40} - 1440 \sqrt{5} + \sqrt{5}b_{21} \\ &+ 6205 \sqrt{2}a_{01} + 21690 \sqrt{2}a_{11} + 117785 \sqrt{2}a_{31} + 8292 \sqrt{2}b_{00} \\ &- 56550 \sqrt{5} + \sqrt{5}a_{10} - 94720 \sqrt{5} + \sqrt{5}a_{20} + 3430 \sqrt{5} + \sqrt{5}b_{01} \\ &- 11650 \sqrt{5} + \sqrt{5}a_{02} + 11650 \sqrt{2}b_{02} + 5210 \sqrt{5} \sqrt{2}b_{02} - 5210 \sqrt{5}a_{02} \sqrt{5} + \sqrt{5} \\ &- 15630 \sqrt{5}a_{00} \sqrt{5} + \sqrt{5} + 3708 \sqrt{2} \sqrt{5}b_{00} + 53945 \sqrt{2}a_{21} + 4626 \sqrt{2} \sqrt{5}b_{30} \\ &- 25290 \sqrt{5} + \sqrt{5} \sqrt{5}a_{10} - 73200 \sqrt{5} + \sqrt{5} \sqrt{5}a_{30} + 2330 \sqrt{5} + \sqrt{5}b_{11} \\ &+ 10344 \sqrt{2}b_{30} + 20232 \sqrt{2}b_{40} + 30500 \sqrt{2}b_{22} - 30500 \sqrt{5} + \sqrt{5}a_{22} \\ &- 163680 \sqrt{5} + \sqrt{5}a_{30} - 18850 \sqrt{5} + \sqrt{5}a_{12} + 13640 \sqrt{2}b_{22} \sqrt{5} \\ &+ 3699 \sqrt{2} \sqrt{5}b_{10} + 52675 \sqrt{2}a_{31} \sqrt{5} - 129750 \sqrt{5} + \sqrt{5} \sqrt{5}a_{40} \\ &+ 18850 \sqrt{2}b_{12} - 34950 \sqrt{5} + \sqrt{5}a_{00} \right) \varepsilon + O(\varepsilon^2). \end{split}$$

(ii) By Lemmas 3.1-3.3, the result can be obtained by the implicit function theorem. $\hfill\square$

The proof of Lemma 3.6. Let $b_k = b_0^k + b_1^k \varepsilon + O(\varepsilon^2)$, $c_k = c_0^k + c_1^k \varepsilon + O(\varepsilon^2)$ and

$$T_k = \begin{pmatrix} 1 & b_k \\ c_k & b_k c_k + 1 \end{pmatrix}, \quad k = 1, 2,$$

where

$$\begin{split} b_0^1 &= b_0^2 = -\frac{\sqrt{-4+2\sqrt{\sqrt{5}+5}}}{\sqrt{\sqrt{5}+5}}, \quad c_0^1 = c_0^2 = \frac{\sqrt{-4+2\sqrt{\sqrt{5}+5}}}{2}, \\ b_1^1 &= -\frac{\sqrt{5}-5}{10((\sqrt{5}-5)\beta_0c_0^1+10b_0^1)} \left(\left(\left(14\sqrt{5}-10\right)\delta_1 + \left(3\sqrt{5}+5\right)\delta_{10} + \left(2\sqrt{5}+20\right)\delta_3 + \left(14\sqrt{5}-10\right)\delta_4 + \left(2\sqrt{5}+20\right)\delta_5 + \left(6\sqrt{5}+10\right)\delta_6 + \left(14\sqrt{5}-10\right)\delta_7 + \left(2\sqrt{5}+20\right)\delta_8 + \left(6\sqrt{5}+10\right)\delta_9 + \left(14\sqrt{5}-10\right)\delta_2\right) (b_0^1)^2 + (10\delta_1+20\delta_2+30\delta_4 + \left(10\sqrt{5}+20\right)\delta_{10} + \left(5\sqrt{5}+5\right)\delta_3 + \left(10\sqrt{5}+10\right)\delta_5 \end{split}$$

$$\begin{split} &+ \left(5\sqrt{5}+15\right)\delta_{6}+40\,\delta_{7}+\left(15\sqrt{5}+15\right)\delta_{8}+\left(10\sqrt{5}+30\right)\delta_{9}\right)b_{0}^{1}+10\beta_{1}^{1}\right),\\ c_{1}^{1}=-\frac{1}{2\left(\sqrt{5}+5\right)^{3/2}\sqrt{-4+2\sqrt{\sqrt{5}+5}}}\left(\left(6\sqrt{5}+2\right)\delta_{1}+\left(2\sqrt{5}+4\right)\delta_{10}\right.\\ &+ \left(3\sqrt{5}+11\right)\delta_{3}+\left(6\sqrt{5}+2\right)\delta_{4}+\left(3\sqrt{5}+11\right)\delta_{5}+\left(4\sqrt{5}+8\right)\delta_{6}\right.\\ &+ \left(3\sqrt{5}+11\right)\delta_{8}+\left(4\sqrt{5}+8\right)\delta_{9}+\left(6\sqrt{5}+2\right)\delta_{2}+\left(6\sqrt{5}+2\right)\delta_{7}\right),\\ b_{1}^{2}=\frac{\sqrt{5}-5}{10\left(\left(\sqrt{5}-5\right)\beta_{0}c_{0}^{2}+10b_{0}^{2}\right)}\left(\left(\left(14\sqrt{5}-10\right)\delta_{1}+\left(3\sqrt{5}+5\right)\delta_{10}\right.\\ &+ \left(2\sqrt{5}+20\right)\delta_{3}+\left(14\sqrt{5}-10\right)\delta_{4}-\left(2\sqrt{5}+20\right)\delta_{5}+\left(6\sqrt{5}+10\right)\delta_{6}\right.\\ &- \left(14\sqrt{5}-10\right)\delta_{7}+\left(2\sqrt{5}+20\right)\delta_{8}-\left(6\sqrt{5}+10\right)\delta_{9}\right.\\ &- \left(14\sqrt{5}-10\right)\delta_{2}\right)\left(b_{0}^{2}\right)^{2}-\left(10\delta_{1}-20\delta_{2}+30\delta_{4}+\left(10\sqrt{5}+20\right)\delta_{10}\right.\\ &- \left(10\sqrt{5}+10\right)\delta_{5}+\left(5\sqrt{5}+15\right)\delta_{6}-40\delta_{7}+\left(15\sqrt{5}+15\right)\delta_{8}\right.\\ &- \left(10\sqrt{5}+30\right)\delta_{9}+\left(5\sqrt{5}+5\right)\delta_{3}\right)b_{0}^{2}-10\beta_{1}^{2}\right),\\ c_{1}^{2}=\frac{1}{2\left(\sqrt{5}+5\right)^{3/2}}\sqrt{-4+2\sqrt{\sqrt{5}+5}}\left(\left(6\sqrt{5}+2\right)\delta_{1}+\left(2\sqrt{5}+4\right)\delta_{10}\right.\\ &+ \left(3\sqrt{5}+11\right)\delta_{3}+\left(6\sqrt{5}+2\right)\delta_{4}-\left(3\sqrt{5}+11\right)\delta_{5}+\left(4\sqrt{5}+8\right)\delta_{6}\right.\\ &+ \left(3\sqrt{5}+11\right)\delta_{8}-\left(4\sqrt{5}+8\right)\delta_{9}-\left(6\sqrt{5}+2\right)\delta_{2}-\left(6\sqrt{5}+2\right)\delta_{7}\right). \end{split}$$

with $\beta_0 = \sqrt{5 + \sqrt{5}}$ and

$$\beta_{1}^{1} = \frac{-1}{10\sqrt{5+\sqrt{5}}} \left(\left(14\sqrt{5}-10 \right) \delta_{1} + \left(3\sqrt{5}+5 \right) \delta_{10} + \left(14\sqrt{5}-10 \right) \delta_{2} \right. \\ \left. + \left(14\sqrt{5}-10 \right) \delta_{4} + \left(2\sqrt{5}+20 \right) \delta_{5} + \left(6\sqrt{5}+10 \right) \delta_{6} + \left(14\sqrt{5}-10 \right) \delta_{7} \right. \\ \left. + \left(2\sqrt{5}+20 \right) \delta_{8} + \left(6\sqrt{5}+10 \right) \delta_{9} + \left(2\sqrt{5}+20 \right) \delta_{3} \right), \\ \beta_{1}^{2} = \frac{1}{10\sqrt{5+\sqrt{5}}} \left(\left(14\sqrt{5}-10 \right) \delta_{1} + \left(3\sqrt{5}+5 \right) \delta_{10} - \left(14\sqrt{5}-10 \right) \delta_{2} \right. \\ \left. + \left(14\sqrt{5}-10 \right) \delta_{4} - \left(2\sqrt{5}+20 \right) \delta_{5} + \left(6\sqrt{5}+10 \right) \delta_{6} - \left(14\sqrt{5}-10 \right) \delta_{7} \right. \\ \left. + \left(2\sqrt{5}+20 \right) \delta_{8} - \left(6\sqrt{5}+10 \right) \delta_{9} + \left(2\sqrt{5}+20 \right) \delta_{3} \right).$$

By the linear transformation

$$\begin{pmatrix} u_k \\ v_k \end{pmatrix} = T_k \begin{pmatrix} x - x_k^c \\ y - y_k^c \end{pmatrix}, \quad k = 1, 2,$$

system (3.2) can be reduced to

$$\begin{pmatrix} \dot{u}_k \\ \dot{v}_k \end{pmatrix} = \begin{pmatrix} \alpha_1^k \varepsilon + O(\varepsilon^2) & -(\beta_0 + \beta_1^k \varepsilon + O(\varepsilon^2)) \\ (\beta_0 + \beta_1^k \varepsilon + O(\varepsilon^2)) & \alpha_1^k \varepsilon + O(\varepsilon^2) \end{pmatrix} \begin{pmatrix} u_k \\ v_k \end{pmatrix} + \begin{pmatrix} f_k(u_k, v_k, \varepsilon) \\ g_k(u_k, v_k, \varepsilon) \end{pmatrix}$$

for k = 1, 2, where

$$\begin{aligned} \alpha_1^1 &= \frac{1}{2}\delta_1 + (\frac{\sqrt{5}}{2} + 1)\delta_{10} + \delta_2 + (\frac{\sqrt{5} + 1}{4})\delta_3 + \frac{3}{2}\delta_4 + (\frac{\sqrt{5} + 1}{2})\delta_5 + (\frac{3 + \sqrt{5}}{4})\delta_6 \\ &+ 2\,\delta_7 + (\frac{3\sqrt{5} + 3}{4})\delta_8 + (\frac{3 + \sqrt{5}}{2})\delta_9, \\ \alpha_1^2 &= \frac{1}{2}\delta_1 + (\frac{\sqrt{5}}{2} + 1)\delta_{10} - \delta_2 + (\frac{\sqrt{5} + 1}{4})\delta_3 + \frac{3}{2}\delta_4 - (\frac{\sqrt{5} + 1}{2})\delta_5 + (\frac{3 + \sqrt{5}}{4})\delta_6 \\ &- 2\,\delta_7 + (\frac{3\sqrt{5} + 3}{4})\delta_8 - (\frac{3 + \sqrt{5}}{2})\delta_9. \end{aligned}$$

From [2], the first order focus value F_k at $\tilde{C}_{k\varepsilon}$ can be expressed as

$$\begin{split} F_k = &\frac{1}{16} \{ (f_k)_{u_k u_k u_k} + (f_k)_{u_k v_k v_k} + (g_k)_{u_k u_k v_k} + (g_k)_{v_k v_k v_k} + \frac{1}{\beta_0} [(f_k)_{u_k v_k} ((f_k)_{u_k u_k} + (f_k)_{v_k v_k}) - (g_k)_{u_k v_k} ((g_k)_{u_k u_k} + (g_k)_{v_k v_k}) - (f_k)_{u_k u_k} (g_k)_{u_k u_k} \\ &+ (f_k)_{v_k v_k} (g_k)_{v_k v_k}] \} |_{u_k = v_k = \varepsilon = 0} \end{split}$$

for k = 1, 2. Thus, we can get (3.3) by the straightforward computation.

Declarations

Conflict of interest. The authors declare that they have no known competing financial interests or personal relationships that could have appeared to influence the work reported in this paper.

Data availability statement. Data sharing not applicable to this article as no datasets were generated or analysed during the current study.

References

- Y. R. Carvalho, L. P. C. Da Cruz and L. F. S. Gouveia, New lower bound for the Hilbert number in low degree Kolmogorov systems, Chaos Solitons Fractals, 2023, 175, 113937.
- [2] S. N. Chow, C. Li and D. Wang, Normal Forms and Bifurcation of Planar Vector Fields, Cambridge University Press, New York, 1994.
- [3] C. J. Christopher and N. G. Lloyd, Polynomial systems: A lower bound for the Hilbert numbers, Proc. Roy. Soc. London Ser. A, 1995, 450(1938), 219–224.
- [4] R. D. Euzébio, J. Llibre and D. J. Tonon, Lower bounds for the number of limit cycles in a generalised Rayleigh-Liénard oscillator, Nonlinearity, 2022, 35(8), 3883–3906.
- [5] L. Gavrilov and I. D. Iliev, The limit cycles in a generalized Rayleigh-Liénard oscillator, Discrete Contin. Dyn. Syst., 2023, 43(6), 2381–2400.

- [6] W. Geng, M. Han, Y. Tian and A Ke, Heteroclinic bifurcation of limit cycles in perturbed cubic Hamiltonian systems by higher-order analysis, J. Diff. Eqs., 2023, 357, 412–435.
- [7] J. Giné, L. F. S. Gouveia and J. Torregrosa, Lower bounds for the local cyclicity for families of centers, J. Diff. Eqs., 2021, 275, 309–331.
- [8] M. Han, Cyclicity of planar homoclinic loops and quadratic integrable systems, Sci. China Ser. A, 1997, 40(12), 1247–1258.
- M. Han and J. Chen, On the number of limit cycles in double homoclinic bifurcations, Sci. China Ser. A, 2000, 43(9), 914–928.
- [10] M. Han, S. Hu and X. Liu, On the stability of double homoclinic and heteroclinic cycles, Nonlinear Anal., 2003, 53(5), 701–713.
- M. Han and J. Li, Lower bounds for the Hilbert number of polynomial systems, J. Diff. Eqs., 2012, 252, 3278–3304.
- [12] M. Han, D. Shang, W. Zheng and P. Yu, Bifurcation of limit cycles in a fourthorder near-Hamiltonian system, Internat. J. Bifur. Chaos, 2007, 17(11), 4117– 4144.
- [13] M. Han, J. Yang and J. Li, General study on limit cycle bifurcation near a double homoclinic loop, J. Diff. Eqs., 2023, 347, 1–23.
- [14] M. Han and Z. Zhang, Cyclicity 1 and 2 conditions for a 2-polycycle of integrable systems on the plane, J. Diff. Eqs., 1999, 155, 245–261.
- [15] C. Li, C. Liu and J. Yang, A cubic system with thirteen limit cycles, J. Diff. Eqs., 2009, 246, 3609–3619.
- [16] J. Li, Hilbert's 16th problem and bifurcations of planar polynomial vector fields, Internat. J. Bifur. Chaos, 2003, 13(1), 47–106.
- [17] J. Li and Y. Liu, New results on the study of Z_q -equivariant planar polynomial vector fields, Qual. Theory Dyn. Syst., 2010, 9(1–2), 167–219.
- [18] S. Liu and M. Han, Bifurcation theory of limit cycles by higher order Melnikov functions and applications, J. Diff. Eqs., 2024, 403, 29–66.
- [19] J. Llibre, Some families of quadratic systems with at most one limit cycle, Bul. Acad. Ştiinţe Repub. Mold. Mat., 2023, 101(1), 8–15.
- [20] P. Yu and M. Han, Small limit cycles bifurcating from fine focus points in cubic order Z₂-equivariant vector fields, Chaos Solitons Fractals, 2005, 24(1), 329–348.
- [21] H. Zang, W. Chen and T. Zhang, Perturbation from a cubic Hamiltonian with three figure eight-loops, Chaos Solitons Fractals, 2004, 22(1), 61–74.
- [22] H. Zang, T. Zhang and M. Han, The number and distributions of limit cycles for a class of cubic near-Hamiltonian systems, J. Math. Anal. Appl., 2006, 316(2), 679–696.
- [23] T. Zhang, M. Han, H. Zang and X. Meng, Bifurcations of limit cycles for a cubic Hamiltonian system under quartic perturbations, Chaos Solitons Fractals, 2004, 22(5), 1127–1138.