

MAPPING PROPERTY FOR BILINEAR Θ -TYPE GENERALIZED FRACTIONAL INTEGRAL OPERATOR ON GRAND VARIABLE EXPONENTS HERZ-MORREY SPACES*

Jinqi Wang¹ and Xijuan Chen^{1,†}

Abstract The aim of this paper is to establish the boundedness of the bilinear θ -type generalized fractional integral operators $B\tilde{T}_{\beta,\theta}$ and their commutators $B\tilde{T}_{\beta,\theta,b_1,b_2}$, generated by $b_1, b_2 \in \text{BMO}(\mathbb{R}^n)$ and $B\tilde{T}_{\beta,\theta}$, on Lebesgue spaces with variable exponent $L^{q(\cdot)}(\mathbb{R}^n)$. Under the assumption that variable exponents $\alpha(\cdot)$ and $q_i(\cdot)$ for $i = 1, 2$ satisfy the log decay at both infinity and the origin, the authors prove that the $B\tilde{T}_{\beta,\theta}$ and $B\tilde{T}_{\beta,\theta,b_1,b_2}$ are bounded on the grand variable exponents Herz spaces $\dot{K}_{q(\cdot)}^{\alpha(\cdot),p,\theta}(\mathbb{R}^n)$ (or $K_{q(\cdot)}^{\alpha(\cdot),p,\theta}(\mathbb{R}^n)$) and grand variable exponents Herz-Morrey spaces $M\dot{K}_{p,\theta,q(\cdot)}^{\alpha(\cdot),\lambda}(\mathbb{R}^n)$ (or $MK_{p,\theta,q(\cdot)}^{\alpha(\cdot),\lambda}(\mathbb{R}^n)$), respectively.

Keywords Bilinear θ -type generalized fractional integral, commutator, space $\text{BMO}(\mathbb{R}^n)$, grand variable exponents Herz-Morrey space, grand variable exponents Herz space.

MSC(2010) 42B20, 42B35, 47A07, 47B47.

1. Introduction

In 2022, Lu and Tao [13] introduced a bilinear θ -type generalized fractional integral operator $B\tilde{T}_{\beta,\theta}$, which is defined on the non-homogeneous metric measure spaces. And also they showed that the $B\tilde{T}_{\beta,\theta}$ is bounded from the product of spaces $L^{p_1}(\mu) \times L^{p_2}(\mu)$ into spaces $L^q(\mu)$, and it is also bounded from the product of Morrey spaces $M_{q_1}^{p_1}(\mu) \times M_{q_2}^{p_2}(\mu)$ into spaces $M_q(\mu)$. The following year, Lu et al. [19] prove that the $B\tilde{T}_{\beta,\theta}$ is bounded from the product of the generalized Morrey spaces $M_{q_1}^{\mu_1}(\mu) \times M_{q_2}^{\mu_2}(\mu)$ into spaces $M_q^\mu(\mu)$ for $\mu_1\mu_2 = \mu$. More researches on the bilinear θ -type generalized fractional and related operators on various functions spaces can be seen in [12, 14–16, 18].

As we also know, the theory of function spaces has many applications in harmonic analysis, including partial differential equations and wavelet analysis [1, 9, 14].

[†]The corresponding author.

¹College of Mathematics and Statistics, Northwest Normal University, Lanzhou 730070, China

*The authors were supported by the Master Foundation of Northwest Normal University (Grant No. KYZZS2025115).

Email: wjqxjy1920@126.com(J. Wang), chenxijuan2023@126.com(X. Chen)

In 1992, T. Iwaniec and C. Sbordone [8] first introduced the grand Lebesgue spaces $L^p(\mathbb{R}^n)$. Since then, many papers have focused on the bounded properties of integral operators on spaces L^p , and we can find in [2, 26]. On these bases, further advances have been made in the study of the grand spaces with variable exponents. For example, in 2020, H. Nafis et al. [20] introduced the grand Herz spaces with variable exponents $\dot{K}_{q(\cdot)}^{\alpha(\cdot), p, \theta}(\mathbb{R}^n)$, and they proved that sublinear operators T is bounded on spaces $\dot{K}_{q(\cdot)}^{\alpha(\cdot), p, \theta}(\mathbb{R}^n)$. In 2022, H. Nafis et al. [21] proved the boundedness of multilinear Calderón-Zygmund operators $\mathcal{T}$ on spaces $\dot{K}_{q(\cdot)}^{\alpha(\cdot), p, \theta}(\mathbb{R}^n)$ for $-n/q_i(\infty) < \alpha_i(0) \leq \alpha_i(\infty) < n(1 - 1/q_i(0))$. In the same year, B. Sultan et al. [22] introduced the grand Herz-Morrey spaces with variable exponents $M\dot{K}_{p, \theta, q(\cdot)}^{\alpha(\cdot), \lambda}(\mathbb{R}^n)$, and proved that the Riesz potential operator I^γ is bounded from spaces $M\dot{K}_{p, \theta, q_1(\cdot)}^{\alpha(\cdot), \lambda}(\mathbb{R}^n)$ into spaces $M\dot{K}_{p, \theta, q_2(\cdot)}^{\alpha(\cdot), \lambda}(\mathbb{R}^n)$. Subsequently, the boundedness of a variety of operators within these spaces has been established, including the fractional type Marcinkiewicz integral operators $\mathcal{M}_{\alpha, \rho, m}$ [3], higher order commutators of Marcinkiewicz integral $[b^m, \mu_\Phi]$ [23], among others. Moreover, more development of the grand spaces can be seen in [17, 18, 23, 24].

It is now necessary to confirm the symbols and nations of this article. C represents an constant independent of the main parameters, but it may be different from row to column. $p(\cdot)$ represents the conjugate exponents defined by $1/p(\cdot) + 1/p'(\cdot) = 1$. The expression $f \approx g$ means $C_1 f \leq g \leq C_2 f$. We also need denote $B_k = \{x \in \mathbb{R}^n : |x| \leq 2^k\}$ and $D_k = B_k \setminus B_{k-1}$.

2. Preliminaries

In this section, we recall some necessary definitions and notations. The definition of a bilinear θ -type generalized fractional integral operator $B\tilde{T}_{\beta, \theta}$ and its commutator $B\tilde{T}_{\beta, \theta, b_1, b_2}$ on $\mathbb{R}^n$ is as follows:

Definition 2.1. Let θ be a non-negative and non-decreasing function defined on $(0, \infty)$ and satisfy

$$\int_0^1 \frac{\theta(t)}{t} \log\left(\frac{1}{t}\right) dt < \infty. \quad (2.1)$$

A function $K_{\beta, \theta}(\cdot, \cdot, \cdot) \in L^1_{\text{loc}}((\mathbb{R}^n)^3 \setminus \{(x, x, x) : x \in \mathbb{R}^n\})$ is called a bilinear θ -type generalized fractional integral kernel if there exists a positive constant C such that,

(a) For all x, x', y_1, y_2 with $x \neq y_i$, $i = 1, 2$,

$$|K_{\beta, \theta}(x, y_1, y_2)| \leq \frac{C}{(|x - y_1|^{(1-\beta)} + |x - y_2|^{(1-\beta)})^{2n}}; \quad (2.2)$$

(b) For all x, y_1, y_2 with $|x - x'| \leq 1/2 \max\{|x - y_1|, |x - y_2|\}$,

$$|K_{\beta, \theta}(x, y_1, y_2) - K_{\beta, \theta}(x', y_1, y_2)| \leq \frac{C\theta\left(\frac{|x - x'|}{|x - y_1| + |x - y_2|}\right)}{(|x - y_1|^{(1-\beta)} + |x - y_2|^{(1-\beta)})^{2n}}; \quad (2.3)$$

(c) For all x, y_1, y'_1, y_2 with $|y_1 - y'_1| \leq 1/2 \max\{|x - y_1|, |x - y_2|\}$,

$$|K_{\beta, \theta}(x, y_1, y_2) - K_{\beta, \theta}(x, y'_1, y_2)| \leq \frac{C\theta\left(\frac{|y_1 - y'_1|}{|x - y_1| + |x - y_2|}\right)}{(|x - y_1|^{(1-\beta)} + |x - y_2|^{(1-\beta)})^{2n}}; \quad (2.4)$$

(d) For all x, y_1, y_2, y'_2 with $|y_2 - y'_2| \leq 1/2 \max\{|x - y_1|, |x - y_2|\}$,

$$|K_{\beta, \theta}(x, y_1, y_2) - K_{\beta, \theta}(x, y_1, y'_2)| \leq \frac{C\theta\left(\frac{|y_2 - y'_2|}{|x - y_1| + |x - y_2|}\right)}{(|x - y_1|^{(1-\beta)} + |x - y_2|^{(1-\beta)})^{2n}}, \quad (2.5)$$

where $\beta \in (0, \infty)$. A bilinear operator $B\tilde{T}_{\beta, \theta}$ is called a bilinear θ -type generalized fractional integral operator with kernels $K_{\beta, \theta}$ satisfying (2.2), (2.3), (2.4) and (2.5) if, for all $f_1, f_2 \in L_c^\infty(\mathbb{R}^n)$ with $x \notin (\text{supp}(f_1) \cap \text{supp}(f_2))$,

$$B\tilde{T}_{\beta, \theta}(f_1, f_2)(x) = \int_{\mathbb{R}^{2n}} K_{\beta, \theta}(x, y_1, y_2) f_1(y_1) f_2(y_2) dy_1 dy_2.$$

Definition 2.2. Given $b_1, b_2 \in \text{BMO}(\mathbb{R}^n)$, the commutators $B\tilde{T}_{\beta, \theta, b_1, b_2}$ generated by b_1, b_2 and the $B\tilde{T}_{\beta, \theta}$ is defined by

$$\begin{aligned} B\tilde{T}_{\beta, \theta, b_1, b_2}(f_1, f_2)(x) = & b_1(x) b_2(x) B\tilde{T}_{\beta, \theta}(f_1, f_2)(x) - b_1(x) B\tilde{T}_{\beta, \theta}(f_1, b_2(\cdot) f_2)(x) \\ & - b_2(x) B\tilde{T}_{\beta, \theta}(b_1(\cdot) f_1, f_2)(x) + B\tilde{T}_{\beta, \theta}(b_1(\cdot) f_1, b_2(\cdot) f_2)(x). \end{aligned}$$

Also, the following definitions are provided for the commutators $B\tilde{T}_{\beta, \theta, b_1}$ and $B\tilde{T}_{\beta, \theta, b_2}$:

$$\begin{aligned} B\tilde{T}_{\beta, \theta, b_1}(f_1, f_2)(x) &= b_1(x) B\tilde{T}_{\beta, \theta}(f_1, f_2)(x) - B\tilde{T}_{\beta, \theta}(b_1(\cdot) f_1, f_2)(x), \\ B\tilde{T}_{\beta, \theta, b_2}(f_1, f_2)(x) &= b_2(x) B\tilde{T}_{\beta, \theta}(f_1, f_2)(x) - B\tilde{T}_{\beta, \theta}(f_1, b_2(\cdot) f_2)(x). \end{aligned}$$

Remark 2.1. (a) For any $\delta \in (0, 1)$ and $t > 0$, if we take the $\theta(t) = t^\delta$, then the above bilinear θ -type generalized fractional integral operators $B\tilde{T}_{\beta, \theta}$ is just the bilinear generalized fractional integral operators $B\tilde{T}_\beta$ associated with kernel K_β (see [5]); (b) If we take $\beta = 0$ in (2.2), (2.3), (2.4) and (2.5), then the operators $B\tilde{T}_{\beta, \theta}$ is just the bilinear θ -type Calderón-Zygmund operators T_θ in [11]; (c) If we take $\theta = t^\delta$ and $\beta = 0$ in (2.2), (2.3), (2.4) and (2.5) then the operators $B\tilde{T}_{\beta, \theta}$ is just the bilinear Calderón-Zygmund operators T in [6].

The following basic facts about Lebesgue spaces with variable exponent are introduced by [25].

Definition 2.3. Let $p(\cdot) : \mathbb{R}^n \rightarrow (0, \infty]$ be a measurable function, for any $p(\cdot)$,

$$1 \leq p_- \leq p(x) \leq p_+ < \infty, \quad (2.6)$$

where

$$p_- = \text{ess inf}_{x \in \mathbb{R}^n} p(x) > 1; \quad p_+ = \text{ess sup}_{x \in \mathbb{R}^n} p(x) < \infty,$$

we denote $p(\cdot) \in \mathcal{P}(\mathbb{R}^n)$ if $p(\cdot)$ satisfy the above inequality, and if

$$I_{p(\cdot)}(f) = \int_{\mathbb{R}^n} |f(x)|^{p(x)} dx < \infty,$$

then $f \in L^{p(\cdot)}(\mathbb{R}^n)$. It is obvious that it is a Banach space being equipped with the norm

$$\|f\|_{L^{p(\cdot)}(\mathbb{R}^n)} = \inf \left\{ \lambda > 0 : I_{p(\cdot)}(f/\lambda) \leq 1 \right\}.$$

The following facts about log-Hölder continuous can be found in [20].

Definition 2.4. A function $g : \mathbb{R}^n \rightarrow \mathbb{R}$ is log-Hölder continuous if there exists a constant $C = C_{\log} > 0$ such that,

$$|g(x) - g(y)| \leq \frac{C}{\log(e + 1/|x - y|)}, \quad |x - y| \leq \frac{1}{2}, \quad x, y \in \mathbb{R}^n.$$

If the following two conditions

$$|g(x) - g_\infty| \leq \frac{C}{\log(e + |x|)}, \quad x \in \mathbb{R}^n \quad (2.7)$$

and

$$|g(x) - g(0)| \leq \frac{C}{\log(e + 1/|x|)}, \quad |x| \leq \frac{1}{2} \quad (2.8)$$

are satisfied. In such cases, the function $g(\cdot)$ is said to have a log-decay at infinity and at the origin, where $g_\infty = \lim_{x \rightarrow \infty} g(x)$.

In conclusion, we recall the definition of grand variable exponents Herz-Morrey spaces (see [22]).

Definition 2.5. Let $0 \leq \lambda < \infty$ and $\theta > 0$. Suppose that the variable exponents $q(\cdot) \in \mathcal{P}(\mathbb{R}^n)$, $\alpha(\cdot) \in L^\infty(\mathbb{R}^n)$ and $\alpha(\cdot) : \mathbb{R}^n \rightarrow \mathbb{R}$, then the homogeneous grand variable exponents Herz-Morrey spaces $M\dot{K}_{p,\theta,q(\cdot)}^{\alpha(\cdot),\lambda}(\mathbb{R}^n)$ is defined by

$$M\dot{K}_{p,\theta,q(\cdot)}^{\alpha(\cdot),\lambda}(\mathbb{R}^n) = \left\{ f \in L_{\text{loc}}^{q(\cdot)}(\mathbb{R}^n \setminus \{0\}) : \|f\|_{M\dot{K}_{p,\theta,q(\cdot)}^{\alpha(\cdot),\lambda}(\mathbb{R}^n)} < \infty \right\},$$

where

$$\|f\|_{M\dot{K}_{p,\theta,q(\cdot)}^{\alpha(\cdot),\lambda}(\mathbb{R}^n)} = \sup_{\epsilon > 0} \sup_{k_0 \in \mathbb{Z}} 2^{-k_0\lambda} \left(\epsilon^\theta \sum_{k=-\infty}^{k_0} \|2^{k\alpha(\cdot)} f \chi_k\|_{L^{q(\cdot)}(\mathbb{R}^n)}^{p(1+\epsilon)} \right)^{\frac{1}{p(1+\epsilon)}}. \quad (2.9)$$

The non-homogeneous grand variable exponents Herz-Morrey spaces is

$$MK_{p,\theta,q(\cdot)}^{\alpha(\cdot),\lambda}(\mathbb{R}^n) = \left\{ f \in L_{\text{loc}}^{q(\cdot)}(\mathbb{R}^n) : \|f\|_{MK_{p,\theta,q(\cdot)}^{\alpha(\cdot),\lambda}(\mathbb{R}^n)} < \infty \right\},$$

where

$$\|f\|_{MK_{p,\theta,q(\cdot)}^{\alpha(\cdot),\lambda}(\mathbb{R}^n)} = \sup_{\epsilon > 0} \sup_{k_0 \in \mathbb{N}_0} 2^{-k_0\lambda} \left(\epsilon^\theta \sum_{k=0}^{k_0} \|2^{k\alpha(\cdot)} f \chi_k\|_{L^{q(\cdot)}(\mathbb{R}^n)}^{p(1+\epsilon)} \right)^{\frac{1}{p(1+\epsilon)}}.$$

Remark 2.2. (a) If we take $\lambda = 0$ in (2.9), then the spaces $M\dot{K}_{p,\theta,q(\cdot)}^{\alpha(\cdot),\lambda}(\mathbb{R}^n)$ is just the homogeneous grand variable exponents Herz spaces $\dot{K}_{q(\cdot)}^{\alpha(\cdot),p,\theta}(\mathbb{R}^n)$.

(b) When $\epsilon = 0$, then $\dot{K}_{q(\cdot)}^{\alpha(\cdot),p,\theta}(\mathbb{R}^n) = \dot{K}_{q(\cdot)}^{\alpha(\cdot),p}(\mathbb{R}^n)$; when $\epsilon = 0$ and $\alpha(\cdot) \equiv \text{const}$, then $\dot{K}_{q(\cdot)}^{\alpha(\cdot),p,\theta}(\mathbb{R}^n) = \dot{K}_{q(\cdot)}^{\alpha,p}(\mathbb{R}^n)$.

3. Estimate for $B\tilde{T}_{\beta,\theta}$ and $B\tilde{T}_{\beta,\theta,b_1,b_2}$ on $M\dot{K}_{p,\theta,q(\cdot)}^{\alpha(\cdot),\lambda}$

In this section, the authors mainly consider the θ -type generalized fractional integral operators $B\tilde{T}_{\beta,\theta}$ and their commutators $B\tilde{T}_{\beta,\theta,b_1,b_2}$ generated by $b_1, b_2 \in \text{BMO}(\mathbb{R}^n)$ and the $B\tilde{T}_{\beta,\theta}$ are bounded on spaces $M\dot{K}_{p,\theta,q(\cdot)}^{\alpha(\cdot),\lambda}(\mathbb{R}^n)$ (or $M\dot{K}_{p,\theta,q(\cdot)}^{\alpha(\cdot),\lambda}(\mathbb{R}^n)$). The main results of this section are stated as follows:

Theorem 3.1. *Let $\theta > 1$, $\alpha(\cdot)$ and $q(\cdot)$ satisfy conditions (2.7) and (2.8). Suppose that $\frac{1}{q(x)} = \frac{1}{r_1(x)} + \frac{1}{r_2(x)}$, $\frac{1}{r_i(x)} = \frac{1}{q_i(x)} - \beta$ and $n\beta - \frac{n}{q_i(\infty)} < \alpha_i(0) \leq \alpha_i(\infty) < n - \frac{n}{q_i(0)}$ are established. Then there exists a positive constant C such that, for any $f_i \in M\dot{K}_{p_i,\theta,q_i(\cdot)}^{\alpha_i(\cdot),\lambda_i}(\mathbb{R}^n)$,*

$$\|B\tilde{T}_{\beta,\theta}(f_1, f_2)\|_{M\dot{K}_{p,\theta,q(\cdot)}^{\alpha(\cdot),\lambda}(\mathbb{R}^n)} \leq C \|f_1\|_{M\dot{K}_{p_1,\theta,q_1(\cdot)}^{\alpha_1(\cdot),\lambda_1}(\mathbb{R}^n)} \|f_2\|_{M\dot{K}_{p_2,\theta,q_2(\cdot)}^{\alpha_2(\cdot),\lambda_2}(\mathbb{R}^n)},$$

where $\lambda = \lambda_1 + \lambda_2$, $\alpha(x) = \alpha_1(x) + \alpha_2(x)$ and $\frac{1}{p} = \frac{1}{p_1} + \frac{1}{p_2}$.

Theorem 3.2. *Let $\theta > 1$, $b_1, b_2 \in \text{BMO}(\mathbb{R}^n)$, $\alpha(\cdot)$ and $q(\cdot)$ satisfy conditions (2.7) and (2.8). Suppose that $\frac{1}{q(x)} = \frac{1}{r_1(x)} + \frac{1}{r_2(x)}$, $\frac{1}{r_i(x)} = \frac{1}{q_i(x)} - \beta$ and $n\beta - \frac{n}{q_i(\infty)} < \alpha_i(0) \leq \alpha_i(\infty) < n - \frac{n}{q_i(0)}$ are established. Then there exists a positive constant C such that, for any $f_i \in M\dot{K}_{p_i,\theta,q_i(\cdot)}^{\alpha_i(\cdot),\lambda_i}(\mathbb{R}^n)$,*

$$\|B\tilde{T}_{\beta,\theta,b_1,b_2}(f_1, f_2)\|_{M\dot{K}_{p,\theta,q(\cdot)}^{\alpha(\cdot),\lambda}(\mathbb{R}^n)} \leq C \|f_1\|_{M\dot{K}_{p_1,\theta,q_1(\cdot)}^{\alpha_1(\cdot),\lambda_1}(\mathbb{R}^n)} \|f_2\|_{M\dot{K}_{p_2,\theta,q_2(\cdot)}^{\alpha_2(\cdot),\lambda_2}(\mathbb{R}^n)},$$

where $\lambda = \lambda_1 + \lambda_2$, $\alpha(x) = \alpha_1(x) + \alpha_2(x)$ and $\frac{1}{p} = \frac{1}{p_1} + \frac{1}{p_2}$.

Remark 3.1. The results of Theorems 3.1 and 3.2 can be applied to non-homogeneous Herz-Morrey spaces with variable exponents using similar arguments.

Furthermore, to prove the main theorems, we need to recall some basic facts.

Lemma 3.1. *Let $1 < p_i^- \leq p_i^+ < \infty$. Suppose that $q_i(\cdot)$ satisfy (2.7), (2.8) and $\frac{1}{q(x)} = \frac{1}{q_1(x)} + \frac{1}{q_2(x)} - 2\beta$. Then there exists a positive constant C such that, for all $f_i \in L^{q_i(\cdot)}(\mathbb{R}^n)$ ($i = 1, 2$),*

$$\|B\tilde{T}_{\beta,\theta}(f_1, f_1)\|_{L^{q(\cdot)}(\mathbb{R}^n)} \leq C \|f_1\|_{L^{q_1(\cdot)}(\mathbb{R}^n)} \|f_2\|_{L^{q_2(\cdot)}(\mathbb{R}^n)}.$$

Lemma 3.2. *Let $1 < p_i^- \leq p_i^+ < \infty$ and $b_i \in \text{BMO}(\mathbb{R}^n)$. Suppose that $q_i(\cdot)$ satisfy (2.7), (2.8) and $\frac{1}{q(x)} = \frac{1}{q_1(x)} + \frac{1}{q_2(x)} - 2\beta$. Then there exists a positive constant C such that, for all $f_i \in L^{q_i(\cdot)}(\mathbb{R}^n)$ ($i = 1, 2$),*

$$\|B\tilde{T}_{\beta,\theta,b_1,b_2}(f_1, f_1)\|_{L^{q(\cdot)}(\mathbb{R}^n)} \leq C \|f_1\|_{L^{q_1(\cdot)}(\mathbb{R}^n)} \|f_2\|_{L^{q_2(\cdot)}(\mathbb{R}^n)}.$$

Remark 3.2. The proofs for Lemma 2.1 and Lemma 2.2 are similar to the proofs of Theorem 1.1 and Theorem 1.2 in [7], respectively. Consequently, they are omitted here for brevity.

The following generalized Hölder's inequality can be found in [4].

Lemma 3.3. *If $p(\cdot), q(\cdot), r(\cdot) \in \mathcal{P}(\mathbb{R}^n)$ by $1/r(x) = 1/p(x) + 1/q(x)$, for any $f \in L^{p(\cdot)}(\mathbb{R}^n)$, $g \in L^{q(\cdot)}(\mathbb{R}^n)$ and $fg \in L^{r(\cdot)}(\mathbb{R}^n)$, such that*

$$\|fg\|_{L^{r(\cdot)}(\mathbb{R}^n)} \leq \|f\|_{L^{p(\cdot)}(\mathbb{R}^n)} \|g\|_{L^{q(\cdot)}(\mathbb{R}^n)}.$$

Finally, we recall the following characterisation of the spaces $\text{BMO}(\mathbb{R}^n)$ introduced in [10].

Lemma 3.4. *Let $b \in \text{BMO}(\mathbb{R}^n)$, $k > l$ and $q(\cdot) \in \mathcal{B}(\mathbb{R}^n)$, then*

$$\sup_{B \in \mathbb{R}^n} \frac{1}{\|\chi_B\|_{L^{q(\cdot)}(\mathbb{R}^n)}} \|(b - b_B)\chi_B\|_{L^{q(\cdot)}(\mathbb{R}^n)} \approx \|b\|_*$$

and

$$\|(b - b_{B_l})\chi_{B_k}\|_{L^{q(\cdot)}(\mathbb{R}^n)} \leq C(k - l)\|b\|_* \|\chi_{B_k}\|_{L^{q(\cdot)}(\mathbb{R}^n)}.$$

Proof of Theorem 3.1. Let $f_i \in M\dot{K}_{p_i, \theta, q_i(\cdot)}^{\alpha_i(\cdot), \lambda_i}(\mathbb{R}^n)$, and decompose

$$f_1(x) = \sum_{l=-\infty}^{\infty} f_1(x)\chi_l(x) = \sum_{l=-\infty}^{\infty} f_1^l(x); f_2(x) = \sum_{j=-\infty}^{\infty} f_2(x)\chi_j(x) = \sum_{j=-\infty}^{\infty} f_2^j(x).$$

By Minkowski's inequality, we obtain

$$\begin{aligned} & \|B\tilde{T}_{\beta, \theta}(f_1, f_2)\|_{M\dot{K}_{p, \theta, q(\cdot)}^{\alpha(\cdot), \lambda}(\mathbb{R}^n)} \\ & \leq \sup_{\epsilon > 0} \sup_{k_0 \in \mathbb{Z}} 2^{-k_0\lambda} \left(\epsilon^\theta \sum_{k=-\infty}^{k_0} \left(\sum_{l=-\infty}^{\infty} \sum_{j=-\infty}^{\infty} \|2^{k\alpha(\cdot)} B\tilde{T}_{\beta, \theta}(f_1^l, f_2^j)\chi_k\|_{q(\cdot)} \right)^{p(1+\epsilon)} \right)^{\frac{1}{p(1+\epsilon)}} \\ & \leq \sup_{\epsilon > 0} \sup_{k_0 \in \mathbb{Z}} 2^{-k_0\lambda} \left(\epsilon^\theta \sum_{k=-\infty}^{k_0} \left(\sum_{l=-\infty}^{k-2} \sum_{j=-\infty}^{k-2} \|2^{k\alpha(\cdot)} B\tilde{T}_{\beta, \theta}(f_1^l, f_2^j)\chi_k\|_{q(\cdot)} \right)^{p(1+\epsilon)} \right)^{\frac{1}{p(1+\epsilon)}} \\ & \quad + \sup_{\epsilon > 0} \sup_{k_0 \in \mathbb{Z}} 2^{-k_0\lambda} \left(\epsilon^\theta \sum_{k=-\infty}^{k_0} \left(\sum_{l=-\infty}^{k-2} \sum_{j=k-1}^{k+1} \|2^{k\alpha(\cdot)} B\tilde{T}_{\beta, \theta}(f_1^l, f_2^j)\chi_k\|_{q(\cdot)} \right)^{p(1+\epsilon)} \right)^{\frac{1}{p(1+\epsilon)}} \\ & \quad + \sup_{\epsilon > 0} \sup_{k_0 \in \mathbb{Z}} 2^{-k_0\lambda} \left(\epsilon^\theta \sum_{k=-\infty}^{k_0} \left(\sum_{l=-\infty}^{k-2} \sum_{j=k+2}^{\infty} \|2^{k\alpha(\cdot)} B\tilde{T}_{\beta, \theta}(f_1^l, f_2^j)\chi_k\|_{q(\cdot)} \right)^{p(1+\epsilon)} \right)^{\frac{1}{p(1+\epsilon)}} \\ & \quad + \sup_{\epsilon > 0} \sup_{k_0 \in \mathbb{Z}} 2^{-k_0\lambda} \left(\epsilon^\theta \sum_{k=-\infty}^{k_0} \left(\sum_{l=k-1}^{k+1} \sum_{j=-\infty}^{k-2} \|2^{k\alpha(\cdot)} B\tilde{T}_{\beta, \theta}(f_1^l, f_2^j)\chi_k\|_{q(\cdot)} \right)^{p(1+\epsilon)} \right)^{\frac{1}{p(1+\epsilon)}} \\ & \quad + \sup_{\epsilon > 0} \sup_{k_0 \in \mathbb{Z}} 2^{-k_0\lambda} \left(\epsilon^\theta \sum_{k=-\infty}^{k_0} \left(\sum_{l=k-1}^{k+1} \sum_{j=k-1}^{k+1} \|2^{k\alpha(\cdot)} B\tilde{T}_{\beta, \theta}(f_1^l, f_2^j)\chi_k\|_{q(\cdot)} \right)^{p(1+\epsilon)} \right)^{\frac{1}{p(1+\epsilon)}} \\ & \quad + \sup_{\epsilon > 0} \sup_{k_0 \in \mathbb{Z}} 2^{-k_0\lambda} \left(\epsilon^\theta \sum_{k=-\infty}^{k_0} \left(\sum_{l=k-1}^{k+1} \sum_{j=k+2}^{\infty} \|2^{k\alpha(\cdot)} B\tilde{T}_{\beta, \theta}(f_1^l, f_2^j)\chi_k\|_{q(\cdot)} \right)^{p(1+\epsilon)} \right)^{\frac{1}{p(1+\epsilon)}} \\ & \quad + \sup_{\epsilon > 0} \sup_{k_0 \in \mathbb{Z}} 2^{-k_0\lambda} \left(\epsilon^\theta \sum_{k=-\infty}^{k_0} \left(\sum_{l=k+2}^{\infty} \sum_{j=-\infty}^{k-2} \|2^{k\alpha(\cdot)} B\tilde{T}_{\beta, \theta}(f_1^l, f_2^j)\chi_k\|_{q(\cdot)} \right)^{p(1+\epsilon)} \right)^{\frac{1}{p(1+\epsilon)}} \\ & \quad + \sup_{\epsilon > 0} \sup_{k_0 \in \mathbb{Z}} 2^{-k_0\lambda} \left(\epsilon^\theta \sum_{k=-\infty}^{k_0} \left(\sum_{l=k+2}^{\infty} \sum_{j=k-1}^{k+1} \|2^{k\alpha(\cdot)} B\tilde{T}_{\beta, \theta}(f_1^l, f_2^j)\chi_k\|_{q(\cdot)} \right)^{p(1+\epsilon)} \right)^{\frac{1}{p(1+\epsilon)}} \end{aligned}$$

$$+ \sup_{\epsilon > 0} \sup_{k_0 \in \mathbb{Z}} 2^{-k_0 \lambda} \left(\epsilon^\theta \sum_{k=-\infty}^{k_0} \left(\sum_{l=k+2}^{\infty} \sum_{j=k+2}^{\infty} \left\| 2^{k\alpha(\cdot)} B\tilde{T}_{\beta, \theta}(f_1^l, f_2^j) \chi_k \right\|_{q(\cdot)} \right)^{p(1+\epsilon)} \right)^{\frac{1}{p(1+\epsilon)}} \\ = E_1 + E_2 + E_3 + E_4 + E_5 + E_6 + E_7 + E_8 + E_9.$$

It is necessary to estimate E_1, E_2, E_3, E_5, E_6 and E_9 , since E_4, E_7 and E_8 can be obtained by following a similar methodology to that used for E_2, E_3 and E_6 , respectively. Then, For E_1 , we first set $l, j \leq k-2, x \in D_k, y_1 \in D_l$ and $y_2 \in D_j$, such that $|x - y_i| \geq C2^k$. Then, for $x \in D_k$, we can obtain

$$\begin{aligned} \left| B\tilde{T}_{\beta, \theta}(f_1^l, f_2^j)(x) \right| &\leq C2^{-2n} 2^{-2nk(1-\beta)} \int_{\mathbb{R}^n} \int_{\mathbb{R}^n} |f_1^l(y_1)| |f_2^j(y_2)| dy_1 dy_2 \\ &\leq C2^{-2nk(1-\beta)} \|f_1^l\|_{L^1} \|f_2^j\|_{L^1}. \end{aligned} \quad (3.1)$$

From the Minkowski's inequality, it follows that

$$\begin{aligned} &\sup_{\epsilon > 0} \sup_{k_0 \in \mathbb{Z}} 2^{-k_0 \lambda} \left(\epsilon^\theta \sum_{k=-\infty}^{k_0} \left(\sum_{l=-\infty}^{k-2} \sum_{j=-\infty}^{k-2} \left\| 2^{k\alpha(\cdot)} B\tilde{T}_{\beta, \theta}(f_1^l, f_2^j) \chi_k \right\|_{q(\cdot)} \right)^{p(1+\epsilon)} \right)^{\frac{1}{p(1+\epsilon)}} \\ &\leq \sup_{\epsilon > 0} \sup_{k_0 \in \mathbb{Z}} 2^{-k_0 \lambda} \left(\epsilon^\theta \sum_{k=-\infty}^{-1} 2^{k\alpha(0)} \left(\sum_{l=-\infty}^{k-2} \sum_{j=-\infty}^{k-2} \left\| B\tilde{T}_{\beta, \theta}(f_1^l, f_2^j) \chi_k \right\|_{q(\cdot)} \right)^{p(1+\epsilon)} \right)^{\frac{1}{p(1+\epsilon)}} \\ &\quad + \sup_{\epsilon > 0} \sup_{k_0 \in \mathbb{Z}} 2^{-k_0 \lambda} \left(\epsilon^\theta \sum_{k=0}^{k_0} 2^{k\alpha(\infty)} \left(\sum_{l=-\infty}^{-1} \sum_{j=-\infty}^{-1} \left\| B\tilde{T}_{\beta, \theta}(f_1^l, f_2^j) \chi_k \right\|_{q(\cdot)} \right)^{p(1+\epsilon)} \right)^{\frac{1}{p(1+\epsilon)}} \\ &\quad + \sup_{\epsilon > 0} \sup_{k_0 \in \mathbb{Z}} 2^{-k_0 \lambda} \left(\epsilon^\theta \sum_{k=0}^{k_0} 2^{k\alpha(\infty)} \left(\sum_{l=-\infty}^{-1} \sum_{j=0}^{k-2} \left\| B\tilde{T}_{\beta, \theta}(f_1^l, f_2^j) \chi_k \right\|_{q(\cdot)} \right)^{p(1+\epsilon)} \right)^{\frac{1}{p(1+\epsilon)}} \\ &\quad + \sup_{\epsilon > 0} \sup_{k_0 \in \mathbb{Z}} 2^{-k_0 \lambda} \left(\epsilon^\theta \sum_{k=0}^{k_0} 2^{k\alpha(\infty)} \left(\sum_{l=0}^{k-2} \sum_{j=-\infty}^{-1} \left\| B\tilde{T}_{\beta, \theta}(f_1^l, f_2^j) \chi_k \right\|_{q(\cdot)} \right)^{p(1+\epsilon)} \right)^{\frac{1}{p(1+\epsilon)}} \\ &\quad + \sup_{\epsilon > 0} \sup_{k_0 \in \mathbb{Z}} 2^{-k_0 \lambda} \left(\epsilon^\theta \sum_{k=0}^{k_0} 2^{k\alpha(\infty)} \left(\sum_{l=0}^{k-2} \sum_{j=0}^{k-2} \left\| B\tilde{T}_{\beta, \theta}(a, f_2^j) \chi_k \right\|_{q(\cdot)} \right)^{p(1+\epsilon)} \right)^{\frac{1}{p(1+\epsilon)}} \\ &= E_{11} + E_{12} + E_{13} + E_{14} + E_{15}. \end{aligned}$$

For E_{11} , by (3.1) and Lemma 3.3, we assume $a_i = \frac{n}{q_i(0)} - n + \alpha_i(0)$ and consider

$$\begin{aligned} E_{11} &\leq C \sup_{\epsilon > 0} \sup_{k_0 \in \mathbb{Z}} 2^{-k_0 \lambda_1} \left(\epsilon^\theta \sum_{k=-\infty}^{-1} \left(\sum_{l=-\infty}^{k-2} 2^{l\alpha_1(0)} 2^{a_1(k-l)} \|f_1^l\|_{q_1(\cdot)} \right)^{p_1(1+\epsilon)} \right)^{\frac{1}{p_1(1+\epsilon)}} \\ &\quad \times \sup_{\epsilon > 0} \sup_{k_0 \in \mathbb{Z}} 2^{-k_0 \lambda_2} \left(\epsilon^\theta \sum_{k=-\infty}^{-1} \left(\sum_{j=-\infty}^{k-2} 2^{j\alpha_2(0)} 2^{a_2(k-j)} \|f_2^j\|_{q_2(\cdot)} \right)^{p_2(1+\epsilon)} \right)^{\frac{1}{p_2(1+\epsilon)}} \\ &= E_{11}^1 \times E_{11}^2. \end{aligned}$$

For E_{11}^1 , we have established that it satisfies the inequality

$$E_{11}^1 \leq C \sup_{\epsilon > 0} \sup_{k_0 \in \mathbb{Z}} 2^{-k_0 \lambda_1} \left(\epsilon^\theta \sum_{k=-\infty}^{-1} \left(\sum_{l=-\infty}^{k-2} 2^{l\alpha_1(0)p_1(1+\epsilon)} 2^{\frac{a_1(k-l)p_1(1+\epsilon)}{2}} \|f_1^l\|_{q_1(\cdot)}^{p_1(1+\epsilon)} \right) \right)$$

$$\times \left(\sum_{l=-\infty}^{k-1} 2^{\frac{a_1(k-l)p'_1(1+\epsilon)}{2}} \right)^{\frac{p_1(1+\epsilon)}{p'_1(1+\epsilon)}}^{\frac{1}{p_1(1+\epsilon)}} \\ \leq C \|f_1\|_{M\dot{K}_{p_1, \theta, q_1(\cdot)}^{\alpha_1(\cdot), \lambda_1}(\mathbb{R}^n)}.$$

With an argument similar to that used in the estimate for E_{11}^1 , it is easy to get

$$E_{11}^2 \leq C \|f_2\|_{M\dot{K}_{p_2, \theta, q_2(\cdot)}^{\alpha_2(\cdot), \lambda_2}(\mathbb{R}^n)}.$$

For E_{12} , since $q_i(0) \leq q_i(\infty)$ and $\alpha_i(0) \leq \alpha_i(\infty)$ are given, we assume $b_i = \frac{n}{q_i(0)} - n + \alpha_i(\infty)$ and deduce

$$E_{12} \leq C \sup_{\epsilon > 0} \sup_{k_0 \in \mathbb{Z}} 2^{-k_0 \lambda_1} \left(\epsilon^\theta \sum_{k=0}^{k_0} \left(\sum_{l=-\infty}^{-1} 2^{l \alpha_1(0)} 2^{b_1(k-l)} \|f_1^l\|_{q_1(\cdot)} \right)^{p_1(1+\epsilon)} \right)^{\frac{1}{p_1(1+\epsilon)}} \\ \times \sup_{\epsilon > 0} \sup_{k_0 \in \mathbb{Z}} 2^{-k_0 \lambda_2} \left(\epsilon^\theta \sum_{k=0}^{k_0} \left(\sum_{j=-\infty}^{-1} 2^{j \alpha_2(0)} 2^{b_2(k-j)} \|f_2^j\|_{q_2(\cdot)} \right)^{p_2(1+\epsilon)} \right)^{\frac{1}{p_2(1+\epsilon)}} \\ = E_{12}^1 \times E_{12}^2.$$

The estimates for E_{12}^1 , E_{12}^2 and E_{11}^1 are similar, hence, it is easy to get

$$E_{12}^1 E_{12}^2 \leq C \|f_1\|_{M\dot{K}_{p_1, \theta, q_1(\cdot)}^{\alpha_1(\cdot), \lambda_1}(\mathbb{R}^n)} \|f_2\|_{M\dot{K}_{p_2, \theta, q_2(\cdot)}^{\alpha_2(\cdot), \lambda_2}(\mathbb{R}^n)}.$$

For E_{13} , by (3.1), $q_1(0) \leq q_1(\infty)$, $\alpha_1(0) \leq \alpha_1(\infty)$ and $c_i = \frac{n}{q_2(\infty)} - n + \alpha_i(\infty)$, we get

$$E_{13} \leq C \sup_{\epsilon > 0} \sup_{k_0 \in \mathbb{Z}} 2^{-k_0 \lambda_1} \left(\epsilon^\theta \sum_{k=0}^{k_0} \left(\sum_{l=-\infty}^{-1} 2^{l \alpha_1(0)} 2^{b_1(k-l)} \|f_1^l\|_{q_1(\cdot)} \right)^{p_1(1+\epsilon)} \right)^{\frac{1}{p_1(1+\epsilon)}} \\ \times \sup_{\epsilon > 0} \sup_{k_0 \in \mathbb{Z}} 2^{-k_0 \lambda_2} \left(\epsilon^\theta \sum_{k=0}^{k_0} \left(\sum_{j=0}^{k-2} 2^{j \alpha_2(\infty)} 2^{c_2(k-j)} \|f_2^j\|_{q_2(\cdot)} \right)^{p_2(1+\epsilon)} \right)^{\frac{1}{p_2(1+\epsilon)}} \\ = E_{13}^1 \times E_{13}^2.$$

With a way similar to that used in the estimate for E_{11} , it is not difficult to get

$$E_{13} \leq C \|f_1\|_{M\dot{K}_{p_1, \theta, q_1(\cdot)}^{\alpha_1(\cdot), \lambda_1}(\mathbb{R}^n)} \|f_2\|_{M\dot{K}_{p_2, \theta, q_2(\cdot)}^{\alpha_2(\cdot), \lambda_2}(\mathbb{R}^n)}.$$

From the symmetry, by E_{13} , we get

$$E_{14} \leq C \|f_1\|_{M\dot{K}_{p_1, \theta, q_1(\cdot)}^{\alpha_1(\cdot), \lambda_1}(\mathbb{R}^n)} \|f_2\|_{M\dot{K}_{p_2, \theta, q_2(\cdot)}^{\alpha_2(\cdot), \lambda_2}(\mathbb{R}^n)}.$$

For E_{15} , from (3.1), Lemma 3.3 and $\frac{n}{q_i(\infty)} - n + \alpha_i(\infty) < 0$, just like E_{11} , we obtain

$$E_{15} \leq C \|f_1\|_{M\dot{K}_{p_1, \theta, q_1(\cdot)}^{\alpha_1(\cdot), \lambda_1}(\mathbb{R}^n)} \|f_2\|_{M\dot{K}_{p_2, \theta, q_2(\cdot)}^{\alpha_2(\cdot), \lambda_2}(\mathbb{R}^n)}.$$

For E_2 , we set $l \leq k-2$, $k-1 \leq j \leq k+1$, $x \in D_k$, $y_1 \in D_l$ and $y_2 \in D_j$, such that $|x - y_1| \geq C2^k$ and $|x - y_2| \geq 0$. For $x \in D_k$, we deduce

$$\begin{aligned} \left| B\tilde{T}_{\beta,\theta}(f_1^l, f_2^j)(x) \right| &\leq C2^{-2nk(1-\beta)} \int_{\mathbb{R}^n} \int_{\mathbb{R}^n} |f_1^l(y_1)| |f_2^j(y_2)| dy_1 dy_2 \\ &\leq C2^{-2nk(1-\beta)} \|f_1^l\|_{L^1} \|f_2^j\|_{L^1}, \end{aligned} \quad (3.2)$$

by this and the Minkowski's inequality, write

$$\begin{aligned} &\sup_{\epsilon > 0} \sup_{k_0 \in \mathbb{Z}} 2^{-k_0\lambda} \left(\epsilon^\theta \sum_{k=-\infty}^{k_0} \left(\sum_{l=-\infty}^{k-2} \sum_{j=k-1}^{k+1} \left\| 2^{k\alpha(\cdot)} B\tilde{T}_{\beta,\theta}(f_1^l, f_2^j) \chi_k \right\|_{q(\cdot)} \right)^{p(1+\epsilon)} \right)^{\frac{1}{p(1+\epsilon)}} \\ &\leq \sup_{\epsilon > 0} \sup_{k_0 \in \mathbb{Z}} 2^{-k_0\lambda} \left(\epsilon^\theta \sum_{k=-\infty}^{-1} 2^{k\alpha(0)} \left(\sum_{l=-\infty}^{k-2} \sum_{j=k-1}^{k+1} \left\| B\tilde{T}_{\beta,\theta}(f_1^l, f_2^j) \chi_k \right\|_{q(\cdot)} \right)^{p(1+\epsilon)} \right)^{\frac{1}{p(1+\epsilon)}} \\ &\quad + \sup_{\epsilon > 0} \sup_{k_0 \in \mathbb{Z}} 2^{-k_0\lambda} \left(\epsilon^\theta \sum_{k=0}^{k_0} 2^{k\alpha(\infty)} \left(\sum_{l=-\infty}^{-1} \sum_{j=k-1}^{k+1} \left\| B\tilde{T}_{\beta,\theta}(f_1^l, f_2^j) \chi_k \right\|_{q(\cdot)} \right)^{p(1+\epsilon)} \right)^{\frac{1}{p(1+\epsilon)}} \\ &\quad + \sup_{\epsilon > 0} \sup_{k_0 \in \mathbb{Z}} 2^{-k_0\lambda} \left(\epsilon^\theta \sum_{k=0}^{k_0} 2^{k\alpha(\infty)} \left(\sum_{l=0}^{k-2} \sum_{j=k-1}^{k+1} \left\| B\tilde{T}_{\beta,\theta}(f_1^l, f_2^j) \chi_k \right\|_{q(\cdot)} \right)^{p(1+\epsilon)} \right)^{\frac{1}{p(1+\epsilon)}} \\ &= E_{21} + E_{22} + E_{23}. \end{aligned}$$

For E_{21} , by (3.2) and Lemma 3.3, we conclude

$$\begin{aligned} E_{21} &\leq C \sup_{\epsilon > 0} \sup_{k_0 \in \mathbb{Z}} 2^{-k_0\lambda_1} \left(\epsilon^\theta \sum_{k=-\infty}^{-1} \left(\sum_{l=-\infty}^{k-2} 2^{l\alpha_1(0)} 2^{a_1(k-l)} \|f_1^l\|_{q_1(\cdot)} \right)^{p_1(1+\epsilon)} \right)^{\frac{1}{p_1(1+\epsilon)}} \\ &\quad \times \sup_{\epsilon > 0} \sup_{k_0 \in \mathbb{Z}} 2^{-k_0\lambda_2} \left(\epsilon^\theta \sum_{k=-\infty}^{-1} \left(\sum_{j=k-1}^{k+1} 2^{k\alpha_2(0)} \|f_2^j\|_{q_2(\cdot)} \right)^{p_2(1+\epsilon)} \right)^{\frac{1}{p_2(1+\epsilon)}} \\ &= E_{21}^1 \times E_{21}^2. \end{aligned}$$

Since E_{21}^1 and E_{11}^1 are the same. Thus, we only turn E_{21}^2 ,

$$\begin{aligned} E_{21}^2 &\leq C \sup_{\epsilon > 0} \sup_{k_0 \in \mathbb{Z}} 2^{-k_0\lambda_2} \left(\epsilon^\theta \sum_{k=-\infty}^{-1} 2^{k\alpha_2(0)p_2(1+\epsilon)} \|f_2^k\|_{q_2(\cdot)}^{p_2(1+\epsilon)} \right)^{\frac{1}{p_2(1+\epsilon)}} \\ &\leq C \|f_2\|_{M\dot{K}_{p_2, \theta, q_2(\cdot)}^{\alpha_2(\cdot), \lambda_2}(\mathbb{R}^n)}. \end{aligned}$$

For E_{22} , by (3.2) and Lemma 3.3, we get

$$\begin{aligned} E_{22} &\leq C \sup_{\epsilon > 0} \sup_{k_0 \in \mathbb{Z}} 2^{-k_0\lambda_1} \left(\epsilon^\theta \sum_{k=0}^{k_0} \left(\sum_{l=-\infty}^{-1} 2^{l\alpha_1(0)} 2^{b_1(k-l)} \|f_1^l\|_{q_1(\cdot)} \right)^{p_1(1+\epsilon)} \right)^{\frac{1}{p_1(1+\epsilon)}} \\ &\quad \times \sup_{\epsilon > 0} \sup_{k_0 \in \mathbb{Z}} 2^{-k_0\lambda_2} \left(\epsilon^\theta \sum_{k=0}^{k_0} \left(\sum_{j=k-1}^{k+1} 2^{k\alpha_2(\infty)} \|f_2^j\|_{q_2(\cdot)} \right)^{p_2(1+\epsilon)} \right)^{\frac{1}{p_2(1+\epsilon)}} \\ &= E_{22}^1 \times E_{22}^2. \end{aligned}$$

Since E_{22}^2 and E_{21}^2 are similar, and $E_{22}^1 = E_{12}^1$, it is easy to get

$$E_{22} \leq C \|f_1\|_{M\dot{K}_{p_1, \theta, q_1(\cdot)}^{\alpha_1(\cdot), \lambda_1}(\mathbb{R}^n)} \|f_2\|_{M\dot{K}_{p_2, \theta, q_2(\cdot)}^{\alpha_2(\cdot), \lambda_2}(\mathbb{R}^n)}.$$

For E_{23} , by (3.2), we obtain

$$\begin{aligned} E_{23} &\leq C \sup_{\epsilon > 0} \sup_{k_0 \in \mathbb{Z}} 2^{-k_0 \lambda_1} \left(\epsilon^\theta \sum_{k=0}^{k_0} \left(\sum_{l=0}^{k-2} 2^{l\alpha_1(\infty)} 2^{c_1(k-l)} \|f_1^l\|_{q_1(\cdot)} \right)^{p_1(1+\epsilon)} \right)^{\frac{1}{p_1(1+\epsilon)}} \\ &\quad \times \sup_{\epsilon > 0} \sup_{k_0 \in \mathbb{Z}} 2^{-k_0 \lambda_2} \left(\epsilon^\theta \sum_{k=0}^{k_0} \left(\sum_{j=k-1}^{k+1} 2^{j\alpha_2(\infty)} \|f_2^j\|_{q_2(\cdot)} \right)^{p_2(1+\epsilon)} \right)^{\frac{1}{p_2(1+\epsilon)}} \\ &= E_{23}^1 \times E_{23}^2, \end{aligned}$$

we can find $E_{23}^1 = E_{15}^1$ and $E_{23}^2 = E_{22}^2$. Therefore, we obtain an estimate of E_{23} .

For E_3 , we set $l \leq k-2$, $j \geq k+2$, $x \in D_k$, $y_1 \in D_l$ and $y_2 \in D_j$, such that $|x - y_1| \geq C2^k$ and $|x - y_2| \geq C2^j$. For $x \in D_k$, we obtain

$$\begin{aligned} |B\tilde{T}_{\beta, \theta}(f_1^l, f_2^j)(x)| &\leq C \int_{\mathbb{R}^n} \int_{\mathbb{R}^n} \frac{|f_1^l(y_1)| |f_2^j(y_2)|}{\left(2^{k(1-\beta)} + 2^{j(1-\beta)}\right)^{2n}} dy_1 dy_2 \\ &\leq C 2^{-nk(1-\beta)} 2^{-nj(1-\beta)} \|f_1^l\|_{L^1} \|f_2^j\|_{L^1}, \end{aligned} \quad (3.3)$$

write

$$\begin{aligned} &\sup_{\epsilon > 0} \sup_{k_0 \in \mathbb{Z}} 2^{-k_0 \lambda} \left(\epsilon^\theta \sum_{k=-\infty}^{k_0} \left(\sum_{l=-\infty}^{k-2} \sum_{j=k+2}^{\infty} \left\| 2^{k\alpha(\cdot)} B\tilde{T}_{\beta, \theta}(f_1^l, f_2^j) \chi_k \right\|_{q(\cdot)} \right)^{p(1+\epsilon)} \right)^{\frac{1}{p(1+\epsilon)}} \\ &\leq \sup_{\epsilon > 0} \sup_{k_0 \in \mathbb{Z}} 2^{-k_0 \lambda} \left(\epsilon^\theta \sum_{k=-\infty}^{-1} 2^{k\alpha(0)} \left(\sum_{l=-\infty}^{k-2} \sum_{j=k+2}^{\infty} \left\| B\tilde{T}_{\beta, \theta}(f_1^l, f_2^j) \chi_k \right\|_{q(\cdot)} \right)^{p(1+\epsilon)} \right)^{\frac{1}{p(1+\epsilon)}} \\ &\quad + \sup_{\epsilon > 0} \sup_{k_0 \in \mathbb{Z}} 2^{-k_0 \lambda} \left(\epsilon^\theta \sum_{k=0}^{k_0} 2^{k\alpha(\infty)} \left(\sum_{l=-\infty}^{-1} \sum_{j=k+2}^{\infty} \left\| B\tilde{T}_{\beta, \theta}(f_1^l, f_2^j) \chi_k \right\|_{q(\cdot)} \right)^{p(1+\epsilon)} \right)^{\frac{1}{p(1+\epsilon)}} \\ &\quad + \sup_{\epsilon > 0} \sup_{k_0 \in \mathbb{Z}} 2^{-k_0 \lambda} \left(\epsilon^\theta \sum_{k=0}^{k_0} 2^{k\alpha(\infty)} \left(\sum_{l=1}^{k-2} \sum_{j=k+2}^{\infty} \left\| B\tilde{T}_{\beta, \theta}(f_1^l, f_2^j) \chi_k \right\|_{q(\cdot)} \right)^{p(1+\epsilon)} \right)^{\frac{1}{p(1+\epsilon)}} \\ &= E_{31} + E_{32} + E_{33}. \end{aligned}$$

For E_{31} , we obtain

$$\begin{aligned} E_{31} &\leq C \sup_{\epsilon > 0} \sup_{k_0 \in \mathbb{Z}} 2^{-k_0 \lambda_1} \left(\epsilon^\theta \sum_{k=-\infty}^{-1} \left(\sum_{l=-\infty}^{k-2} 2^{l\alpha_1(0)} 2^{a_1(k-l)} \|f_1^l\|_{q_1(\cdot)} \right)^{p_1(1+\epsilon)} \right)^{\frac{1}{p_1(1+\epsilon)}} \\ &\quad \times \sup_{\epsilon > 0} \sup_{k_0 \in \mathbb{Z}} 2^{-k_0 \lambda_2} \left(\epsilon^\theta \sum_{k=-\infty}^{-1} \left(\sum_{j=k+2}^{\infty} 2^{k\alpha_2(0)} \right. \right. \\ &\quad \left. \left. \times 2^{(k-j)\left(\frac{n}{q_2(\infty)} - n\beta\right)} \|f_2^j\|_{q_2(\cdot)} \right)^{p_2(1+\epsilon)} \right)^{\frac{1}{p_2(1+\epsilon)}} \end{aligned}$$

$$= E_{31}^1 E_{31}^2.$$

It is easy to see that $E_{31}^1 = E_{11}^1$, hence, we only estimate E_{31}^2 . Write

$$\begin{aligned} E_{31}^2 &\leq C \sup_{\epsilon > 0} \sup_{k_0 \in \mathbb{Z}} 2^{-k_0 \lambda_2} \left(\epsilon^\theta \sum_{k=-\infty}^{-1} \left(\sum_{j=k+2}^{-1} 2^{j\alpha_2(0)} 2^{d_2(k-j)} \|f_2^j\|_{q_2(\cdot)} \right)^{p_2(1+\epsilon)} \right)^{\frac{1}{p_2(1+\epsilon)}} \\ &\quad + \sup_{\epsilon > 0} \sup_{k_0 \in \mathbb{Z}} 2^{-k_0 \lambda_2} \left(\epsilon^\theta \sum_{k=-\infty}^{-1} \left(\sum_{j=0}^{\infty} 2^{j\alpha_2(\infty)} 2^{d_2(k-j)} \|f_2^j\|_{q_2(\cdot)} \right)^{p_2(1+\epsilon)} \right)^{\frac{1}{p_2(1+\epsilon)}} \\ &= D_1 + D_2, \end{aligned}$$

where $d_i = \frac{n}{q_i(\infty)} - n\beta + \alpha_i(0)$. With an argument similar to that used in E_{11}^1 , it is easy to get E_{31}^2 . For E_{32} , by (3.3), $q_1(0) \leq q_1(\infty)$ and $e_i = \frac{n}{q_i(\infty)} - n\beta + \alpha_i(\infty)$, write

$$\begin{aligned} E_{32} &\leq C \sup_{\epsilon > 0} \sup_{k_0 \in \mathbb{Z}} 2^{-k_0 \lambda_1} \left(\epsilon^\theta \sum_{k=0}^{k_0} \left(\sum_{l=-\infty}^{-1} 2^{l\alpha_1(0)} 2^{b_1(k-l)} \|f_1^l\|_{q_1(\cdot)} \right)^{p_1(1+\epsilon)} \right)^{\frac{1}{p_1(1+\epsilon)}} \\ &\quad \times \sup_{\epsilon > 0} \sup_{k_0 \in \mathbb{Z}} 2^{-k_0 \lambda_2} \left(\epsilon^\theta \sum_{k=0}^{k_0} \left(\sum_{j=k+2}^{\infty} 2^{j\alpha_2(\infty)} 2^{e_2(k-j)} \|f_2^j\|_{q_2(\cdot)} \right)^{p_2(1+\epsilon)} \right)^{\frac{1}{p_2(1+\epsilon)}} \\ &= E_{32}^1 E_{32}^2. \end{aligned}$$

It is obvious to find that $E_{32}^1 = E_{12}^1$ and the estimate of E_{32}^2 is similar to D_2 . Hence, we have

$$E_{32} \leq C \|f_1\|_{M\dot{K}_{p_1, \theta, q_1(\cdot)}^{\alpha_1(\cdot), \lambda_1}(\mathbb{R}^n)} \|f_2\|_{M\dot{K}_{p_2, \theta, q_2(\cdot)}^{\alpha_2(\cdot), \lambda_2}(\mathbb{R}^n)}.$$

For E_{33} , just like E_{32} , we obtain

$$\begin{aligned} E_{33} &\leq C \sup_{\epsilon > 0} \sup_{k_0 \in \mathbb{Z}} 2^{-k_0 \lambda_1} \left(\epsilon^\theta \sum_{k=0}^{k_0} \left(\sum_{l=0}^{k-2} 2^{l\alpha_1(\infty)} 2^{c_1(k-l)} \|f_1^l\|_{q_1(\cdot)} \right)^{p_1(1+\epsilon)} \right)^{\frac{1}{p_1(1+\epsilon)}} \\ &\quad \times \sup_{\epsilon > 0} \sup_{k_0 \in \mathbb{Z}} 2^{-k_0 \lambda_2} \left(\epsilon^\theta \sum_{k=0}^{k_0} \left(\sum_{j=k+2}^{\infty} 2^{j\alpha_2(\infty)} 2^{e_2(k-j)} \|f_2^j\|_{q_2(\cdot)} \right)^{p_2(1+\epsilon)} \right)^{\frac{1}{p_2(1+\epsilon)}} \\ &= E_{33}^1 E_{33}^2, \end{aligned}$$

we have $E_{33}^1 = E_{23}^1$ and $E_{33}^2 = E_{32}^2$, thus E_{33} is estimated.

For E_5 , by Minkowski's inequality, write

$$\begin{aligned} &\sup_{\epsilon > 0} \sup_{k_0 \in \mathbb{Z}} 2^{-k_0 \lambda} \left(\epsilon^\theta \sum_{k=-\infty}^{k_0} \left(\sum_{l=k-1}^{k+1} \sum_{j=k-1}^{k+1} \left\| 2^{k\alpha(\cdot)} B\tilde{T}_{\beta, \theta}(f_1^l, f_2^j) \chi_k \right\|_{q(\cdot)} \right)^{p(1+\epsilon)} \right)^{\frac{1}{p(1+\epsilon)}} \\ &\leq \sup_{\epsilon > 0} \sup_{k_0 \in \mathbb{Z}} 2^{-k_0 \lambda} \left(\epsilon^\theta \sum_{k=-\infty}^{-1} 2^{k\alpha(0)} \left(\sum_{l=k-1}^{k+1} \sum_{j=k-1}^{k+1} \left\| B\tilde{T}_{\beta, \theta}(f_1^l, f_2^j) \chi_k \right\|_{q(\cdot)} \right)^{p(1+\epsilon)} \right)^{\frac{1}{p(1+\epsilon)}} \\ &\quad + \sup_{\epsilon > 0} \sup_{k_0 \in \mathbb{Z}} 2^{-k_0 \lambda} \left(\epsilon^\theta \sum_{k=0}^{k_0} 2^{k\alpha(\infty)} \left(\sum_{l=k-1}^{k+1} \sum_{j=k-1}^{k+1} \left\| B\tilde{T}_{\beta, \theta}(f_1^l, f_2^j) \chi_k \right\|_{q(\cdot)} \right)^{p(1+\epsilon)} \right)^{\frac{1}{p(1+\epsilon)}} \end{aligned}$$

$$=E_{51} + E_{52}.$$

Applying Lemma 3.1, we obtain the following result

$$\begin{aligned} E_{51} &\leq C \sup_{\epsilon > 0} \sup_{k_0 \in \mathbb{Z}} 2^{-k_0 \lambda_1} \left(\epsilon^\theta \sum_{k=-\infty}^{-1} \left(\sum_{l=k-1}^{k+1} 2^{k\alpha_1(0)} \|f_1^l\|_{q_1(\cdot)} \right)^{p_1(1+\epsilon)} \right)^{\frac{1}{p_1(1+\epsilon)}} \\ &\quad \times \sup_{\epsilon > 0} \sup_{k_0 \in \mathbb{Z}} 2^{-k_0 \lambda_2} \left(\epsilon^\theta \sum_{k=-\infty}^{-1} \left(\sum_{j=k-1}^{k+1} 2^{k\alpha_2(0)} \|f_2^j\|_{q_2(\cdot)} \right)^{p_2(1+\epsilon)} \right)^{\frac{1}{p_2(1+\epsilon)}} \\ &= C \|f_1\|_{M\dot{K}_{p_1, \theta, q_1(\cdot)}^{\alpha_1(\cdot), \lambda_1}(\mathbb{R}^n)} \|f_2\|_{M\dot{K}_{p_2, \theta, q_2(\cdot)}^{\alpha_2(\cdot), \lambda_2}(\mathbb{R}^n)}, \end{aligned}$$

as with the estimation of E_{51} , the estimation of E_{52} follows a similar procedure and is therefore easy to obtain.

For E_6 , we set $k-1 \leq l \leq k+1$, $j \geq k+2$, $x \in D_k$, $y_1 \in D_l$ and $y_2 \in D_j$, such that $|x - y_1| \geq 0$ and $|x - y_2| \geq C2^j$. For $x \in D_k$, we have

$$\begin{aligned} |B\tilde{T}_{\beta, \theta}(f_1^l, f_2^j)(x)| &\leq C 2^{-2nj(1-\beta)} \int_{\mathbb{R}^n} \int_{\mathbb{R}^n} |f_1^l(y_1)| |f_2^j(y_2)| dy_1 dy_2 \\ &\leq C 2^{-nk(1-\beta)} 2^{-nj(1-\beta)} \|f_1^l\|_{L^1} \|f_2^j\|_{L^1}, \end{aligned} \quad (3.4)$$

from this and the Minkowski's inequality, it then follows that

$$\begin{aligned} &\sup_{\epsilon > 0} \sup_{k_0 \in \mathbb{Z}} 2^{-k_0 \lambda} \left(\epsilon^\theta \sum_{k=-\infty}^{k_0} \left(\sum_{l=k-1}^{k+1} \sum_{j=k+2}^{\infty} \left\| 2^{k\alpha(\cdot)} B\tilde{T}_{\beta, \theta}(f_1^l, f_2^j) \chi_k \right\|_{q(\cdot)} \right)^{p(1+\epsilon)} \right)^{\frac{1}{p(1+\epsilon)}} \\ &\leq \sup_{\epsilon > 0} \sup_{k_0 \in \mathbb{Z}} 2^{-k_0 \lambda} \left(\epsilon^\theta \sum_{k=-\infty}^{-1} 2^{k\alpha(0)} \left(\sum_{l=k-1}^{k+1} \sum_{j=k+2}^{\infty} \left\| B\tilde{T}_{\beta, \theta}(f_1^l, f_2^j) \chi_k \right\|_{q(\cdot)} \right)^{p(1+\epsilon)} \right)^{\frac{1}{p(1+\epsilon)}} \\ &\quad + \sup_{\epsilon > 0} \sup_{k_0 \in \mathbb{Z}} 2^{-k_0 \lambda} \left(\epsilon^\theta \sum_{k=0}^{k_0} 2^{k\alpha(\infty)} \left(\sum_{l=k-1}^{k+1} \sum_{j=k+2}^{\infty} \left\| B\tilde{T}_{\beta, \theta}(f_1^l, f_2^j) \chi_k \right\|_{q(\cdot)} \right)^{p(1+\epsilon)} \right)^{\frac{1}{p(1+\epsilon)}} \\ &= E_{61} + E_{62}. \end{aligned}$$

For E_{61} , we conclude the following inequality holds

$$\begin{aligned} E_{61} &\leq C \sup_{\epsilon > 0} \sup_{k_0 \in \mathbb{Z}} 2^{-k_0 \lambda_1} \left(\epsilon^\theta \sum_{k=-\infty}^{-1} \left(\sum_{l=k-1}^{k+1} 2^{k\alpha_1(0)} \|f_1^l\|_{q_1(\cdot)} \right)^{p_1(1+\epsilon)} \right)^{\frac{1}{p_1(1+\epsilon)}} \\ &\quad \times \sup_{\epsilon > 0} \sup_{k_0 \in \mathbb{Z}} 2^{-k_0 \lambda_2} \left(\epsilon^\theta \sum_{k=-\infty}^{-1} \left(\sum_{j=k+2}^{\infty} 2^{k\alpha_2(0)} \right. \right. \\ &\quad \times \left. \left. 2^{(k-j)\left(\frac{n}{q_2(\infty)} - n\beta\right)} \|f_2^j\|_{q_2(\cdot)} \right)^{p_2(1+\epsilon)} \right)^{\frac{1}{p_2(1+\epsilon)}} \\ &= E_{61}^1 E_{61}^2. \end{aligned}$$

E_{61}^1 is similar to E_{21}^2 , and since $E_{61}^2 = E_{31}^2$, it is easy to get

$$E_{61} \leq C \|f_1\|_{M\dot{K}_{p_1, \theta, q_1(\cdot)}^{\alpha_1(\cdot), \lambda_1}(\mathbb{R}^n)} \|f_2\|_{M\dot{K}_{p_2, \theta, q_2(\cdot)}^{\alpha_2(\cdot), \lambda_2}(\mathbb{R}^n)}.$$

For E_{62} , by (3.4) and Lemma 2.4, we consider

$$\begin{aligned} E_{62} &\leq C \sup_{\epsilon > 0} \sup_{k_0 \in \mathbb{Z}} 2^{-k_0 \lambda_1} \left(\epsilon^\theta \sum_{k=0}^{k_0} \left(\sum_{l=k-1}^{k+1} 2^{k\alpha_1(\infty)} \|f_1^l\|_{q_1(\cdot)} \right)^{p_1(1+\epsilon)} \right)^{\frac{1}{p_1(1+\epsilon)}} \\ &\quad \times \sup_{\epsilon > 0} \sup_{k_0 \in \mathbb{Z}} 2^{-k_0 \lambda_2} \left(\epsilon^\theta \sum_{k=0}^{k_0} \left(\sum_{j=k+2}^{\infty} 2^{j\alpha_2(\infty)} 2^{e_2(k-j)} \|f_2^j\|_{q_2(\cdot)} \right)^{p_2(1+\epsilon)} \right)^{\frac{1}{p_2(1+\epsilon)}} \\ &= E_{62}^1 E_{62}^2. \end{aligned}$$

Since the estimates of E_{62}^1 and E_{62}^2 are similar E_{22}^2 and E_{31}^2 , respectively, it is easy to see that

$$E_{62} \leq C \|f_1\|_{M\dot{K}_{p_1, \theta, q_1(\cdot)}^{\alpha_1(\cdot), \lambda_1}(\mathbb{R}^n)} \|f_2\|_{M\dot{K}_{p_2, \theta, q_2(\cdot)}^{\alpha_2(\cdot), \lambda_2}(\mathbb{R}^n)}.$$

For E_9 , we set $l, j \geq k+2$, $x \in D_k$, $y_1 \in D_l$ and $y_2 \in D_j$, such that $|x - y_1| \geq C2^l$ and $|x - y_2| \geq C2^j$. For $x \in D_k$, we have

$$\begin{aligned} \left| B\tilde{T}_{\beta, \theta}(f_1^l, f_2^j)(x) \right| &\leq C \int_{\mathbb{R}^n} \int_{\mathbb{R}^n} \frac{|f_1^l(y_1)| |f_2^j(y_2)|}{\left(2^{l(1-\beta)} + 2^{j(1-\beta)} \right)^{2n}} dy_1 dy_2 \\ &\leq C 2^{-nl(1-\beta)} \|f_1^l\|_{L^1} 2^{-nj(1-\beta)} \|f_2^j\|_{L^1}, \end{aligned} \quad (3.5)$$

by Minkowski's inequality, we have

$$\begin{aligned} &\sup_{\epsilon > 0} \sup_{k_0 \in \mathbb{Z}} 2^{-k_0 \lambda} \left(\epsilon^\theta \sum_{k=-\infty}^{k_0} \left(\sum_{l=k+2}^{\infty} \sum_{j=k+2}^{\infty} \left\| 2^{k\alpha(\cdot)} B\tilde{T}_{\beta, \theta}(f_1^l, f_2^j) \chi_k \right\|_{q(\cdot)} \right)^{p(1+\epsilon)} \right)^{\frac{1}{p(1+\epsilon)}} \\ &\leq \sup_{\epsilon > 0} \sup_{k_0 \in \mathbb{Z}} 2^{-k_0 \lambda} \left(\epsilon^\theta \sum_{k=-\infty}^{-1} 2^{k\alpha(0)} \left(\sum_{l=k+2}^{\infty} \sum_{j=k+2}^{\infty} \left\| B\tilde{T}_{\beta, \theta}(f_1^l, f_2^j) \chi_k \right\|_{q(\cdot)} \right)^{p(1+\epsilon)} \right)^{\frac{1}{p(1+\epsilon)}} \\ &\quad + \sup_{\epsilon > 0} \sup_{k_0 \in \mathbb{Z}} 2^{-k_0 \lambda} \left(\epsilon^\theta \sum_{k=0}^{k_0} 2^{k\alpha(\infty)} \left(\sum_{l=k+2}^{\infty} \sum_{j=k+2}^{\infty} \left\| B\tilde{T}_{\beta, \theta}(f_1^l, f_2^j) \chi_k \right\|_{q(\cdot)} \right)^{p(1+\epsilon)} \right)^{\frac{1}{p(1+\epsilon)}} \\ &= E_{91} + E_{92}. \end{aligned}$$

For E_{91} , by (3.5) and Lemma 3.3, we obtain the following result

$$\begin{aligned} E_{91} &\leq C \sup_{\epsilon > 0} \sup_{k_0 \in \mathbb{Z}} 2^{-k_0 \lambda_1} \left(\epsilon^\theta \sum_{k=-\infty}^{-1} \left(\sum_{l=k+2}^{\infty} 2^{k\alpha_1(0)} 2^{(k-l)\left(\frac{n}{q_1(\infty)} - n\beta\right)} \right. \right. \\ &\quad \left. \left. \times \|f_1^l\|_{q_1(\cdot)} \right)^{p_1(1+\epsilon)} \right)^{\frac{1}{p_1(1+\epsilon)}} \\ &\quad \times \sup_{\epsilon > 0} \sup_{k_0 \in \mathbb{Z}} 2^{-k_0 \lambda_2} \left(\epsilon^\theta \sum_{k=-\infty}^{-1} \left(\sum_{j=k+2}^{\infty} 2^{k\alpha_2(0)} 2^{(k-j)\left(\frac{n}{q_2(\infty)} - n\beta\right)} \right. \right. \\ &\quad \left. \left. \times \|f_2^j\|_{q_2(\cdot)} \right)^{p_2(1+\epsilon)} \right)^{\frac{1}{p_2(1+\epsilon)}} \end{aligned}$$

$$= E_{91}^1 E_{91}^2.$$

It is easy to find that E_{91}^1 and E_{91}^2 are similar to E_{32}^2 , hence, here we omit the details of the proof.

For E_{92} , it is easy to get

$$\begin{aligned} & \sup_{\epsilon > 0} \sup_{k_0 \in \mathbb{Z}} 2^{-k_0 \lambda} \left(\epsilon^\theta \sum_{k=0}^{k_0} 2^{k\alpha(\infty)} \left(\sum_{l=k+2}^{\infty} \sum_{j=k+2}^{\infty} \|B\tilde{T}_{\beta, \theta}(f_1^l, f_2^j) \chi_k\|_{q(\cdot)} \right)^{p(1+\epsilon)} \right)^{\frac{1}{p(1+\epsilon)}} \\ & \leq C \sup_{\epsilon > 0} \sup_{k_0 \in \mathbb{Z}} 2^{-k_0 \lambda_1} \left(\epsilon^\theta \sum_{k=0}^{k_0} \left(\sum_{l=k+2}^{\infty} 2^{l\alpha_1(\infty)} 2^{e_1(k-l)} \|f_1^l\|_{q_1(\cdot)} \right)^{p_1(1+\epsilon)} \right)^{\frac{1}{p_1(1+\epsilon)}} \\ & \quad \times \sup_{\epsilon > 0} \sup_{k_0 \in \mathbb{Z}} 2^{-k_0 \lambda_2} \left(\epsilon^\theta \sum_{k=0}^{k_0} \left(\sum_{j=k+2}^{\infty} 2^{j\alpha_2(\infty)} 2^{e_2(k-j)} \|f_2^j\|_{q_2(\cdot)} \right)^{p_2(1+\epsilon)} \right)^{\frac{1}{p_2(1+\epsilon)}} \\ & = E_{92}^1 E_{92}^2. \end{aligned}$$

Since the estimates for E_{92}^1 and E_{92}^2 are similar to E_{32}^2 . So, it is not difficult to obtain that

$$E_{92} \leq C \|f_1\|_{M\dot{K}_{p_1, \theta, q_1(\cdot)}^{\alpha_1(\cdot), \lambda_1}(\mathbb{R}^n)} \|f_2\|_{M\dot{K}_{p_2, \theta, q_2(\cdot)}^{\alpha_2(\cdot), \lambda_2}(\mathbb{R}^n)}.$$

Which, combining the estimates for $E_1, E_2, \dots, E_9$, yields the desired result. Hence, the proof of Theorem 3.1 is finished. $\square$

Proof of Theorem 3.2. Let $\|b_1\|_* = \|b_2\|_* = 1$, $f_i \in M\dot{K}_{p_i, \theta, q_i(\cdot)}^{\alpha_i(\cdot), \lambda_i}(\mathbb{R}^n)$, and decompose

$$f_1(x) = \sum_{l=-\infty}^{\infty} f(x) \chi_l(x) = \sum_{l=-\infty}^{\infty} f_1^l(x), \quad f_2(x) = \sum_{j=-\infty}^{\infty} f(x) \chi_j(x) = \sum_{l=-\infty}^{\infty} f_2^j(x).$$

Since the decomposition method for $\|B\tilde{T}_{\beta, \theta, b_1, b_2}(f_1, f_2)\|_{M\dot{K}_{p, \theta, q(\cdot)}^{\alpha(\cdot), \lambda}}$ is the same as that for $\|B\tilde{T}_{\beta, \theta}(f_1, f_2)\|_{M\dot{K}_{p, \theta, q(\cdot)}^{\alpha(\cdot), \lambda}}$, it is omitted here. For the sake of brevity, we set $B\tilde{T}_{\beta, \theta, b_1, b_2} = B\tilde{T}_{\beta, \theta, \tilde{b}}$ in the proof. It is necessary to estimate F_1, F_2, F_3, F_5, F_6 and F_9 , since F_4, F_7 and F_8 can be obtained by following a similar methodology to that used for F_2, F_3 and F_6 , respectively. For F_1 , we set $l, j \leq k-2$, $x \in D_k$, $y_1 \in D_l$ and $y_2 \in D_j$, such that $|x - y_i| \geq C2^k$. For $x \in D_k$, we can deduce that

$$\begin{aligned} & \left| B\tilde{T}_{\beta, \theta, \tilde{b}}(f_1^l, f_2^j)(x) \right| \\ & \leq C 2^{-2nk(1-\beta)} \int_{\mathbb{R}^n} |b_1(x) - b_1(y_1)| |f_1^l(y_1)| dy_1 \int_{\mathbb{R}^n} |b_2(x) - b_2(y_2)| |f_2^j(y_2)| dy_2 \\ & \leq C 2^{-2nk(1-\beta)} \left(|b_1(x) - b_{B_l}| \int_{\mathbb{R}^n} |f_1^l(y_1)| dy_1 + \int_{\mathbb{R}^n} |b_{B_l} - b_1(y_1)| |f_1^l(y_1)| dy_1 \right) \\ & \quad \times \left(|b_2(x) - b_{B_j}| \int_{\mathbb{R}^n} |f_2^j(y_2)| dy_2 + \int_{\mathbb{R}^n} |b_{B_j} - b_2(y_2)| |f_2^j(y_2)| dy_2 \right) \\ & \leq C 2^{-2nk(1-\beta)} \|f_1^l\|_{q_1(\cdot)} \left(|b_1(\cdot) - b_{B_l}| \|\chi_l\|_{q_1'(\cdot)} + \|b_{B_l} - b_1\|_{q_1'(\cdot)} \right) \end{aligned}$$

$$\times \|f_2^j\|_{q_2(\cdot)} \left(\|b_2(\cdot) - b_{B_j}\| \| \chi_j \|_{q_2'(\cdot)} + \|b_{B_j} - b_2\| \chi_j \|_{q_2'(\cdot)} \right). \quad (3.6)$$

From Minkowski's inequality, it follows that

$$\begin{aligned} & \sup_{\epsilon > 0} \sup_{k_0 \in \mathbb{Z}} 2^{-k_0 \lambda} \left(\epsilon^\theta \sum_{k=-\infty}^{k_0} \left(\sum_{l=-\infty}^{k-2} \sum_{j=-\infty}^{k-2} \left\| 2^{k\alpha(\cdot)} B\tilde{T}_{\beta, \theta, \tilde{b}}(f_1^l, f_2^j) \chi_k \right\|_{q(\cdot)} \right)^{p(1+\epsilon)} \right)^{\frac{1}{p(1+\epsilon)}} \\ & \leq \sup_{\epsilon > 0} \sup_{k_0 \in \mathbb{Z}} 2^{-k_0 \lambda} \left(\epsilon^\theta \sum_{k=-\infty}^{-1} \left(\sum_{l=-\infty}^{k-2} \sum_{j=-\infty}^{k-2} \left\| 2^{k\alpha(\cdot)} B\tilde{T}_{\beta, \theta, \tilde{b}}(f_1^l, f_2^j) \chi_k \right\|_{q(\cdot)} \right)^{p(1+\epsilon)} \right)^{\frac{1}{p(1+\epsilon)}} \\ & \quad + \sup_{\epsilon > 0} \sup_{k_0 \in \mathbb{Z}} 2^{-k_0 \lambda} \left(\epsilon^\theta \sum_{k=0}^{k_0} \left(\sum_{l=-\infty}^{-1} \sum_{j=-\infty}^{-1} \left\| 2^{k\alpha(\cdot)} B\tilde{T}_{\beta, \theta, \tilde{b}}(f_1^l, f_2^j) \chi_k \right\|_{q(\cdot)} \right)^{p(1+\epsilon)} \right)^{\frac{1}{p(1+\epsilon)}} \\ & \quad + \sup_{\epsilon > 0} \sup_{k_0 \in \mathbb{Z}} 2^{-k_0 \lambda} \left(\epsilon^\theta \sum_{k=0}^{k_0} \left(\sum_{l=-\infty}^{-1} \sum_{j=0}^{k-2} \left\| 2^{k\alpha(\cdot)} B\tilde{T}_{\beta, \theta, \tilde{b}}(f_1^l, f_2^j) \chi_k \right\|_{q(\cdot)} \right)^{p(1+\epsilon)} \right)^{\frac{1}{p(1+\epsilon)}} \\ & \quad + \sup_{\epsilon > 0} \sup_{k_0 \in \mathbb{Z}} 2^{-k_0 \lambda} \left(\epsilon^\theta \sum_{k=0}^{k_0} \left(\sum_{l=0}^{k-2} \sum_{j=-\infty}^{-1} \left\| 2^{k\alpha(\cdot)} B\tilde{T}_{\beta, \theta, \tilde{b}}(f_1^l, f_2^j) \chi_k \right\|_{q(\cdot)} \right)^{p(1+\epsilon)} \right)^{\frac{1}{p(1+\epsilon)}} \\ & \quad + \sup_{\epsilon > 0} \sup_{k_0 \in \mathbb{Z}} 2^{-k_0 \lambda} \left(\epsilon^\theta \sum_{k=0}^{k_0} \left(\sum_{l=0}^{k-2} \sum_{j=0}^{k-2} \left\| 2^{k\alpha(\cdot)} B\tilde{T}_{\beta, \theta, \tilde{b}}(f_1^l, f_2^j) \chi_k \right\|_{q(\cdot)} \right)^{p(1+\epsilon)} \right)^{\frac{1}{p(1+\epsilon)}} \\ & = F_{11} + F_{12} + F_{13} + F_{14} + F_{15}. \end{aligned}$$

For F_{11} , by (3.1), (3.6), Hölder's inequality, Lemmas 2.4 and 2.5, we consider

$$\begin{aligned} & \sup_{\epsilon > 0} \sup_{k_0 \in \mathbb{Z}} 2^{-k_0 \lambda} \left(\epsilon^\theta \sum_{k=-\infty}^{-1} \left(\sum_{l=-\infty}^{k-2} \sum_{j=-\infty}^{k-2} \left\| 2^{k\alpha(\cdot)} B\tilde{T}_{\beta, \theta, \tilde{b}}(f_1^l, f_2^j) \chi_k \right\|_{q(\cdot)} \right)^{p(1+\epsilon)} \right)^{\frac{1}{p(1+\epsilon)}} \\ & \leq C \sup_{\epsilon > 0} \sup_{k_0 \in \mathbb{Z}} 2^{-k_0 \lambda_1} \left(\epsilon^\theta \sum_{k=-\infty}^{-1} \left(\sum_{l=-\infty}^{k-2} (k-l) 2^{l\alpha_1(0)+a_1(k-l)} \|f_1^l\|_{q_1(\cdot)} \right)^{p_1(1+\epsilon)} \right)^{\frac{1}{p_1(1+\epsilon)}} \\ & \quad \times \sup_{\epsilon > 0} \sup_{k_0 \in \mathbb{Z}} 2^{-k_0 \lambda_2} \left(\epsilon^\theta \sum_{k=-\infty}^{-1} \left(\sum_{j=-\infty}^{k-2} (k-j) 2^{j\alpha_2(0)+a_2(k-j)} \|f_2^j\|_{q_2(\cdot)} \right)^{p_2(1+\epsilon)} \right)^{\frac{1}{p_2(1+\epsilon)}} \\ & = F_{11}^1 \times F_{11}^2. \end{aligned}$$

For F_{11}^1 , we have

$$\begin{aligned} F_{11}^1 & \leq C \sup_{\epsilon > 0} \sup_{k_0 \in \mathbb{Z}} 2^{-k_0 \lambda_1} \left(\epsilon^\theta \sum_{k=-\infty}^{-1} \left(\sum_{l=-\infty}^{k-1} 2^{l\alpha_1(0)p_1(1+\epsilon)} 2^{\frac{a_1(k-l)p_1(1+\epsilon)}{2}} \|f_1^l\|_{q_1(\cdot)}^{p_1(1+\epsilon)} \right) \right. \\ & \quad \times \left. \left(\sum_{l=-\infty}^{k-1} (k-l) p_1'(1+\epsilon) 2^{\frac{a_1(k-l)p_1'(1+\epsilon)}{2}} \right)^{\frac{p_1(1+\epsilon)}{p_1'(1+\epsilon)}} \right)^{\frac{1}{p_1(1+\epsilon)}} \\ & \leq C \|f_1\|_{M\dot{K}_{p_1, \theta, q_1(\cdot)}^{\alpha_1(\cdot), \lambda_1}(\mathbb{R}^n)}. \end{aligned}$$

With an argument similar to that used in the estimate for F_{11}^1 , it is easy to get

$$F_{11}^2 \leq C \|f_2\|_{M\dot{K}_{p_2, \theta, q_2(\cdot)}^{\alpha_2(\cdot), \lambda_2}(\mathbb{R}^n)}.$$

For F_{12} , by (3.1), (3.6), Lemmas 3.3 and 3.4, we deduce

$$\begin{aligned} & \sup_{\epsilon > 0} \sup_{k_0 \in \mathbb{Z}} 2^{-k_0 \lambda} \left(\epsilon^\theta \sum_{k=0}^{k_0} \left(\sum_{l=-\infty}^{-1} \sum_{j=-\infty}^{-1} \left\| 2^{k\alpha(\cdot)} B\tilde{T}_{\beta, \theta, \tilde{b}}(f_1^l, f_2^j) \chi_k \right\|_{q(\cdot)} \right)^{p(1+\epsilon)} \right)^{\frac{1}{p(1+\epsilon)}} \\ & \leq C \sup_{\epsilon > 0} \sup_{k_0 \in \mathbb{Z}} 2^{-k_0 \lambda_1} \left(\epsilon^\theta \sum_{k=0}^{k_0} \left(\sum_{l=-\infty}^{-1} (k-l) 2^{l\alpha_1(0)+b_1(k-l)} \|f_1^l\|_{q_1(\cdot)} \right)^{p_1(1+\epsilon)} \right)^{\frac{1}{p_1(1+\epsilon)}} \\ & \quad \times \sup_{\epsilon > 0} \sup_{k_0 \in \mathbb{Z}} 2^{-k_0 \lambda_2} \left(\epsilon^\theta \sum_{k=0}^{\infty} \left(\sum_{j=-\infty}^{-1} (k-j) 2^{j\alpha_2(0)+b_2(k-j)} \|f_2^j\|_{q_2(\cdot)} \right)^{p_2(1+\epsilon)} \right)^{\frac{1}{p_2(1+\epsilon)}} \\ & = F_{12}^1 \times F_{12}^2. \end{aligned}$$

The estimates for F_{12}^1 , F_{12}^2 and F_{11}^1 are similar, hence, it is easy to get

$$F_{12} \leq C \|f_2\|_{M\dot{K}_{p_1, \theta, q_1(\cdot)}^{\alpha_1(\cdot), \lambda_1}(\mathbb{R}^n)} \|f_2\|_{M\dot{K}_{p_2, \theta, q_2(\cdot)}^{\alpha_2(\cdot), \lambda_2}(\mathbb{R}^n)}.$$

For F_{13} , by $q_2(0) \leq q_2(\infty)$, we have

$$\begin{aligned} & \sum_{l=-\infty}^{-1} \sum_{j=0}^{k-2} \|B\tilde{T}_{\beta, \theta, \tilde{b}}(f_1^l, f_2^j) \chi_k\|_{q(\cdot)} \\ & \leq C \sum_{l=-\infty}^{-1} (k-l) 2^{(k-l)(\frac{n}{q_1(0)}-n)} \|f_1^l\|_{q_1(\cdot)} \sum_{j=0}^{k-2} (k-j) 2^{(k-j)(\frac{n}{q_2(\infty)}-n)} \|f_2^j\|_{q_2(\cdot)}. \end{aligned}$$

With a way similar to that used in the estimate for F_{12} , it is not difficult to get

$$F_{13} \leq C \|f_1\|_{M\dot{K}_{p_1, \theta, q_1(\cdot)}^{\alpha_1(\cdot), \lambda_1}(\mathbb{R}^n)} \|f_2\|_{M\dot{K}_{p_2, \theta, q_2(\cdot)}^{\alpha_2(\cdot), \lambda_2}(\mathbb{R}^n)}.$$

From the symmetry, by F_{13} we get

$$F_{14} \leq C \|f_1\|_{M\dot{K}_{p_1, \theta, q_1(\cdot)}^{\alpha_1(\cdot), \lambda_1}(\mathbb{R}^n)} \|f_2\|_{M\dot{K}_{p_2, \theta, q_2(\cdot)}^{\alpha_2(\cdot), \lambda_2}(\mathbb{R}^n)}.$$

For F_{15} , from (3.6), Lemmas 3.3 and 3.4 and $\frac{n}{q_i(\infty)} - n + \alpha_i(\infty) < 0$, just like F_{11} , we obtain

$$F_{15} \leq C \|f_1\|_{M\dot{K}_{p_1, \theta, q_1(\cdot)}^{\alpha_1(\cdot), \lambda_1}(\mathbb{R}^n)} \|f_2\|_{M\dot{K}_{p_2, \theta, q_2(\cdot)}^{\alpha_2(\cdot), \lambda_2}(\mathbb{R}^n)}.$$

For F_2 , we set $l \leq k-2$, $k-1 \leq j \leq k+1$, $x \in D_k$, $y_1 \in D_l$ and $y_2 \in D_j$, such that $|x - y_1| \geq C2^k$ and $|x - y_1| \geq 0$. For $x \in D_k$, we deduce

$$\begin{aligned} & \left| B\tilde{T}_{\beta, \theta, \tilde{b}}(f_1^l, f_2^j)(x) \right| \\ & \leq C 2^{-2nk(1-\beta)} \|f_1^l\|_{q_1(\cdot)} \left(|b_1(\cdot) - b_{B_l}| \|\chi_l\|_{q'_1(\cdot)} + \|b_{B_l} - b_1\| \|\chi_l\|_{q'_1(\cdot)} \right) \end{aligned}$$

$$\times \|f_2^j\|_{q_2(\cdot)} \left(\|b_2(\cdot) - b_{B_j}\| \|\chi_j\|_{q'_2(\cdot)} + \|b_{B_j} - b_2\| \|\chi_j\|_{q'_2(\cdot)} \right), \quad (3.7)$$

by this and Minkowski's inequality, write

$$\begin{aligned} & \sup_{\epsilon > 0} \sup_{k_0 \in \mathbb{Z}} 2^{-k_0 \lambda} \left(\epsilon^\theta \sum_{k=-\infty}^{k_0} \left(\sum_{l=-\infty}^{k-2} \sum_{j=k-1}^{k+1} \left\| 2^{k\alpha(\cdot)} B\tilde{T}_{\beta, \theta, \tilde{b}}(f_1^l, f_2^j) \chi_k \right\|_{q(\cdot)} \right)^{p(1+\epsilon)} \right)^{\frac{1}{p(1+\epsilon)}} \\ & \leq \sup_{\epsilon > 0} \sup_{k_0 \in \mathbb{Z}} 2^{-k_0 \lambda} \left(\epsilon^\theta \sum_{k=-\infty}^{-1} \left(\sum_{l=-\infty}^{k-2} \sum_{j=k-1}^{k+1} \left\| 2^{k\alpha(\cdot)} B\tilde{T}_{\beta, \theta, \tilde{b}}(f_1^l, f_2^j) \chi_k \right\|_{q(\cdot)} \right)^{p(1+\epsilon)} \right)^{\frac{1}{p(1+\epsilon)}} \\ & \quad + \sup_{\epsilon > 0} \sup_{k_0 \in \mathbb{Z}} 2^{-k_0 \lambda} \left(\epsilon^\theta \sum_{k=0}^{k_0} \left(\sum_{l=-\infty}^{-1} \sum_{j=k-1}^{k+1} \left\| 2^{k\alpha(\cdot)} B\tilde{T}_{\beta, \theta, \tilde{b}}(f_1^l, f_2^j) \chi_k \right\|_{q(\cdot)} \right)^{p(1+\epsilon)} \right)^{\frac{1}{p(1+\epsilon)}} \\ & \quad + \sup_{\epsilon > 0} \sup_{k_0 \in \mathbb{Z}} 2^{-k_0 \lambda} \left(\epsilon^\theta \sum_{k=0}^{k_0} \left(\sum_{l=0}^{k-2} \sum_{j=k-1}^{k+1} \left\| 2^{k\alpha(\cdot)} B\tilde{T}_{\beta, \theta, \tilde{b}}(f_1^l, f_2^j) \chi_k \right\|_{q(\cdot)} \right)^{p(1+\epsilon)} \right)^{\frac{1}{p(1+\epsilon)}} \\ & = F_{21} + F_{22} + F_{23}. \end{aligned}$$

For F_{21} , by (3.2), (3.7), Lemmas 3.3 and 3.4, we conclude

$$\begin{aligned} & \sup_{\epsilon > 0} \sup_{k_0 \in \mathbb{Z}} 2^{-k_0 \lambda} \left(\epsilon^\theta \sum_{k=-\infty}^{-1} \left(\sum_{l=-\infty}^{k-2} \sum_{j=k-1}^{k+1} \left\| 2^{k\alpha(\cdot)} B\tilde{T}_{\beta, \theta, \tilde{b}}(f_1^l, f_2^j) \chi_k \right\|_{q(\cdot)} \right)^{p(1+\epsilon)} \right)^{\frac{1}{p(1+\epsilon)}} \\ & \leq C \sup_{\epsilon > 0} \sup_{k_0 \in \mathbb{Z}} 2^{-k_0 \lambda_1} \left(\epsilon^\theta \sum_{k=-\infty}^{-1} \left(\sum_{l=-\infty}^{-1} (k-l) 2^{l\alpha_1(0)+a_1(k-l)} \|f_1^l\|_{q_1(\cdot)} \right)^{p_1(1+\epsilon)} \right)^{\frac{1}{p_1(1+\epsilon)}} \\ & \quad \times \sup_{\epsilon > 0} \sup_{k_0 \in \mathbb{Z}} 2^{-k_0 \lambda_2} \left(\epsilon^\theta \sum_{k=-\infty}^{-1} \left(\sum_{j=k-1}^{k+1} 2^{k\alpha_2(0)} \|f_2^j\|_{q_2(\cdot)} \right)^{p_2(1+\epsilon)} \right)^{\frac{1}{p_2(1+\epsilon)}} \\ & = F_{21}^1 \times F_{21}^2. \end{aligned}$$

Since F_{21}^1 and F_{11}^1 are the same. Thus, we only turn F_{21}^2 ,

$$\begin{aligned} F_{21}^2 & \leq C \sup_{\epsilon > 0} \sup_{k_0 \in \mathbb{Z}} 2^{-k_0 \lambda_2} \left(\epsilon^\theta \sum_{k=-\infty}^{-1} 2^{k\alpha_2(0)p_2(1+\epsilon)} \|f_2^k\|_{q_2(\cdot)}^{p_2(1+\epsilon)} \right)^{\frac{1}{p_2(1+\epsilon)}} \\ & \leq C \|f_2\|_{MK_{p_2, \theta, q_2(\cdot)}^{\alpha_2(\cdot), \lambda_2}(\mathbb{R}^n)}. \end{aligned}$$

For F_{22} , by (3.2), (3.7), Lemmas 3.3 and 3.4, we get

$$\begin{aligned} F_{22} & \leq C \sup_{\epsilon > 0} \sup_{k_0 \in \mathbb{Z}} 2^{-k_0 \lambda_1} \left(\epsilon^\theta \sum_{k=0}^{k_0} \left(\sum_{l=-\infty}^{-1} (k-l) 2^{l\alpha_1(0)+b_1(k-l)} \|f_1^l\|_{q_1(\cdot)} \right)^{p_1(1+\epsilon)} \right)^{\frac{1}{p_1(1+\epsilon)}} \\ & \quad \times \sup_{\epsilon > 0} \sup_{k_0 \in \mathbb{Z}} 2^{-k_0 \lambda_2} \left(\epsilon^\theta \sum_{k=0}^{k_0} \left(\sum_{j=k-1}^{k+1} 2^{k\alpha_2(\infty)} \|f_2^j\|_{q_2(\cdot)} \right)^{p_2(1+\epsilon)} \right)^{\frac{1}{p_2(1+\epsilon)}} \\ & = F_{22}^1 \times F_{22}^2. \end{aligned}$$

Since F_{22}^2 and F_{22}^1 are similar, and $F_{22}^1 = F_{12}^1$, it is easy to get

$$F_{22} \leq C \|f_1\|_{MK_{p_1, \theta, q_1(\cdot)}^{\alpha_1(\cdot), \lambda_1}(\mathbb{R}^n)} \|f_2\|_{MK_{p_2, \theta, q_2(\cdot)}^{\alpha_2(\cdot), \lambda_2}(\mathbb{R}^n)}.$$

For F_{23} , by (3.2), (3.7), Lemmas 3.3 and 3.4, we obtain the following inequality:

$$\begin{aligned} F_{23} &\leq C \sup_{\epsilon > 0} \sup_{k_0 \in \mathbb{Z}} 2^{-k_0 \lambda_1} \left(\epsilon^\theta \sum_{k=0}^{k_0} \left(\sum_{l=0}^{k-2} (k-l) 2^{l\alpha_1(\infty)} 2^{c_1(k-l)} \|f_1^l\|_{q_1(\cdot)} \right)^{p_1(1+\epsilon)} \right)^{\frac{1}{p_1(1+\epsilon)}} \\ &\quad \times \sup_{\epsilon > 0} \sup_{k_0 \in \mathbb{Z}} 2^{-k_0 \lambda_2} \left(\epsilon^\theta \sum_{k=0}^{k_0} \left(\sum_{j=k-1}^{k+1} 2^{k\alpha_2(\infty)} 2^{\|f_2^j\|_{q_2(\cdot)}} \right)^{p_2(1+\epsilon)} \right)^{\frac{1}{p_2(1+\epsilon)}} \\ &= F_{23}^1 \times F_{23}^2, \end{aligned}$$

we can find $F_{23}^1 = F_{15}^1$ and $F_{23}^2 = F_{22}^2$. Therefore, we obtain an estimate of F_{23} .

For F_3 , we set $l \leq k-2$, $j \geq k+2$, $x \in D_k$, $y_1 \in D_l$ and $y_2 \in D_j$, such that $|x - y_1| \geq C2^k$ and $|x - y_1| \geq C2^j$. For $x \in D_k$, we obtain

$$\begin{aligned} &\left| BT_{\beta, \theta, \tilde{b}}(f_l, f_j)(x) \right| \\ &\leq C 2^{-nk(1-\beta)} \|f_1^l\|_{q_1(\cdot)} \left(\|b_1(\cdot) - b_{B_l}\| \|\chi_l\|_{q'_1(\cdot)} + \|b_{B_l} - b_1\| \|\chi_l\|_{q'_1(\cdot)} \right) \\ &\quad \times 2^{-nj(1-\beta)} \|f_2^j\|_{q_2(\cdot)} \left(\|b_2(\cdot) - b_{B_k}\| \|\chi_j\|_{q'_2(\cdot)} + \|b_{B_k} - b_2\| \|\chi_j\|_{q'_2(\cdot)} \right), \quad (3.8) \end{aligned}$$

write

$$\begin{aligned} &\sup_{\epsilon > 0} \sup_{k_0 \in \mathbb{Z}} 2^{-k_0 \lambda} \left(\epsilon^\theta \sum_{k=-\infty}^{k_0} \left(\sum_{l=-\infty}^{k-2} \sum_{j=k+2}^{\infty} \|2^{k\alpha(\cdot)} BT_{\beta, \theta, \tilde{b}}(f_1^l, f_2^j) \chi_k\|_{q(\cdot)} \right)^{p(1+\epsilon)} \right)^{\frac{1}{p(1+\epsilon)}} \\ &\leq \sup_{\epsilon > 0} \sup_{k_0 \in \mathbb{Z}} 2^{-k_0 \lambda} \left(\epsilon^\theta \sum_{k=-\infty}^{-1} \left(\sum_{l=-\infty}^{k-2} \sum_{j=k+2}^{\infty} \|2^{k\alpha(\cdot)} BT_{\beta, \theta, \tilde{b}}(f_1^l, f_2^j) \chi_k\|_{q(\cdot)} \right)^{p(1+\epsilon)} \right)^{\frac{1}{p(1+\epsilon)}} \\ &\quad + \sup_{\epsilon > 0} \sup_{k_0 \in \mathbb{Z}} 2^{-k_0 \lambda} \left(\epsilon^\theta \sum_{k=0}^{k_0} \left(\sum_{l=-\infty}^{-1} \sum_{j=k+2}^{\infty} \|2^{k\alpha(\cdot)} BT_{\beta, \theta, \tilde{b}}(f_1^l, f_2^j) \chi_k\|_{q(\cdot)} \right)^{p(1+\epsilon)} \right)^{\frac{1}{p(1+\epsilon)}} \\ &\quad + \sup_{\epsilon > 0} \sup_{k_0 \in \mathbb{Z}} 2^{-k_0 \lambda} \left(\epsilon^\theta \sum_{k=0}^{k_0} \left(\sum_{l=0}^{k-2} \sum_{j=k+2}^{\infty} \|2^{k\alpha(\cdot)} BT_{\beta, \theta, \tilde{b}}(f_1^l, f_2^j) \chi_k\|_{q(\cdot)} \right)^{p(1+\epsilon)} \right)^{\frac{1}{p(1+\epsilon)}} \\ &= F_{31} + F_{32} + F_{33}. \end{aligned}$$

For F_{31} , by (3.3), (3.8), Lemmas 3.3 and 3.4, we consider

$$\begin{aligned} &\sup_{\epsilon > 0} \sup_{k_0 \in \mathbb{Z}} 2^{-k_0 \lambda} \left(\epsilon^\theta \sum_{k=-\infty}^{-1} 2^{k\alpha(0)} \left(\sum_{l=-\infty}^{k-2} \sum_{j=k+2}^{\infty} \|BT_{\beta, \theta, \tilde{b}}(f_1^l, f_2^j) \chi_k\|_{q(\cdot)} \right)^{p(1+\epsilon)} \right)^{\frac{1}{p(1+\epsilon)}} \\ &\leq C \sup_{\epsilon > 0} \sup_{k_0 \in \mathbb{Z}} 2^{-k_0 \lambda_1} \left(\epsilon^\theta \sum_{k=-\infty}^{-1} \left(\sum_{l=-\infty}^{k-2} (k-l) 2^{l\alpha_1(0) + a_1(k-l)} \|f_1^l\|_{q_1(\cdot)} \right)^{p_1(1+\epsilon)} \right)^{\frac{1}{p_1(1+\epsilon)}} \\ &\quad \times \sup_{\epsilon > 0} \sup_{k_0 \in \mathbb{Z}} 2^{-k_0 \lambda_2} \left(\epsilon^\theta \sum_{k=-\infty}^{-1} \left(\sum_{j=k+2}^{\infty} (j-k) 2^{k\alpha_2(0)} \right)^{p_2(1+\epsilon)} \right)^{\frac{1}{p_2(1+\epsilon)}} \end{aligned}$$

$$\times 2^{(k-j)\left(\frac{n}{q_2(\infty)} - n\beta\right)} \|f_2^j\|_{q_2(\cdot)} \Big)^{\frac{p_2(1+\epsilon)}{p_2(1+\epsilon)}} \Big)^{\frac{1}{p_2(1+\epsilon)}} \\ = F_{31}^1 F_{31}^2.$$

It is easy to see that $F_{31}^1 = F_{11}^1$, hence, we only estimate F_{31}^2 . Write

$$\begin{aligned} F_{31}^2 &\leq C \sup_{\epsilon > 0} \sup_{k_0 \in \mathbb{Z}} 2^{-k_0 \lambda_2} \left(\epsilon^\theta \sum_{k=-\infty}^{-1} \left(\sum_{j=k+2}^{-1} (j-k) 2^{j\alpha_2(0)} 2^{d_2(k-j)} \right. \right. \\ &\quad \times \|f_2^j\|_{q_2(\cdot)} \Big)^{\frac{p_2(1+\epsilon)}{p_2(1+\epsilon)}} \Big)^{\frac{1}{p_2(1+\epsilon)}} \\ &\quad + C \sup_{\epsilon > 0} \sup_{k_0 \in \mathbb{Z}} 2^{-k_0 \lambda_2} \left(\epsilon^\theta \sum_{k=-\infty}^{-1} \left(\sum_{j=0}^{\infty} (j-k) 2^{j\alpha_2(\infty)} 2^{d_2(k-j)} \right. \right. \\ &\quad \times \|f_2^j\|_{q_2(\cdot)} \Big)^{\frac{p_2(1+\epsilon)}{p_2(1+\epsilon)}} \Big)^{\frac{1}{p_2(1+\epsilon)}} \\ &= D_3 + D_4. \end{aligned}$$

For D_3 , by the Hölder's inequality, we have

$$\begin{aligned} D_3 &\leq C \sup_{\epsilon > 0} \sup_{k_0 \in \mathbb{Z}} 2^{-k_0 \lambda_2} \left(\epsilon^\theta \sum_{k=-\infty}^{-1} \left(\sum_{j=k+2}^{-1} 2^{j\alpha_2(0)p_2(1+\epsilon)} 2^{\frac{d_2(k-j)p_2(1+\epsilon)}{2}} \|f_2^j\|_{q_2(\cdot)}^{p_2(1+\epsilon)} \right. \right. \\ &\quad \times \left. \left(\sum_{j=k+2}^{-1} (j-k)^{p_2'(1+\epsilon)} 2^{j\alpha_2(0)p_2'(1+\epsilon)} 2^{\frac{d_2(k-j)p_2'(1+\epsilon)}{2}} \right)^{\frac{p_2(1+\epsilon)}{p_2'(1+\epsilon)}} \right)^{\frac{1}{p_2(1+\epsilon)}} \\ &\leq C \|f_2\|_{M\dot{K}_{p_2, \theta, q_2(\cdot)}^{\alpha_2(\cdot), \lambda_2}(\mathbb{R}^n)}. \end{aligned}$$

Just like the D_3 , we also have

$$D_4 \leq C \|f_2\|_{M\dot{K}_{p_2, \theta, q_2(\cdot)}^{\alpha_2(\cdot), \lambda_2}(\mathbb{R}^n)}.$$

For F_{32} , by (3.3), (3.8), Lemmas 3.3 and 3.4, write

$$\begin{aligned} &\sup_{\epsilon > 0} \sup_{k_0 \in \mathbb{Z}} 2^{-k_0 \lambda} \left(\epsilon^\theta \sum_{k=0}^{k_0} \left(\sum_{l=-\infty}^{-1} \sum_{j=k+2}^{\infty} \left\| 2^{k\alpha(\cdot)} B\tilde{T}_{\beta, \theta, \tilde{b}}(f_1^l, f_2^j) \chi_k \right\|_{q(\cdot)} \right)^{\frac{p(1+\epsilon)}{p(1+\epsilon)}} \right)^{\frac{1}{p(1+\epsilon)}} \\ &\leq C \sup_{\epsilon > 0} \sup_{k_0 \in \mathbb{Z}} 2^{-k_0 \lambda_1} \left(\epsilon^\theta \sum_{k=0}^{k_0} \left(\sum_{l=-\infty}^{-1} (k-l) 2^{l\alpha_1(0)} 2^{b_1(k-l)} \|f_1^l\|_{q_1(\cdot)} \right)^{\frac{p_1(1+\epsilon)}{p_1(1+\epsilon)}} \right)^{\frac{1}{p_1(1+\epsilon)}} \\ &\quad \times \sup_{\epsilon > 0} \sup_{k_0 \in \mathbb{Z}} 2^{-k_0 \lambda_2} \left(\epsilon^\theta \sum_{k=0}^{k_0} \left(\sum_{j=k+2}^{\infty} (j-k) 2^{j\alpha_2(\infty)+e_2(k-j)} \|f_2^j\|_{q_2(\cdot)} \right)^{\frac{p_2(1+\epsilon)}{p_2(1+\epsilon)}} \right)^{\frac{1}{p_2(1+\epsilon)}} \\ &= F_{32}^1 F_{32}^2. \end{aligned}$$

It is obvious to find that $F_{32}^1 = F_{12}^1$ and the estimate of F_{32}^2 is similar to D_3 . Hence, we have

$$F_{32} \leq C \|f_1\|_{M\dot{K}_{p_1, \theta, q_1(\cdot)}^{\alpha_1(\cdot), \lambda_1}(\mathbb{R}^n)} \|f_2\|_{M\dot{K}_{p_2, \theta, q_2(\cdot)}^{\alpha_2(\cdot), \lambda_2}(\mathbb{R}^n)}.$$

For F_{33} , just like F_{32} , we obtain

$$\begin{aligned} & \sup_{\epsilon > 0} \sup_{k_0 \in \mathbb{Z}} 2^{-k_0 \lambda} \left(\epsilon^\theta \sum_{k=0}^{k_0} \left(\sum_{l=0}^{k-2} \sum_{j=k+2}^{\infty} \|2^{k\alpha(\cdot)} B\tilde{T}_{\beta, \theta, \tilde{b}}(f_1^l, f_2^j) \chi_k\|_{q(\cdot)} \right)^{p(1+\epsilon)} \right)^{\frac{1}{p(1+\epsilon)}} \\ & \leq C \sup_{\epsilon > 0} \sup_{k_0 \in \mathbb{Z}} 2^{-k_0 \lambda_1} \left(\epsilon^\theta \sum_{k=0}^{k_0} \left(\sum_{l=0}^{k-2} (k-l) 2^{l\alpha_1(\infty)} 2^{c_1(k-l)} \|f_1^l\|_{q_1(\cdot)} \right)^{p_1(1+\epsilon)} \right)^{\frac{1}{p_1(1+\epsilon)}} \\ & \quad \times \sup_{\epsilon > 0} \sup_{k_0 \in \mathbb{Z}} 2^{-k_0 \lambda_2} \left(\epsilon^\theta \sum_{k=0}^{k_0} \left(\sum_{j=k+2}^{\infty} (j-k) 2^{j\alpha_2(\infty)+e_2(k-j)} \|f_2^j\|_{q_2(\cdot)} \right)^{p_2(1+\epsilon)} \right)^{\frac{1}{p_2(1+\epsilon)}} \\ & = F_{33}^1 F_{33}^2, \end{aligned}$$

we have $F_{33}^1 = F_{23}^1$ and $F_{33}^2 = F_{32}^2$, thus, F_{33} is estimated.

For F_5 , by Minkowski's inequality, write

$$\begin{aligned} & \sup_{\epsilon > 0} \sup_{k_0 \in \mathbb{Z}} 2^{-k_0 \lambda} \left(\epsilon^\theta \sum_{k=-\infty}^{k_0} \left(\sum_{l=k-1}^{k+1} \sum_{j=k-1}^{k+1} \|2^{k\alpha(\cdot)} B\tilde{T}_{\beta, \theta, \tilde{b}}(f_1^l, f_2^j) \chi_k\|_{q(\cdot)} \right)^{p(1+\epsilon)} \right)^{\frac{1}{p(1+\epsilon)}} \\ & \leq \sup_{\epsilon > 0} \sup_{k_0 \in \mathbb{Z}} 2^{-k_0 \lambda} \left(\epsilon^\theta \sum_{k=-\infty}^{-1} \left(\sum_{l=k-1}^{k+1} \sum_{j=k-1}^{k+1} \|2^{k\alpha(\cdot)} B\tilde{T}_{\beta, \theta, \tilde{b}}(f_1^l, f_2^j) \chi_k\|_{q(\cdot)} \right)^{p(1+\epsilon)} \right)^{\frac{1}{p(1+\epsilon)}} \\ & \quad + \sup_{\epsilon > 0} \sup_{k_0 \in \mathbb{Z}} 2^{-k_0 \lambda} \left(\epsilon^\theta \sum_{k=0}^{k_0} \left(\sum_{l=k-1}^{k+1} \sum_{j=k-1}^{k+1} \|2^{k\alpha(\cdot)} B\tilde{T}_{\beta, \theta, \tilde{b}}(f_1^l, f_2^j) \chi_k\|_{q(\cdot)} \right)^{p(1+\epsilon)} \right)^{\frac{1}{p(1+\epsilon)}} \\ & = F_{51} + F_{52}. \end{aligned}$$

By Lemmas 3.3 and 3.4, we obtain

$$\begin{aligned} F_{51} & \leq C \sup_{\epsilon > 0} \sup_{k_0 \in \mathbb{Z}} 2^{-k_0 \lambda_1} \left(\epsilon^\theta \sum_{k=-\infty}^{-1} \left(\sum_{l=k-1}^{k+1} 2^{k\alpha_1(0)} \|f_1^l\|_{q_1(\cdot)} \right)^{p_1(1+\epsilon)} \right)^{\frac{1}{p_1(1+\epsilon)}} \\ & \quad \times \sup_{\epsilon > 0} \sup_{k_0 \in \mathbb{Z}} 2^{-k_0 \lambda_2} \left(\epsilon^\theta \sum_{k=-\infty}^{-1} \left(\sum_{j=k-1}^{k+1} 2^{k\alpha_2(0)} \|f_2^j\|_{q_2(\cdot)} \right)^{p_2(1+\epsilon)} \right)^{\frac{1}{p_2(1+\epsilon)}} \\ & = C \|f_1\|_{\dot{M}_{p_1, \theta, q_1(\cdot)}^{\alpha_1(\cdot), \lambda_1}(\mathbb{R}^n)} \|f_2\|_{\dot{M}_{p_2, \theta, q_2(\cdot)}^{\alpha_2(\cdot), \lambda_2}(\mathbb{R}^n)}, \end{aligned}$$

as with the estimation of F_{51} , the estimation of F_{52} follows a similar procedure and is therefore easy to obtain.

For F_6 , we set $k-1 \leq l \leq k+1$, $j \geq k+2$, $x \in D_k$, $y_1 \in D_l$ and $y_2 \in D_j$, such that $|x - y_1| \geq 0$ and $|x - y_1| \geq C2^j$. Then, we have

$$\begin{aligned} & \left| B\tilde{T}_{\beta, \theta, b_1, b_2}(f_1^l, f_2^j)(x) \right| \\ & \leq C 2^{-nk(1-\beta)} 2^{-nj(1-\beta)} \|f_1^l\|_{q_1(\cdot)} \left(\|b_1(\cdot) - b_{B_l}\| \|\chi_l\|_{q'_1(\cdot)} + \|b_{B_l} - b_1\| \|\chi_l\|_{q'_1(\cdot)} \right) \\ & \quad \times \|f_2^j\|_{q_2(\cdot)} \left(\|b_2(\cdot) - b_{B_k}\| \|\chi_j\|_{q'_2(\cdot)} + \|b_{B_k} - b_2\| \|\chi_j\|_{q'_2(\cdot)} \right), \end{aligned} \quad (3.9)$$

from this and Minkowski's inequality, it then follows that

$$\begin{aligned}
 & \sup_{\epsilon > 0} \sup_{k_0 \in \mathbb{Z}} 2^{-k_0 \lambda} \left(\epsilon^\theta \sum_{k=-\infty}^{k_0} \left(\sum_{l=k-1}^{k+1} \sum_{j=k+2}^{\infty} \left\| 2^{k\alpha(\cdot)} B\tilde{T}_{\beta, \theta, \tilde{b}}(f_1^l, f_2^j) \chi_k \right\|_{q(\cdot)} \right)^{p(1+\epsilon)} \right)^{\frac{1}{p(1+\epsilon)}} \\
 & \leq \sup_{\epsilon > 0} \sup_{k_0 \in \mathbb{Z}} 2^{-k_0 \lambda} \left(\epsilon^\theta \sum_{k=-\infty}^{-1} \left(\sum_{l=k-1}^{k+1} \sum_{j=k+2}^{\infty} \left\| 2^{k\alpha(\cdot)} B\tilde{T}_{\beta, \theta, \tilde{b}}(f_1^l, f_2^j) \chi_k \right\|_{q(\cdot)} \right)^{p(1+\epsilon)} \right)^{\frac{1}{p(1+\epsilon)}} \\
 & \quad + \sup_{\epsilon > 0} \sup_{k_0 \in \mathbb{Z}} 2^{-k_0 \lambda} \left(\epsilon^\theta \sum_{k=0}^{k_0} \left(\sum_{l=k-1}^{k+1} \sum_{j=k+2}^{\infty} \left\| 2^{k\alpha(\cdot)} B\tilde{T}_{\beta, \theta, \tilde{b}}(f_1^l, f_2^j) \chi_k \right\|_{q(\cdot)} \right)^{p(1+\epsilon)} \right)^{\frac{1}{p(1+\epsilon)}} \\
 & = F_{61} + F_{62}.
 \end{aligned}$$

For F_{61} , by (3.4), (3.9) and Lemmas 3.3 and 3.4, we conclude that F_{61} satisfies the inequality

$$\begin{aligned}
 & \sup_{\epsilon > 0} \sup_{k_0 \in \mathbb{Z}} 2^{-k_0 \lambda} \left(\epsilon^\theta \sum_{k=-\infty}^{-1} \left(\sum_{l=k-1}^{k+1} \sum_{j=k+2}^{\infty} \left\| 2^{k\alpha(\cdot)} B\tilde{T}_{\beta, \theta, \tilde{b}}(f_1^l, f_2^j) \chi_k \right\|_{q(\cdot)} \right)^{p(1+\epsilon)} \right)^{\frac{1}{p(1+\epsilon)}} \\
 & \leq C \sup_{\epsilon > 0} \sup_{k_0 \in \mathbb{Z}} 2^{-k_0 \lambda_1} \left(\epsilon^\theta \sum_{k=-\infty}^{-1} \left(\sum_{l=k-1}^{k+1} 2^{k\alpha_1(0)} \|f_1^l\|_{q_1(\cdot)} \right)^{p_1(1+\epsilon)} \right)^{\frac{1}{p_1(1+\epsilon)}} \\
 & \quad \times \sup_{\epsilon > 0} \sup_{k_0 \in \mathbb{Z}} 2^{-k_0 \lambda_2} \left(\epsilon^\theta \sum_{k=-\infty}^{-1} \left(\sum_{j=k+2}^{\infty} (j-k) 2^{k\alpha_2(0) + (k-j)(\frac{n}{q_2(\infty)} - n\beta)} \right. \right. \\
 & \quad \left. \left. \times \|f_2^j\|_{q_2(\cdot)} \right)^{p_2(1+\epsilon)} \right)^{\frac{1}{p_2(1+\epsilon)}} \\
 & = F_{61}^1 F_{61}^2,
 \end{aligned}$$

F_{61}^1 is similar to F_{21}^2 and $F_{61}^2 = F_{31}^2$, we obtain

$$F_{61} \leq C \|f_1\|_{M\dot{K}_{p_1, \theta, q_1(\cdot)}^{\alpha_1(\cdot), \lambda_1}(\mathbb{R}^n)} \|f_2\|_{M\dot{K}_{p_2, \theta, q_2(\cdot)}^{\alpha_2(\cdot), \lambda_2}(\mathbb{R}^n)}.$$

For F_{62} , by (3.9), we consider

$$\begin{aligned}
 & \sup_{\epsilon > 0} \sup_{k_0 \in \mathbb{Z}} 2^{-k_0 \lambda} \left(\epsilon^\theta \sum_{k=0}^{k_0} \left(\sum_{l=k-1}^{k+1} \sum_{j=k+2}^{\infty} \left\| 2^{k\alpha(\cdot)} B\tilde{T}_{\beta, \theta, \tilde{b}}(f_1^l, f_2^j) \chi_k \right\|_{q(\cdot)} \right)^{p(1+\epsilon)} \right)^{\frac{1}{p(1+\epsilon)}} \\
 & \leq C \sup_{\epsilon > 0} \sup_{k_0 \in \mathbb{Z}} 2^{-k_0 \lambda_1} \left(\epsilon^\theta \sum_{k=0}^{k_0} \left(\sum_{l=k-1}^{k+1} 2^{k\alpha_1(\infty)} \|f_1^l\|_{q_1(\cdot)} \right)^{p_1(1+\epsilon)} \right)^{\frac{1}{p_1(1+\epsilon)}} \\
 & \quad \times \sup_{\epsilon > 0} \sup_{k_0 \in \mathbb{Z}} 2^{-k_0 \lambda_2} \left(\epsilon^\theta \sum_{k=0}^{k_0} \left(\sum_{j=k+2}^{\infty} (j-k) 2^{j\alpha_2(\infty) + e_2(k-j)} \|f_2^j\|_{q_2(\cdot)} \right)^{p_2(1+\epsilon)} \right)^{\frac{1}{p_2(1+\epsilon)}} \\
 & = F_{62}^1 F_{62}^2.
 \end{aligned}$$

Since the estimates of F_{62}^1 and F_{62}^2 are similar F_{23}^2 and F_{23}^1 , respectively, we conclude

$$F_{62} \leq C \|f_1\|_{M\dot{K}_{p_1, \theta, q_1(\cdot)}^{\alpha_1(\cdot), \lambda_1}(\mathbb{R}^n)} \|f_2\|_{M\dot{K}_{p_2, \theta, q_2(\cdot)}^{\alpha_2(\cdot), \lambda_2}(\mathbb{R}^n)}.$$

For F_9 , we set $l, j \geq k+2$, $x \in D_k$, $y_1 \in D_l$ and $y_2 \in D_j$, such that $|x - y_1| \geq C2^l$ and $|x - y_2| \geq C2^j$. For $x \in D_k$, we have

$$\begin{aligned} & \left| B\tilde{T}_{\beta, \theta, \tilde{b}}(f_1^l, f_2^j)(x) \right| \\ & \leq C2^{-nl(1-\beta)}2^{-nj(1-\beta)} \|f_1^l\|_{q_1(\cdot)} \left(\|b_1(\cdot) - b_{B_k}\| \|\chi_l\|_{q'_1(\cdot)} + \|b_{B_k} - b_1\| \|\chi_l\|_{q'_1(\cdot)} \right) \\ & \quad \times \|f_2^j\|_{q_2(\cdot)} \left(\|b_2(\cdot) - b_{B_k}\| \|\chi_j\|_{q'_2(\cdot)} + \|b_{B_k} - b_2\| \|\chi_j\|_{q'_2(\cdot)} \right), \end{aligned} \quad (3.10)$$

by Minkowski's inequality, write

$$\begin{aligned} & \sup_{\epsilon > 0} \sup_{k_0 \in \mathbb{Z}} 2^{-k_0\lambda} \left(\epsilon^\theta \sum_{k=-\infty}^{k_0} \left(\sum_{l=k+2}^{\infty} \sum_{j=k+2}^{\infty} \left\| 2^{k\alpha(\cdot)} B\tilde{T}_{\beta, \theta, \tilde{b}}(f_1^l, f_2^j) \chi_k \right\|_{q(\cdot)} \right)^{p(1+\epsilon)} \right)^{\frac{1}{p(1+\epsilon)}} \\ & \leq \sup_{\epsilon > 0} \sup_{k_0 \in \mathbb{Z}} 2^{-k_0\lambda} \left(\epsilon^\theta \sum_{k=-\infty}^{-1} \left(\sum_{l=k+2}^{\infty} \sum_{j=k+2}^{\infty} \left\| 2^{k\alpha(\cdot)} B\tilde{T}_{\beta, \theta, \tilde{b}}(f_1^l, f_2^j) \chi_k \right\|_{q(\cdot)} \right)^{p(1+\epsilon)} \right)^{\frac{1}{p(1+\epsilon)}} \\ & \quad + \sup_{\epsilon > 0} \sup_{k_0 \in \mathbb{Z}} 2^{-k_0\lambda} \left(\epsilon^\theta \sum_{k=0}^{k_0} \left(\sum_{l=k+2}^{\infty} \sum_{j=k+2}^{\infty} \left\| 2^{k\alpha(\cdot)} B\tilde{T}_{\beta, \theta, \tilde{b}}(f_1^l, f_2^j) \chi_k \right\|_{q(\cdot)} \right)^{p(1+\epsilon)} \right)^{\frac{1}{p(1+\epsilon)}} \\ & = F_{91} + F_{92}. \end{aligned}$$

For F_{91} , by (3.5), (3.10), Lemmas 3.3 and 3.4, we obtain

$$\begin{aligned} F_{91} & \leq C \sup_{\epsilon > 0} \sup_{k_0 \in \mathbb{Z}} 2^{-k_0\lambda_1} \left(\epsilon^\theta \sum_{k=-\infty}^{-1} \left(\sum_{l=k+2}^{\infty} (l-k) 2^{k\alpha_1(0)+(k-l)\left(\frac{n}{q_1(\infty)} - n\beta\right)} \right. \right. \\ & \quad \times \|f_1^l\|_{q_1(\cdot)} \left. \left. \right)^{p_1(1+\epsilon)} \right)^{\frac{1}{p_1(1+\epsilon)}} \\ & \quad \times \sup_{\epsilon > 0} \sup_{k_0 \in \mathbb{Z}} 2^{-k_0\lambda_2} \left(\epsilon^\theta \sum_{k=-\infty}^{-1} \left(\sum_{j=k+2}^{\infty} (j-k) 2^{k\alpha_2(0)+(k-j)\left(\frac{n}{q_2(\infty)} - n\beta\right)} \right. \right. \\ & \quad \times \|f_2^j\|_{q_2(\cdot)} \left. \left. \right)^{p_2(1+\epsilon)} \right)^{\frac{1}{p_2(1+\epsilon)}} \\ & = F_{91}^1 F_{91}^2. \end{aligned}$$

It is easy to find that F_{91}^1 is similar to F_{32}^2 , hence, here we omit the details of the proof.

For F_{92} , it is easy to get

$$\begin{aligned} & \sup_{\epsilon > 0} \sup_{k_0 \in \mathbb{Z}} 2^{-k_0\lambda} \left(\epsilon^\theta \sum_{k=0}^{k_0} \left(\sum_{l=k+2}^{\infty} \sum_{j=k+2}^{\infty} \left\| 2^{k\alpha(\cdot)} B\tilde{T}_{\beta, \theta, \tilde{b}}(f_1^l, f_2^j) \chi_k \right\|_{q(\cdot)} \right)^{p(1+\epsilon)} \right)^{\frac{1}{p(1+\epsilon)}} \\ & \leq C \sup_{\epsilon > 0} \sup_{k_0 \in \mathbb{Z}} 2^{-k_0\lambda_1} \left(\epsilon^\theta \sum_{k=0}^{k_0} \left(\sum_{l=k+2}^{\infty} (l-k) 2^{l\alpha_1(\infty)+e_1(k-l)} \|f_1^l\|_{q_1(\cdot)} \right)^{p_1(1+\epsilon)} \right)^{\frac{1}{p_1(1+\epsilon)}} \\ & \quad \times \sup_{\epsilon > 0} \sup_{k_0 \in \mathbb{Z}} 2^{-k_0\lambda_2} \left(\epsilon^\theta \sum_{k=0}^{k_0} \left(\sum_{j=k+2}^{\infty} (j-k) 2^{j\alpha_2(\infty)+e_2(k-j)} \|f_2^j\|_{q_2(\cdot)} \right)^{p_2(1+\epsilon)} \right)^{\frac{1}{p_2(1+\epsilon)}} \end{aligned}$$

$$= F_{92}^1 F_{92}^2.$$

Since the estimates for F_{92}^1 and F_{92}^2 are similar to F_{32}^2 . So, we have

$$F_{92} \leq C \|f_1\|_{MK_{p_1(\cdot), \theta, q_1(\cdot)}^{\alpha_1(\cdot), \lambda_1}(\mathbb{R}^n)} \|f_2\|_{MK_{p_2(\cdot), \theta, q_2(\cdot)}^{\alpha_2(\cdot), \lambda_2}(\mathbb{R}^n)}.$$

Which, combining the estimates for $F_1, F_2, \dots, F_9$, yields the desired result. Hence, the proof of Theorem 3.2 is finished. $\square$

4. Estimate for $B\tilde{T}_{\beta, \theta}$ and $B\tilde{T}_{\beta, \theta, b_1, b_2}$ on $\dot{K}_{q(\cdot)}^{\alpha(\cdot), p, \theta}(\mathbb{R}^n)$

In this section, the authors mainly consider the θ -type generalized fractional integral operators $B\tilde{T}_{\beta, \theta}$ and their commutators $B\tilde{T}_{\beta, \theta, b_1, b_2}$ generated by $b_1, b_2 \in \text{BMO}(\mathbb{R}^n)$ and the $B\tilde{T}_{\beta, \theta}$ are bounded on grand variable exponents Herz spaces $\dot{K}_{q(\cdot)}^{\alpha(\cdot), p, \theta}(\mathbb{R}^n)$ (or $K_{q(\cdot)}^{\alpha(\cdot), p, \theta}(\mathbb{R}^n)$). The main results of this section are stated as follows:

Theorem 4.1. *Let $\theta > 1$, $b_1, b_2 \in \text{BMO}(\mathbb{R}^n)$, $\alpha(\cdot)$ and $q(\cdot)$ satisfy conditions (2.7) and (2.8). Suppose that $\frac{1}{q(x)} = \frac{1}{r_1(x)} + \frac{1}{r_2(x)}$, $\frac{1}{r_i(x)} = \frac{1}{q_i(x)} - \beta$ and $n\beta - \frac{n}{q_i(\infty)} < \alpha_i(0) \leq \alpha_i(\infty) < n - \frac{n}{q_i(0)}$ are established. Then there exists a positive constant C such that, for any $f_i \in \dot{K}_{q(\cdot)}^{\alpha(\cdot), p, \theta}(\mathbb{R}^n)$,*

$$\|B\tilde{T}_{\beta, \theta}(f_1, f_2)\|_{\dot{K}_{q(\cdot)}^{\alpha(\cdot), p, \theta}(\mathbb{R}^n)} \leq C \|f_1\|_{\dot{K}_{q(\cdot)}^{\alpha(\cdot), p, \theta}(\mathbb{R}^n)} \|f_2\|_{\dot{K}_{q(\cdot)}^{\alpha(\cdot), p, \theta}(\mathbb{R}^n)},$$

where $\alpha(x) = \alpha_1(x) + \alpha_2(x)$ and $\frac{1}{p} = \frac{1}{p_1} + \frac{1}{p_2}$.

Theorem 4.2. *Let $\theta > 1$, $b_1, b_2 \in \text{BMO}(\mathbb{R}^n)$, $\alpha(\cdot)$ and $q(\cdot)$ satisfy conditions (2.7) and (2.8). Suppose that $\frac{1}{q(x)} = \frac{1}{r_1(x)} + \frac{1}{r_2(x)}$, $\frac{1}{r_i(x)} = \frac{1}{q_i(x)} - \beta$ and $n\beta - \frac{n}{q_i(\infty)} < \alpha_i(0) \leq \alpha_i(\infty) < n - \frac{n}{q_i(0)}$ are established. Then there exists a positive constant C such that, for any $f_i \in \dot{K}_{q(\cdot)}^{\alpha(\cdot), p, \theta}(\mathbb{R}^n)$,*

$$\|B\tilde{T}_{\beta, \theta, b_1, b_2}(f_1, f_2)\|_{\dot{K}_{q(\cdot)}^{\alpha(\cdot), p, \theta}(\mathbb{R}^n)} \leq C \|b_1\|_* \|b_2\|_* \|f_1\|_{\dot{K}_{q(\cdot)}^{\alpha(\cdot), p, \theta}(\mathbb{R}^n)} \|f_2\|_{\dot{K}_{q(\cdot)}^{\alpha(\cdot), p, \theta}(\mathbb{R}^n)},$$

where $\alpha(x) = \alpha_1(x) + \alpha_2(x)$ and $\frac{1}{p} = \frac{1}{p_1} + \frac{1}{p_2}$.

Remark 4.1. (a) Theorems 4.1 and 4.2 can be shown to be true in the same way as Theorems 3.1 and 3.2. Consequently, a detailed presentation of the respective proofs is not included here; (b) Given that the proof for the non-homogeneous case can be treated by a similar method, the results presented of theorems 4.1 and 4.2 are applicable to the non-homogeneous grand variable exponents Herz spaces.

Acknowledgements

The authors would like to thank the referees for their valuable comments and suggestions, which improved the quality of this paper.

References

- [1] I. Aarab and M. A. Tagmouti, *Harmonic oscillator perturbed by a decreasing scalar potential*, J. Pseudo-Differ. Oper. Appl., 2020, 11(1), 141–157.
- [2] E. Bereznoi and A. Karapetyants, *Grand and small norms in Lebesgue spaces*, Math. Methods Appl. Sci., 2024, 47(2), 725–741.
- [3] X. J. Chen, G. H. Lu and W. W. Tao, *Fractional type Marcinkiewicz integral and its commutator on grand variable Herz-Morrey spaces*, J. Pseudo-Differ. Oper. Appl., 2024, 15(3), 1–19.
- [4] D. Cruz-Uribe, A. Fiorenza, J. M. Martell and C. Pérez, *The boundedness of classical operators on variable L^p spaces*, Ann. Acad. Sci. Fenn. Math., 2006, 31(1), 239–264.
- [5] X. Fu, D. C. Yang and W. Yuan, *Generalized fractional integrals and their commutators over non-homogeneous metric measure spaces*, Taiwan. J. Math., 2014, 18, 509–557.
- [6] G. E. Hu, Y. Meng and D. C. Yang, *Weighted norm inequalities for multilinear Calderón-Zygmund operators on non-homogeneous metric measure spaces*, Forum Math., 2014, 26(5), 1289–1322.
- [7] A. W. Huang and J. S. Xu, *Multilinear singular integrals and commutators in variable exponent Lebesgue spaces*, Appl. Math. J. Chinese Univ. Ser., 2010, 25(1), 69–77.
- [8] T. Iwaniec and C. Sbordone, *On the integrability of the Jacobian under minimal hypotheses*, Arch. Ration. Mech. Anal., 1992, 119(2), 129–143.
- [9] M. Izuki, *Herz and amalgam spaces with variable exponent, the Haar wavelets and greediness of the wavelet system*, East J. Approx., 2009, 15(1), 87–109.
- [10] M. Izuki, *Boundedness of commutators on Herz spaces with variable exponent*, Rend. Circ. Mat. Palermo., 2010, 59(2), 199–213.
- [11] G. H. Lu, *Bilinear θ -type Calderón-Zygmund operator and its commutator on non-homogeneous weighted Morrey spaces*, Rev. R. Acad. Cienc. Exactas Fís. Nat. Ser. A Mat. RACSAM, 2021, 115(1), 1–15.
- [12] G. H. Lu and L. Rui, *θ -type generalized fractional integral and its commutator on some non-homogeneous variable exponent spaces*, AIMS Math., 2021, 6(9), 9619–9632.
- [13] G. H. Lu and S. P. Tao, *Bilinear θ -type generalized fractional integral operator and its commutator on some non-homogeneous spaces*, Bull. Sci. Math., 2022, 174, 1–32.
- [14] G. H. Lu and S. P. Tao, *Two-weighted estimate for generalized fractional integral and its commutator on generalized fractional Morrey spaces*, J. Math. Study., 2023, 56(4), 345–356.
- [15] G. H. Lu and S. P. Tao, *Bilinear θ -type generalized fractional integral and its commutator on nonhomogeneous metric measure spaces*, Rocky Mountain J. Math., 2023, 53(3), 839–857.
- [16] G. H. Lu and S. P. Tao, *Two classes of bilinear fractional integral operators and their commutators on generalized fractional Morrey spaces*, J. Pseudo-Differ. Oper. Appl., 2021, 12(4), 1–24.

- [17] G. H. Lu, S. P. Tao and R. H. Liu, *θ -type Calderón-Zygmund operator and its commutator on (grand) generalized weighted variable exponent Morrey space over RD-spaces*, J. Pseudo-Differ. Oper. Appl., 2023, 14(4), 1–22.
- [18] G. H. Lu, S. P. Tao and M. M. Wang, *Estimates for bilinear generalized fractional integral operator and its commutator on generalized Morrey spaces over RD-spaces*, Ann. Funct. Anal., 2024, 15(1), 1–47.
- [19] G. H. Lu, M. M. Wang and S. P. Tao, *Estimates for bilinear θ -type generalized fractional integral and its commutator on new non-homogeneous generalized Morrey spaces*, Anal. Geom. Metr. Spaces, 2023, 11(1), 1–24.
- [20] H. Nafis, H. Rafeiro and M. A. Zaighum, *A note on the boundedness of sublinear operators on grand variable Herz spaces*, J. Inequal. Appl., 2020, 2020(1), 1–13.
- [21] H. Nafis, H. Rafeiro and M. A. Zaighum, *Boundedness of multilinear Calderón-Zygmund operators on grand variable Herz spaces*, J. Funct. Spaces, 2022, 2022(1), 1–11.
- [22] B. Sultan, F. Azmi, M. Sultan, M. Mehmood and N. Mlaiki, *Boundedness of Riesz potential operator on grand Herz-Morrey spaces*, Axioms, 2022, 11(583), 1–14.
- [23] B. Sultan, M. Sultan and F. Gübüç, *BMO estimate for the higher order commutators of Marcinkiewicz integral operator on grand variable Herz-Morrey spaces*, Commun. Fac. Sci. Univ. Ank. Ser. A1. Math. Stat., 2023, 72(4), 1000–1018.
- [24] M. Sultan, B. Sultan and A. Hussain, *Grand Herz-Morrey spaces with variable exponent*, Math. Notes., 2023, 114(5–6), 957–977.
- [25] L. W. Wang, *Parametrized Littlewood-Paley operators on grand variable Herz spaces*, Ann. Funct. Anal., 2022, 13(4), 1–26.
- [26] S. Yang, J. W. Sun and B. D. Li, *Maximal and Calderón-Zygmund operators in grand variable Lebesgue spaces*, Banach J. Math. Anal., 2023, 17(3), 1–28.

Received November 2024; Accepted February 2025; Available online April 2025.