

# EXISTENCE AND STABILITY OF SOLUTIONS FOR HADAMARD TYPE FRACTIONAL DIFFERENTIAL SYSTEM ON BENZENE GRAPHS\*

Yunzhe Zhang<sup>1,2</sup>, Youhui Su<sup>1,†</sup> and Yongzhen Yun<sup>1</sup>

**Abstract** This paper is mainly concerned with the existence of solutions for a class of Hadamard type fractional differential systems on benzene graphs, and the Hyers-Ulam stability of the systems is also proved. Furthermore, an example is presented on a formic acid graph to demonstrate the applicability of the conclusions obtained. The interesting of this paper lies in the integration of fractional differential equations with graph theory, utilizing the formic acid graph as a specific case for numerical simulation, and providing an approximate solution graph after iterations.

**Keywords** Fractional differential equation, benzene graphs, Hyers-Ulam stability, numerical simulation.

**MSC(2010)** 34A08, 34K37, 34B18.

## 1. Introduction

The fractional differential equation is a generalization of an integer-order differential equation, allowing for a more accurate description of complex phenomena in nature and engineering. For instance, fractional differential equations provide a more appropriate model for describing diffusion processes, wave phenomena and memory effects [9, 10, 13, 16, 18], and possess a diverse array of applications across numerous fields, encompassing stochastic equations, fluid flow, dynamical systems theory, physics, biology, and other domains [4–6, 20, 22].

Star graph  $G = (V, E)$  consists of a finite set of nodes or vertices  $V(G) = \{v_0, v_1, \dots, v_k\}$  and a set of edges  $E(G) = \{e_1 = \overrightarrow{v_1 v_0}, e_2 = \overrightarrow{v_2 v_0}, \dots, e_k = \overrightarrow{v_k v_0}\}$  connecting these nodes, where  $v_0$  is the joint point and  $e_i$  is the length of  $l_i$  the edge connecting the nodes  $v_i$  and  $v_0$ , i.e.  $l_i = |\overrightarrow{v_i v_0}|$ .

<sup>†</sup>The corresponding author.

<sup>1</sup>School of Mathematics and Statistics, Xuzhou University of Technology, Xuzhou 221018, China

<sup>2</sup>College of science, Shenyang University of Technology, Shenyang 110870, China

\*The authors were supported by the Xuzhou Science and Technology Plan Project (KC23058), the Natural Science Foundation of Xinjiang Uygur Autonomous Region(2023D01C51) and the Natural Science Research Project of Jiangsu Colleges and Universities (22KJB110026).

Email: 18015151008@163.com(Y. Zhang), suyh02@163.com(Y. Su), yongzhen0614@163.com(Y. Yun)

Graph theory is a mathematical discipline that investigates graphs and networks. A network is a graph, such as computer network extensions, transportation route maps, molecules in medicine and biology, and so on [1, 15]. Graph theory has become widely applied in sociology, traffic management, telecommunications and other fields [7, 23].

As is well known, differential equations on star graphs can be applied to different fields, such as chemistry, bioengineering and so on [8, 11]. Mehendiratta et al [12] explored the fractional differential system on star graphs with  $n + 1$  nodes and  $n$  edges

$$\begin{cases} {}^C D_{0,x}^\alpha u_i(x) = f_i(x, u_i, {}^C D_{0,x}^\beta u_i(x)), & 0 < x < l_i, i = 1, 2, \dots, k, \\ u_i(0) = 0, & i = 1, 2, \dots, k, \\ u_i(l_i) = u_j(l_j), & i, j = 1, 2, \dots, k, i \neq j, \\ \sum_{i=1}^k u_i' = 0, & i = 1, 2, \dots, k, \end{cases}$$


where  ${}^C D_{0,x}^\alpha$ ,  ${}^C D_{0,x}^\beta$  are the Caputo fractional derivative operator,  $1 < \alpha \leq 2$ ,  $0 < \beta \leq \alpha - 1$ ,  $f_i, i = 1, 2, \dots, k$  are continuous functions on  $C([0, 1] \times \mathbb{R} \times \mathbb{R})$ . By a transformation, the equivalent fractional differential system defined on  $[0, 1]$  is obtained. The author studied a nonlinear Caputo fractional boundary value problem on star graphs and established the existence and uniqueness results by fixed point theory.

Zhang et al. [25] discussed the fractional boundary value problem on star graphs, and obtained the existence and uniqueness results of solutions by fixed point theory. In addition, Wang et al [24] discussed the existence and stability of a fractional differential equation with Hadamard derivative. For more papers on the existence of solutions to fractional differential equations on graphs, refer to [2, 3, 17, 21]. By numerically simulating the solution of fractional differential systems, scholars can solve problems more clearly and accurately. However, numerical simulation has been rarely used to describe the solutions of fractional differential systems on graphs [14, 19, 26].


Here, we introduce a novel modeling of fractional boundary value problems on the benzene graph (Figure 1). The molecular structure of the benzene is as ring containing six carbon atoms and six hydrogen atoms. Benzene stands as a pivotal raw material in the petrochemical industry, encompassing a diverse array of applications. Therefore, a thorough understanding of its properties is of utmost importance.

By this structure, we consider atoms of carbons and hydrogens as vertices of the graph and also the existing chemical bonds between atoms are considered as edges of the graph. To investigate the existence of solutions for our fractional boundary value problems in the sequel, we label vertices of the benzene graph in the form of labeled vertices by two values 0 or 1 and the length of each edge is fixed at  $e$  ( $|\vec{e}_i| = e$ ,  $i = 1, 2, \dots, 12$ ) (Figure 2). In this case, we construct a local coordinate system on the benzene graph and the orientation of each vertex is determined by the orientation of its corresponding edge. The labels of the beginning and ending vertices are taken into account as values 0 and 1, respectively, as we move along any edge.

Motivated by the above work and relevant literatures [2, 3, 12, 14, 17, 19, 21, 24–26],



**Figure 1.** Molecular structure of benzene.



**Figure 2.** Benzene graphs with vertices 0 or 1.

we study a boundary value problem consisting of nonlinear fractional differential equations defined on  $|\vec{e}_i| = e$ ,  $i = 1, 2, \dots, 12$  by

$${}^H D_{1+}^\alpha u_i(t) = \lambda_i^\alpha f_i(s, u_i(s), {}^H D_{1+}^\beta u_i(s)), \quad t \in [1, e],$$

and the boundary conditions defined at boundary nodes  $e_1, e_2, \dots, e_{12}$ , and

$$u_i(1) = 0, u_i(e) = u_j(e), \quad i, j = 1, 2, \dots, 12, \quad i \neq j,$$

together with conditions of conjunctions at 0 or 1 with

$$\sum_{i=1}^k \lambda_i^{-1} u'_i(e) = 0, \quad i = 1, 2, \dots, 12.$$

Overall, we consider the existence and stability of solutions to the following nonlinear boundary value problem on benzene graphs

$$\begin{cases} {}^H D_{1+}^\alpha u_i(t) = \lambda_i^\alpha f_i(s, u_i(s), {}^H D_{1+}^\beta u_i(s)), \quad t \in [1, e], \\ u_i(1) = 0, \quad i = 1, 2, \dots, 12, \\ u_i(e) = u_j(e), \quad i, j = 1, 2, \dots, 12, \quad i \neq j, \\ \sum_{i=1}^k \lambda_i^{-1} u'_i(e) = 0, \quad i = 1, 2, \dots, 12, \end{cases} \quad (1.1)$$

where  ${}^H D_{1+}^\alpha, {}^H D_{1+}^\beta$  represents the the Hadamard fractional derivative,  $\alpha \in (2, 3]$ ,  $\beta \in (1, 2]$ ,  $f_i \in C([1, e] \times \mathbb{R} \times \mathbb{R})$  and  $\lambda_i, i = 1, 2, \dots, 12$  is real constant. The existence and Hyers-Ulam stability of the solutions to the system (1.1) are discussed. Moreover, the approximate graphs of the solution are obtained.

It is also noteworthy that solutions obtained from the problem (1.1) can be depicted in various rational applications of organic chemistry. More precisely, any solution on an arbitrary edge can be described as the amount of bond polarity, bond power, bond energy etc. The interesting of this paper lies in the integration of fractional differential equations with graph theory, utilizing the formic acid graph as a specific case for numerical simulation, and providing an approximate solution graph after iterations.

## 2. Preliminaries

In this section, for conveniently researching the problem, several properties and lemmas of fractional calculus are given, forming the indispensable premises for obtaining the main conclusions.

**Definition 2.1.** [3,9] The Hadamard fractional integral of order  $\alpha$ , for a function  $g \in L^p[a, b]$ ,  $0 \leq a \leq t \leq b \leq \infty$ , is defined as

$${}^H I_{a+}^\alpha g(t) = \frac{1}{\Gamma(\alpha)} \int_a^t \left( \log \frac{t}{s} \right)^{\alpha-1} \frac{g(s)}{s} ds.$$

**Definition 2.2.** [3,9] Let  $[a, b] \subset \mathbb{R}$ ,  $\delta = t \frac{d}{dt}$  and  $AC_\delta^n[a, b] = \{g : [a, b] \rightarrow \mathbb{R} : \delta^{n-1}(g(t)) \in AC[a, b]\}$ . The Hadamard derivative of fractional order  $\alpha$  for a function  $g \in AC_\delta^n[a, b]$  is defined as

$${}^H D_{a+}^\alpha g(t) = \delta^n ({}^H I_{a+}^{n-\alpha})(t) = \frac{1}{\Gamma(n-\alpha)} \left( t \frac{d}{dt} \right)^n \int_a^t \left( \log \frac{t}{s} \right)^{n-\alpha-1} \frac{g(s)}{s} ds,$$

where  $n-1 < \alpha < n$ ,  $n = [\alpha] + 1$  and  $[\alpha]$  denotes the integer part of the real number  $\alpha$  and  $\log(\cdot) = \log_e(\cdot)$ .

**Lemma 2.1.** [3] For  $y \in AC_\delta^n[a, b]$ , the following result hold

$${}^H I_{0+}^\alpha ({}^H D_{0+}^\alpha) y(t) = y(t) - \sum_{k=0}^{n-1} c_k \left( \log \frac{t}{a} \right)^k,$$

where  $c_i \in \mathbb{R}$ ,  $i = 0, 1, \dots, n-1$ .

**Lemma 2.2.** (Scheafer's fixed point theorem) [8] Let  $X$  be a Banach space and let  $F : X \rightarrow X$  be a completely continuous operator (i.e., an operator that restricted to any bounded set in  $X$  is compact). Then either

- (i) The set  $\{x \in X : x = \mu Fx \text{ for some } \mu \in (0, 1)\}$  is unbounded, or
- (ii)  $F$  has at least one fixed point in  $X$ .

**Lemma 2.3.** Let  $h_i(t) \in AC([1, e], \mathbb{R})$ ,  $i = 1, 2, \dots, 12$ , then the solution of the fractional differential equations

$$\begin{cases} {}^H D_{1+}^\alpha u_i(t) = h_i(t), & t \in [1, e], \\ u_i(1) = 0, & i = 1, 2, \dots, 12, \\ u_i(e) = u_j(e), & i, j = 1, 2, \dots, 12, \quad i \neq j, \\ \sum_{i=1}^k \lambda_i^{-1} u'_i(e) = 0, & i = 1, 2, \dots, 12, \end{cases} \quad (2.1)$$

is given by

$$\begin{aligned} u_i(t) = & \frac{1}{\Gamma(\alpha)} \int_1^t \left( \log \frac{t}{s} \right)^{\alpha-1} \frac{h_i(s)}{s} ds \\ & - \log t \left[ \frac{1}{\Gamma(\alpha-1)} \sum_{j=1}^k \left( \frac{\lambda_j^{-1}}{\sum_{j=1}^k \lambda_j^{-1}} \right) \int_1^e \left( \log \frac{e}{s} \right)^{\alpha-2} \frac{h_j(s)}{s} ds \right. \\ & \left. - \frac{1}{\Gamma(\alpha)} \sum_{\substack{j=1 \\ j \neq i}}^k \left( \frac{\lambda_j^{-1}}{\sum_{j=1}^k \lambda_j^{-1}} \right) \left( \int_1^e \left( \log \frac{e}{s} \right)^{\alpha-1} \left( \frac{h_j(s) - h_i(s)}{s} \right) ds \right) \right]. \end{aligned} \quad (2.2)$$

**Proof.** By Lemma 2.1, we have

$$u_i(t) = {}^H I_{1+}^\alpha h_i(t) - c_i^{(1)} - c_i^{(2)} \log t, \quad i = 1, 2, \dots, 12,$$

where  $c_i^{(1)}$ ,  $c_i^{(2)}$  are constants. The boundary condition  $u_i(1) = 0$ , gives  $c_i^{(1)} = 0$ , for  $i = 1, 2, \dots, 12$ .

Hence,

$$\begin{aligned} u_i(t) &= {}^H I_{1+}^\alpha h_i(t) - c_i^{(2)} \log t \\ &= \frac{1}{\Gamma(\alpha)} \int_1^t \left( \log \frac{t}{s} \right)^{\alpha-1} \frac{h(s)}{s} ds - c_i^{(2)} \log t, \quad i = 1, 2, \dots, 12. \end{aligned} \quad (2.3)$$

Also

$$u'_i(t) = \frac{1}{\Gamma(\alpha-1)} \int_1^t \frac{1}{t} \left( \log \frac{t}{s} \right)^{\alpha-2} \frac{h(s)}{s} ds - \frac{1}{t} c_i^{(2)}.$$

Now, the boundary conditions  $u_i(e) = u_j(e)$  and  $\sum_{i=1}^k \lambda_i^{-1} u'_i(e) = 0$  implies that  $c_i^{(2)}$  must satisfy

$$\frac{1}{\Gamma(\alpha)} \int_1^e \left( \log \frac{e}{s} \right)^{\alpha-1} \frac{h_i(s)}{s} ds - c_i^{(2)} = \frac{1}{\Gamma(\alpha)} \int_1^e \left( \log \frac{e}{s} \right)^{\alpha-1} \frac{h_j(s)}{s} ds - c_j^{(2)}, \quad (2.4)$$

$$\sum_{i=1}^k \lambda_i^{-1} \left( \frac{1}{\Gamma(\alpha-1)} \int_1^e \left( \log \frac{e}{s} \right)^{\alpha-2} \frac{h_i(s)}{s} ds - c_i^{(2)} \right) = 0. \quad (2.5)$$

On solving above equations 2.4 and 2.5, we have

$$\sum_{j=1}^k \lambda_j^{-1} \left( \frac{1}{\Gamma(\alpha-1)} \int_1^e \left( \log \frac{e}{s} \right)^{\alpha-2} \frac{h_j(s)}{s} ds \right) - \lambda_i^{-1} c_i^{(2)}$$

$$= \sum_{\substack{j=1 \\ j \neq i}}^k \lambda_j^{-1} \left[ \frac{1}{\Gamma(\alpha)} \int_1^e \left( \log \frac{e}{s} \right)^{\alpha-1} \frac{h_j(s)}{s} ds - \frac{1}{\Gamma(\alpha)} \int_1^e \left( \log \frac{e}{s} \right)^{\alpha-1} \frac{h_i(s)}{s} ds + c_i^{(2)} \right],$$

which implies

$$\begin{aligned} \sum_{j=1}^k \lambda_j^{-1} c_i^{(2)} &= \sum_{j=1}^k \lambda_j^{-1} \frac{1}{\Gamma(\alpha-1)} \int_1^e \left( \log \frac{e}{s} \right)^{\alpha-2} \frac{h_j(s)}{s} ds \\ &\quad - \sum_{\substack{j=1 \\ j \neq i}}^k \lambda_j^{-1} \frac{1}{\Gamma(\alpha)} \int_1^e \left( \log \frac{e}{s} \right)^{\alpha-1} \left( \frac{h_j(s) - h_i(s)}{s} \right) ds. \end{aligned}$$

Hence, we get

$$\begin{aligned} c_i^{(2)} &= \frac{1}{\Gamma(\alpha-1)} \sum_{j=1}^k \left( \frac{\lambda_j^{-1}}{\sum_{j=1}^k \lambda_j^{-1}} \right) \int_1^e \left( \log \frac{e}{s} \right)^{\alpha-2} \frac{h_j(s)}{s} ds \\ &\quad - \frac{1}{\Gamma(\alpha)} \sum_{\substack{j=1 \\ j \neq i}}^k \left( \frac{\lambda_j^{-1}}{\sum_{j=1}^k \lambda_j^{-1}} \right) \left( \int_1^e \left( \log \frac{e}{s} \right)^{\alpha-1} \left( \frac{h_j(s) - h_i(s)}{s} \right) ds \right). \quad (2.6) \end{aligned}$$

Hence, inserting the values of  $c_i^{(2)}$ , we get the solution (2.2). This completes the proof.  $\square$

### 3. Main results

In this section, the existence and uniqueness to solutions of system (1.1) are discussed.

We define the space  $X = \{u : u \in C([1, e], \mathbb{R}), {}^H D_{1+}^\beta u \in C([1, e], \mathbb{R})\}$  with the norm

$$\|u\|_X = \|u\| + \left\| {}^H D_{1+}^\beta u \right\| = \sup_{t \in [1, e]} |u(t)| + \sup_{t \in [1, e]} \left| {}^H D_{1+}^\beta u(t) \right|.$$

Then,  $(X, \|\cdot\|_X)$  is a Banach space and accordingly, the product space  $(X^k = X_1 \times X_2 \cdots \times X_{12}, \|\cdot\|_{X^k})$  is a Banach space with norm

$$\|u\|_{X^k} = \|(u_1, u_2, \dots, u_{12})\|_X = \sum_{i=1}^k \|u_i\|_X, \quad (u_1, u_2, \dots, u_k) \in X^k.$$

In view of Lemma 2.3, we define the operator  $T : X^k \rightarrow X^k$  by

$$T(u_1, u_2, \dots, u_k)(t) := (T_1(u_1, u_2, \dots, u_k)(t), \dots, T_k(u_1, u_2, \dots, u_k)(t)),$$

where

$$\begin{aligned} &T_i(u_1, u_2, \dots, u_k)(t) \\ &= \frac{\lambda_i^\alpha}{\Gamma(\alpha)} \int_1^t \left( \log \frac{t}{s} \right)^{\alpha-1} \frac{f_i(s, u_i(s), {}^H D_{1+}^\beta u_i(s))}{s} ds \end{aligned}$$

$$\begin{aligned}
& -\frac{\log t}{\Gamma(\alpha-1)} \sum_{j=1}^k \left( \frac{\lambda_j^{-1}}{\sum_{j=1}^k \lambda_j^{-1}} \right) \lambda_j^\alpha \int_1^e \left( \log \frac{e}{s} \right)^{\alpha-2} \frac{f_j(s, u_j(s), {}^H D_{1+}^\beta u_j(s))}{s} ds \\
& + \frac{\log t}{\Gamma(\alpha)} \sum_{\substack{j=1 \\ j \neq i}}^k \left( \frac{\lambda_j^{-1}}{\sum_{j=1}^k \lambda_j^{-1}} \right) \lambda_j^\alpha \int_1^e \left( \log \frac{e}{s} \right)^{\alpha-1} \frac{f_j(s, u_j(s), {}^H D_{1+}^\beta u_j(s))}{s} ds \\
& - \frac{\log t}{\Gamma(\alpha)} \sum_{\substack{j=1 \\ j \neq i}}^k \left( \frac{\lambda_j^{-1}}{\sum_{j=1}^k \lambda_j^{-1}} \right) \lambda_i^\alpha \int_1^e \left( \log \frac{e}{s} \right)^{\alpha-1} \frac{f_i(s, u_i(s), {}^H D_{1+}^\beta u_i(s))}{s} ds.
\end{aligned}$$

Let  $\frac{f_i(s, u_i(s), {}^H D_{1+}^\beta u_i(s))}{s} = g_i(s, u_i(s), {}^H D_{1+}^\beta u_i(s))$ , then

$$\begin{aligned}
& T_i(u_1, u_2, \dots, u_k)(t) \\
& = \frac{\lambda_i^\alpha}{\Gamma(\alpha)} \int_1^t \left( \log \frac{t}{s} \right)^{\alpha-1} g_i(s, u_i(s), {}^H D_{1+}^\beta u_i(s)) ds \\
& - \frac{\log t}{\Gamma(\alpha-1)} \sum_{j=1}^k \left( \frac{\lambda_j^{-1}}{\sum_{j=1}^k \lambda_j^{-1}} \right) \lambda_j^\alpha \int_1^e \left( \log \frac{e}{s} \right)^{\alpha-2} g_j(s, u_j(s), {}^H D_{1+}^\beta u_j(s)) ds \\
& + \frac{\log t}{\Gamma(\alpha)} \sum_{\substack{j=1 \\ j \neq i}}^k \left( \frac{\lambda_j^{-1}}{\sum_{j=1}^k \lambda_j^{-1}} \right) \lambda_j^\alpha \int_1^e \left( \log \frac{e}{s} \right)^{\alpha-1} g_j(s, u_j(s), {}^H D_{1+}^\beta u_j(s)) ds \\
& - \frac{\log t}{\Gamma(\alpha)} \sum_{\substack{j=1 \\ j \neq i}}^k \left( \frac{\lambda_j^{-1}}{\sum_{j=1}^k \lambda_j^{-1}} \right) \lambda_i^\alpha \int_1^e \left( \log \frac{e}{s} \right)^{\alpha-1} g_i(s, u_i(s), {}^H D_{1+}^\beta u_i(s)) ds. \quad (3.1)
\end{aligned}$$

Assume that the following conditions hold:

(H<sub>1</sub>)  $g_i : [1, e] \times \mathbb{R} \times \mathbb{R} \rightarrow \mathbb{R}, i = 1, 2, \dots, 12$  be continuous functions and there exists nonnegative functions  $l_i(t) \in C[1, e]$  such that

$$|g_i(t, x, y) - g_i(t, x_1, y_1)| \leq l_i(t)(|x - x_1| + |y - y_1|),$$

where  $t \in [1, e], (x, y), (x_1, y_1) \in \mathbb{R}^2$ ;

(H<sub>2</sub>)  $\omega_i = \sup_{t \in [1, e]} |l_i(t)|, i = 1, 2, \dots, 12$ ;

(H<sub>3</sub>) There exists  $A_i > 0$ , such that

$$|g_i(t, x, y)| \leq A_i, \quad t \in [1, e], \quad (x, y) \in \mathbb{R} \times \mathbb{R}, \quad i = 1, 2, \dots, 12.$$

For computational convenience, we also set the following quantities

$$\begin{aligned}
B_i = & e(\lambda_i^\alpha + \lambda_i^{\alpha-\beta}) \left[ \frac{1}{\Gamma(\alpha)} + \frac{2}{\Gamma(\alpha+1)} + \frac{1}{\Gamma(\alpha-\beta+1)} \right. \\
& \left. + \frac{1}{\Gamma(\alpha)\Gamma(2-\beta)} + \frac{1}{\Gamma(\alpha+1)\Gamma(2-\beta)} \right]
\end{aligned}$$

$$+ e \sum_{\substack{j=1 \\ j \neq i}}^k (\lambda_j^\alpha + \lambda_j^{\alpha-\beta}) \left[ \frac{1}{\Gamma(\alpha)} + \frac{1}{\Gamma(\alpha+1)} + \frac{1}{\Gamma(\alpha)\Gamma(2-\beta)} + \frac{1}{\Gamma(\alpha+1)\Gamma(2-\beta)} \right], \quad (3.2)$$

$$C_i = e \lambda_i^\alpha \left[ \frac{1}{\Gamma(\alpha)} + \frac{2}{\Gamma(\alpha+1)} + \frac{1}{\Gamma(\alpha-\beta+1)} + \frac{1}{\Gamma(\alpha)\Gamma(2-\beta)} + \frac{1}{\Gamma(\alpha+1)\Gamma(2-\beta)} \right] \\ + e \sum_{\substack{j=1 \\ j \neq i}}^k \lambda_j^\alpha \left[ \frac{1}{\Gamma(\alpha)} + \frac{1}{\Gamma(\alpha+1)} + \frac{1}{\Gamma(\alpha)\Gamma(2-\beta)} + \frac{1}{\Gamma(\alpha+1)\Gamma(2-\beta)} \right]. \quad (3.3)$$

**Theorem 3.1.** Assume that (H1) and (H2) hold, then the fractional differential system (1.1) has a unique solution on  $[1, e]$  if

$$\left( \sum_{i=1}^k B_i \right) \left( \sum_{i=1}^k \omega_i \right) < 1,$$

where  $B_i$ ,  $i = 1, 2, \dots, 12$  are given by equation (3.2).

**Proof.** Let  $u = (u_1, u_2, \dots, u_{12})$ ,  $v = (v_1, v_2, \dots, v_{12}) \in X^k$ ,  $t \in [1, e]$ , we have

$$\begin{aligned} & |T_i u(t) - T_i v(t)| \\ & \leq \frac{\lambda_i^\alpha}{\Gamma(\alpha)} \int_1^t \left( \log \frac{t}{s} \right)^{\alpha-1} \left| g_i(s, u_i(s), {}^H D_{1+}^\beta u_i(s)) - g_i(s, v_i(s), {}^H D_{1+}^\beta v_i(s)) \right| ds \\ & \quad + \frac{\log t}{\Gamma(\alpha-1)} \sum_{j=1}^k \left( \frac{\lambda_j^{-1}}{\sum_{j=1}^k \lambda_j^{-1}} \right) \lambda_j^\alpha \int_1^e \left( \log \frac{e}{s} \right)^{\alpha-2} \left[ \left| g_j(s, u_j(s), {}^H D_{1+}^\beta u_j(s)) \right. \right. \\ & \quad \left. \left. - g_j(s, v_j(s), {}^H D_{1+}^\beta v_j(s)) \right| ds \right] \\ & \quad + \frac{\log t}{\Gamma(\alpha)} \sum_{\substack{j=1 \\ j \neq i}}^k \left( \frac{\lambda_j^{-1}}{\sum_{j=1}^k \lambda_j^{-1}} \right) \lambda_j^\alpha \int_1^e \left( \log \frac{e}{s} \right)^{\alpha-1} \left[ \left| g_j(s, u_j(s), {}^H D_{1+}^\beta u_j(s)) \right. \right. \\ & \quad \left. \left. - g_j(s, v_j(s), {}^H D_{1+}^\beta v_j(s)) \right| ds \right] \\ & \quad + \frac{\log t}{\Gamma(\alpha)} \sum_{\substack{j=1 \\ j \neq i}}^k \left( \frac{\lambda_j^{-1}}{\sum_{j=1}^k \lambda_j^{-1}} \right) \lambda_i^\alpha \int_1^e \left( \log \frac{e}{s} \right)^{\alpha-1} \left[ \left| g_i(s, u_i(s), {}^H D_{1+}^\beta u_i(s)) \right. \right. \\ & \quad \left. \left. - g_i(s, v_i(s), {}^H D_{1+}^\beta v_i(s)) \right| ds \right]. \end{aligned}$$

Using (H1) and (H2),  $t \in [1, e]$  and  $\left( \frac{\lambda_j^{-1}}{\sum_{j=1}^k \lambda_j^{-1}} \right) < 1$  for  $j = 1, 2, \dots, k$ , we obtain

$$\begin{aligned} & |T_i u(t) - T_i v(t)| \\ & \leq \frac{e \lambda_i^\alpha}{\Gamma(\alpha+1)} \omega_i \|u_i - v_i\| + \frac{e \lambda_i^{\alpha-\beta}}{\Gamma(\alpha+1)} \omega_i \left\| {}^H D_{1+}^\beta u_i - {}^H D_{1+}^\beta v_i \right\| \end{aligned}$$



$$\begin{aligned}
& + e \sum_{j=1}^k \left( \frac{\lambda_j^\alpha}{\Gamma(\alpha)} \omega_j \|u_j - v_j\| + \frac{\lambda_j^{\alpha-\beta}}{\Gamma(\alpha)} \omega_j \left\| {}^H D_{1+}^\beta u_j - {}^H D_{1+}^\beta v_j \right\| \right) \\
& + e \sum_{\substack{j=1 \\ j \neq i}}^k \left( \frac{\lambda_j^\alpha}{\Gamma(\alpha+1)} \omega_j \|u_j - v_j\| + \frac{\lambda_j^{\alpha-\beta}}{\Gamma(\alpha+1)} \omega_j \left\| {}^H D_{1+}^\beta u_j - {}^H D_{1+}^\beta v_j \right\| \right) \\
& + \frac{e\lambda_i^\alpha}{\Gamma(\alpha+1)} \omega_i \|u_i - v_i\| + \frac{e\lambda_i^{\alpha-\beta}}{\Gamma(\alpha+1)} \omega_i \left\| {}^H D_{1+}^\beta u_i - {}^H D_{1+}^\beta v_i \right\| \\
& \leq \frac{2e(\lambda_i^\alpha + \lambda_i^{\alpha-\beta})}{\Gamma(\alpha+1)} \omega_i \left( \|u_i - v_i\| + \left\| {}^H D_{1+}^\beta u_i - {}^H D_{1+}^\beta v_i \right\| \right) \\
& + e \sum_{j=1}^k \frac{\lambda_j^\alpha + \lambda_j^{\alpha-\beta}}{\Gamma(\alpha)} \omega_j \left( \|u_j - v_j\| + \left\| {}^H D_{1+}^\beta u_j - {}^H D_{1+}^\beta v_j \right\| \right) \\
& + e \sum_{\substack{j=1 \\ j \neq i}}^k \frac{\lambda_j^\alpha + \lambda_j^{\alpha-\beta}}{\Gamma(\alpha+1)} \omega_j \left( \|u_j - v_j\| + \left\| {}^H D_{1+}^\beta u_j - {}^H D_{1+}^\beta v_j \right\| \right) \\
& = e \left( \frac{1}{\Gamma(\alpha)} + \frac{2}{\Gamma(\alpha+1)} \right) (\lambda_i^\alpha + \lambda_i^{\alpha-\beta}) \omega_i \left( \|u_i - v_i\| + \left\| {}^H D_{1+}^\beta u_i - {}^H D_{1+}^\beta v_i \right\| \right) \\
& + e \left( \frac{1}{\Gamma(\alpha)} + \frac{1}{\Gamma(\alpha+1)} \right) \sum_{\substack{j=1 \\ j \neq i}}^k (\lambda_j^\alpha + \lambda_j^{\alpha-\beta}) \omega_j \left( \|u_j - v_j\| + \left\| {}^H D_{1+}^\beta u_j - {}^H D_{1+}^\beta v_j \right\| \right).
\end{aligned}$$

Hence,

$$\begin{aligned}
& \|T_i u(t) - T_i v(t)\| \\
& \leq e \left( \frac{1}{\Gamma(\alpha)} + \frac{2}{\Gamma(\alpha+1)} \right) (\lambda_i^\alpha + \lambda_i^{\alpha-\beta}) \omega_i \left( \|u_i - v_i\| + \left\| {}^H D_{1+}^\beta u_i - {}^H D_{1+}^\beta v_i \right\| \right) \\
& + e \left( \frac{1}{\Gamma(\alpha)} + \frac{1}{\Gamma(\alpha+1)} \right) \sum_{\substack{j=1 \\ j \neq i}}^k (\lambda_j^\alpha + \lambda_j^{\alpha-\beta}) \\
& \quad \times \omega_j \left( \|u_j - v_j\| + \left\| {}^H D_{1+}^\beta u_j - {}^H D_{1+}^\beta v_j \right\| \right). \tag{3.4}
\end{aligned}$$

By the formula in reference [10]

$${}^H D_{1+}^\beta \left( \log \frac{t}{s} \right)^{\beta-1} = \frac{\Gamma(\beta)}{\Gamma(\beta-\alpha)} \left( \log \frac{t}{s} \right)^{\beta-\alpha-1}, \quad \beta > 1,$$

we have

$$\begin{aligned}
& |{}^H D_{1+}^\beta T_i u(t) - {}^H D_{1+}^\beta T_i v(t)| \\
& \leq \frac{\lambda_i^\alpha}{\Gamma(\alpha-\beta)} \int_1^t \left( \log \frac{t}{s} \right)^{\alpha-\beta-1} \left| g_i(s, u_i(s), {}^H D_{1+}^\beta u_i(s)) - g_i(s, v_i(s), {}^H D_{1+}^\beta v_i(s)) \right| ds \\
& + \frac{(\log t)^{1-\beta}}{\Gamma(\alpha-1)\Gamma(2-\beta)} \sum_{j=1}^k \left( \frac{\lambda_j^{-1}}{\sum_{j=1}^k \lambda_j^{-1}} \right) \lambda_j^\alpha \int_1^e \left( \log \frac{e}{s} \right)^{\alpha-2} \\
& \quad \times \left[ \left| g_j(s, u_j(s), {}^H D_{1+}^\beta u_j(s)) - g_j(s, v_j(s), {}^H D_{1+}^\beta v_j(s)) \right| ds \right]
\end{aligned}$$

$$\begin{aligned}
& + \frac{(\log t)^{1-\beta}}{\Gamma(\alpha)\Gamma(2-\beta)} \sum_{\substack{j=1 \\ j \neq i}}^k \left( \frac{\lambda_j^{-1}}{\sum_{j=1}^k \lambda_j^{-1}} \right) \lambda_j^\alpha \int_1^e \left( \log \frac{e}{s} \right)^{\alpha-1} \left[ \left| g_j(s, u_j(s), {}^H D_{1+}^\beta u_j(s)) \right. \right. \\
& \quad \left. \left. - g_j(s, v_j(s), {}^H D_{1+}^\beta v_j(s)) \right| ds \right] \\
& + \frac{(\log t)^{1-\beta}}{\Gamma(\alpha)\Gamma(2-\beta)} \sum_{\substack{j=1 \\ j \neq i}}^k \left( \frac{\lambda_j^{-1}}{\sum_{j=1}^k \lambda_j^{-1}} \right) \lambda_i^\alpha \int_1^e \left( \log \frac{e}{s} \right)^{\alpha-1} \left[ \left| g_i(s, u_i(s), {}^H D_{1+}^\beta u_i(s)) \right. \right. \\
& \quad \left. \left. - g_i(s, v_i(s), {}^H D_{1+}^\beta v_i(s)) \right| ds \right].
\end{aligned}$$

Using (H1) and (H2),  $\Gamma(2-\beta) \leq 1$  and  $\left( \frac{\lambda_j^{-1}}{\sum_{j=1}^k \lambda_j^{-1}} \right) < 1$  for  $j = 1, 2, \dots, k$ , we obtain

$$\begin{aligned}
& |{}^H D_{1+}^\beta T_i u(t) - {}^H D_{1+}^\beta T_i v(t)| \\
& \leq \frac{e\lambda_i^\alpha}{\Gamma(\alpha-\beta+1)} \omega_i \|u_i - v_i\| + \frac{e\lambda_i^{\alpha-\beta}}{\Gamma(\alpha-\beta+1)} \omega_i \left\| {}^H D_{1+}^\beta u_i - {}^H D_{1+}^\beta v_i \right\| \\
& \quad + e \sum_{j=1}^k \left( \frac{\lambda_i^\alpha}{\Gamma(\alpha)\Gamma(2-\beta)} \omega_j \|u_j - v_j\| + \frac{\lambda_i^{\alpha-\beta}}{\Gamma(\alpha)\Gamma(2-\beta)} \omega_j \left\| {}^H D_{1+}^\beta u_j - {}^H D_{1+}^\beta v_j \right\| \right) \\
& \quad + e \sum_{\substack{j=1 \\ j \neq i}}^k \left( \frac{\lambda_j^\alpha}{\Gamma(\alpha+1)} \omega_j \|u_j - v_j\| + \frac{\lambda_j^{\alpha-\beta}}{\Gamma(\alpha+1)} \omega_j \left\| {}^H D_{1+}^\beta u_j - {}^H D_{1+}^\beta v_j \right\| \right) \\
& \quad + \frac{e\lambda_i^\alpha}{\Gamma(\alpha+1)\Gamma(2-\beta)} \omega_i \|u_i - v_i\| + \frac{e\lambda_i^{\alpha-\beta}}{\Gamma(\alpha+1)\Gamma(2-\beta)} \omega_i \left\| {}^H D_{1+}^\beta u_i - {}^H D_{1+}^\beta v_i \right\| \\
& \leq \frac{e(\lambda_i^\alpha + \lambda_i^{\alpha-\beta})}{\Gamma(\alpha-\beta+1)} \omega_i \left( \|u_i - v_i\| + \left\| {}^H D_{1+}^\beta u_i - {}^H D_{1+}^\beta v_i \right\| \right) \\
& \quad + e \sum_{j=1}^k \frac{\lambda_j^\alpha + \lambda_j^{\alpha-\beta}}{\Gamma(\alpha)\Gamma(2-\beta)} \omega_j \left( \|u_j - v_j\| + \left\| {}^H D_{1+}^\beta u_j - {}^H D_{1+}^\beta v_j \right\| \right) \\
& \quad + e \sum_{\substack{j=1 \\ j \neq i}}^k \frac{\lambda_j^\alpha + \lambda_j^{\alpha-\beta}}{\Gamma(\alpha+1)\Gamma(2-\beta)} \omega_j \left( \|u_j - v_j\| + \left\| {}^H D_{1+}^\beta u_j - {}^H D_{1+}^\beta v_j \right\| \right) \\
& \quad + \frac{e(\lambda_i^\alpha + \lambda_i^{\alpha-\beta})}{\Gamma(\alpha+1)\Gamma(2-\beta)} \omega_i \left( \|u_i - v_i\| + \left\| {}^H D_{1+}^\beta u_i - {}^H D_{1+}^\beta v_i \right\| \right).
\end{aligned}$$

Hence,

$$\begin{aligned}
& \|{}^H D_{1+}^\beta T_i u(t) - {}^H D_{1+}^\beta T_i v(t)\| \\
& \leq e \left( \frac{1}{\Gamma(\alpha-\beta+1)} + \frac{1}{\Gamma(\alpha)\Gamma(2-\beta)} + \frac{1}{\Gamma(\alpha+1)\Gamma(2-\beta)} \right) \\
& \quad \times (\lambda_i^\alpha + \lambda_i^{\alpha-\beta}) \omega_i \left( \|u_i - v_i\| + \left\| {}^H D_{1+}^\beta u_i - {}^H D_{1+}^\beta v_i \right\| \right)
\end{aligned}$$

$$\begin{aligned}
& + e \left( \frac{1}{\Gamma(\alpha)\Gamma(2-\beta)} + \frac{1}{\Gamma(\alpha+1)\Gamma(2-\beta)} \right) \sum_{\substack{j=1 \\ j \neq i}}^k (\lambda_j^\alpha + \lambda_j^{\alpha-\beta}) \\
& \times \omega_j \left( \|u_j - v_j\| + \left\| {}^H D_{1+}^\beta u_j - {}^H D_{1+}^\beta v_j \right\| \right). \tag{3.5}
\end{aligned}$$

From (3.4) and (3.5), we have

$$\begin{aligned}
& \|T_i u(t) - T_i v(t)\| + \left\| {}^H D_{1+}^\beta T_i u(t) - {}^H D_{1+}^\beta T_i v(t) \right\| \\
& \leq e \left( \frac{1}{\Gamma(\alpha)} + \frac{2}{\Gamma(\alpha+1)} + \frac{1}{\Gamma(\alpha-\beta+1)} + \frac{1}{\Gamma(\alpha)\Gamma(2-\beta)} + \frac{1}{\Gamma(\alpha+1)\Gamma(2-\beta)} \right) \\
& \quad \times (\lambda_i^\alpha + \lambda_i^{\alpha-\beta}) \omega_i \left( \|u_i - v_i\| + \left\| {}^H D_{1+}^\beta u_i - {}^H D_{1+}^\beta v_i \right\| \right) \\
& \quad + \left( \frac{1}{\Gamma(\alpha)} + \frac{1}{\Gamma(\alpha+1)} + \frac{1}{\Gamma(\alpha)\Gamma(2-\beta)} + \frac{1}{\Gamma(\alpha+1)\Gamma(2-\beta)} \right) \sum_{\substack{j=1 \\ j \neq i}}^k (\lambda_j^\alpha + \lambda_j^{\alpha-\beta}) \\
& \quad \times \omega_j \left( \|u_j - v_j\| + \left\| {}^H D_{1+}^\beta u_j - {}^H D_{1+}^\beta v_j \right\| \right).
\end{aligned}$$

Hence,

$$\begin{aligned}
& \|T_i u(t) - T_i v(t)\|_X \\
& \leq e \left[ \left( \frac{1}{\Gamma(\alpha)} + \frac{2}{\Gamma(\alpha+1)} + \frac{1}{\Gamma(\alpha-\beta+1)} + \frac{1}{\Gamma(\alpha)\Gamma(2-\beta)} + \frac{1}{\Gamma(\alpha+1)\Gamma(2-\beta)} \right) \right. \\
& \quad \times (\lambda_i^\alpha + \lambda_i^{\alpha-\beta}) + e \left( \frac{1}{\Gamma(\alpha)} + \frac{1}{\Gamma(\alpha+1)} + \frac{1}{\Gamma(\alpha)\Gamma(2-\beta)} + \frac{1}{\Gamma(\alpha+1)\Gamma(2-\beta)} \right) \\
& \quad \times \sum_{\substack{j=1 \\ j \neq i}}^k (\lambda_j^\alpha + \lambda_j^{\alpha-\beta}) \left. \right] \left( \sum_{i=1}^k \omega_i \right) \left( \|u_j - v_j\| + \left\| {}^H D_{1+}^\beta u_j - {}^H D_{1+}^\beta v_j \right\| \right) \\
& = B_i \left( \sum_{i=1}^k \omega_i \right) \|u - v\|_{X^k}, \tag{3.6}
\end{aligned}$$

where  $B_i$ ,  $i = 1, 2, \dots, k$  are given by (3.2).

From the above equation (3.6), it follows that

$$\begin{aligned}
\|T_u - T_v\|_{X^k} &= \sum_{i=1}^k \|T_i u - T_i v\|_X \\
&\leq \left( \sum_{i=1}^k B_i \right) \left( \sum_{i=1}^k \omega_i \right) \|u - v\|_{X^k}.
\end{aligned}$$

Since

$$\left( \sum_{i=1}^k B_i \right) \left( \sum_{i=1}^k \omega_i \right) < 1,$$

we obtain that  $T$  is a contraction map. According to Banach's contraction principle, the original system (1.1) has a unique solution on  $[1, e]$ .  $\square$

**Theorem 3.2.** Assume that (H1) and (H2) hold, then system (2.1) has at least one solution on  $[1, e]$ .

**Proof.** We demonstrate that the operator  $T : X^k \rightarrow X^k$  is completely continuous. In view of continuity of the functions  $f_i, i = 1, 2, \dots, k$ , we obtain that the operator  $T$  is continuous.

Let  $\Omega$  be any bounded subset of  $X^k$ , for  $u = (u_1, u_2, \dots, u_k) \in \Omega$ ,  $t \in [1, e]$ , we can get

$$\begin{aligned} |T_i u(t)| &\leq \frac{\lambda_i^\alpha}{\Gamma(\alpha)} \int_1^t \left( \log \frac{t}{s} \right)^{\alpha-1} \left| g_i(s, u_i(s), {}^H D_{1+}^\beta u_i(s)) \right| ds \\ &\quad + \frac{\log t}{\Gamma(\alpha-1)} \sum_{j=1}^k \left( \frac{\lambda_j^{-1}}{\sum_{j=1}^k \lambda_j^{-1}} \right) \lambda_j^\alpha \int_1^e \left( \log \frac{e}{s} \right)^{\alpha-2} \\ &\quad \times \left| g_j(s, u_j(s), {}^H D_{1+}^\beta u_j(s)) \right| ds \\ &\quad + \frac{\log t}{\Gamma(\alpha)} \sum_{\substack{j=1 \\ j \neq i}}^k \left( \frac{\lambda_j^{-1}}{\sum_{j=1}^k \lambda_j^{-1}} \right) \lambda_j^\alpha \int_1^e \left( \log \frac{e}{s} \right)^{\alpha-1} \left| g_j(s, u_j(s), {}^H D_{1+}^\beta u_j(s)) \right| ds \\ &\quad + \frac{\log t}{\Gamma(\alpha)} \sum_{\substack{j=1 \\ j \neq i}}^k \left( \frac{\lambda_j^{-1}}{\sum_{j=1}^k \lambda_j^{-1}} \right) \lambda_i^\alpha \int_1^e \left( \log \frac{e}{s} \right)^{\alpha-1} \left| g_i(s, u_i(s), {}^H D_{1+}^\beta u_i(s)) \right| ds. \end{aligned}$$

Using (H3), we can write

$$\begin{aligned} |T_i u(t)| &\leq \frac{2eA_i \lambda_i^\alpha}{\Gamma(\alpha+1)} + \sum_{j=1}^k \frac{eA_j \lambda_j^\alpha}{\Gamma(\alpha)} + \sum_{\substack{j=1 \\ j \neq i}}^k \frac{eA_j \lambda_j^\alpha}{\Gamma(\alpha+1)} \\ &= eA_i \lambda_i^\alpha \left( \frac{1}{\Gamma(\alpha)} + \frac{2}{\Gamma(\alpha+1)} \right) + e \sum_{\substack{j=1 \\ j \neq i}}^k A_j \lambda_j^\alpha \left( \frac{1}{\Gamma(\alpha)} + \frac{1}{\Gamma(\alpha+1)} \right). \end{aligned}$$

Thus,

$$\|T_i u\| \leq eA_i \lambda_i^\alpha \left( \frac{1}{\Gamma(\alpha)} + \frac{2}{\Gamma(\alpha+1)} \right) + e \sum_{\substack{j=1 \\ j \neq i}}^k A_j \lambda_j^\alpha \left( \frac{1}{\Gamma(\alpha)} + \frac{1}{\Gamma(\alpha+1)} \right). \quad (3.7)$$

On the other hand,

$$\begin{aligned} &\left| {}^H D_{1+}^\beta T_i u(t) \right| \\ &\leq \frac{\lambda_i^\alpha}{\Gamma(\alpha-\beta)} \int_1^t \left( \log \frac{t}{s} \right)^{\alpha-\beta-1} \left| g_i(s, u_i(s), {}^H D_{1+}^\beta u_i(s)) \right| ds \\ &\quad + \frac{(\log t)^{1-\beta}}{\Gamma(\alpha-1)\Gamma(2-\beta)} \sum_{j=1}^k \left( \frac{\lambda_j^{-1}}{\sum_{j=1}^k \lambda_j^{-1}} \right) \lambda_j^\alpha \int_1^e \left( \log \frac{e}{s} \right)^{\alpha-2} \\ &\quad \times \left| g_j(s, u_j(s), {}^H D_{1+}^\beta u_j(s)) \right| ds \end{aligned}$$

$$\begin{aligned}
& + \frac{(\log t)^{1-\beta}}{\Gamma(\alpha)\Gamma(2-\beta)} \sum_{\substack{j=1 \\ j \neq i}}^k \left( \frac{\lambda_j^{-1}}{\sum_{j=1}^k \lambda_j^{-1}} \right) \lambda_j^\alpha \int_1^e \left( \log \frac{e}{s} \right)^{\alpha-1} \left| g_j(s, u_j(s), {}^H D_{1+}^\beta u_j(s)) \right| ds \\
& + \frac{(\log t)^{1-\beta}}{\Gamma(\alpha)\Gamma(2-\beta)} \sum_{\substack{j=1 \\ j \neq i}}^k \left( \frac{\lambda_j^{-1}}{\sum_{j=1}^k \lambda_j^{-1}} \right) \lambda_i^\alpha \int_1^e \left( \log \frac{e}{s} \right)^{\alpha-1} \left| g_i(s, u_i(s), {}^H D_{1+}^\beta u_i(s)) \right| ds.
\end{aligned}$$

By (H3),  $(\log t)^{1-\beta} \leq 1$  and  $\left( \frac{\lambda_j^{-1}}{\sum_{j=1}^k \lambda_j^{-1}} \right) < 1$  for  $j = 1, 2, \dots, k$ , we get

$$\begin{aligned}
& \left| {}^H D_{1+}^\beta T_i u(t) \right| \\
& \leq \frac{e A_i \lambda_i^\alpha}{\Gamma(\alpha - \beta + 1)} + \frac{e A_i \lambda_i^\alpha}{\Gamma(\alpha + 1)\Gamma(2 - \beta)} + \sum_{j=1}^k \frac{e A_j \lambda_j^\alpha}{\Gamma(\alpha)\Gamma(2 - \beta)} + \sum_{\substack{j=1 \\ j \neq i}}^k \frac{e A_j \lambda_j^\alpha}{\Gamma(\alpha + 1)\Gamma(2 - \beta)} \\
& = e A_i \lambda_i^\alpha \left( \frac{1}{\Gamma(\alpha - \beta + 1)} + \frac{1}{\Gamma(\alpha)\Gamma(2 - \beta)} + \frac{1}{\Gamma(\alpha + 1)\Gamma(2 - \beta)} \right) \\
& \quad + e \sum_{\substack{j=1 \\ j \neq i}}^k A_j \lambda_j^\alpha \left( \frac{1}{\Gamma(\alpha)\Gamma(2 - \beta)} + \frac{1}{\Gamma(\alpha + 1)\Gamma(2 - \beta)} \right).
\end{aligned}$$

Hence

$$\begin{aligned}
\left\| {}^H D_{1+}^\beta T_i u \right\| & \leq e A_i \lambda_i^\alpha \left( \frac{1}{\Gamma(\alpha - \beta + 1)} + \frac{1}{\Gamma(\alpha)\Gamma(2 - \beta)} + \frac{1}{\Gamma(\alpha + 1)\Gamma(2 - \beta)} \right) \\
& \quad + e \sum_{\substack{j=1 \\ j \neq i}}^k A_j \lambda_j^\alpha \left( \frac{1}{\Gamma(\alpha)\Gamma(2 - \beta)} + \frac{1}{\Gamma(\alpha + 1)\Gamma(2 - \beta)} \right). \quad (3.8)
\end{aligned}$$

From (3.7) and (3.8), we have

$$\begin{aligned}
& \|T_i u\| + \left\| {}^H D_{1+}^\beta T_i u \right\| \\
& \leq e A_i \lambda_i^\alpha \left( \frac{1}{\Gamma(\alpha)} + \frac{2}{\Gamma(\alpha + 1)} + \frac{1}{\Gamma(\alpha - \beta + 1)} + \frac{1}{\Gamma(\alpha)\Gamma(2 - \beta)} + \frac{1}{\Gamma(\alpha + 1)\Gamma(2 - \beta)} \right) \\
& \quad + e \sum_{\substack{j=1 \\ j \neq i}}^k A_i \lambda_i^\alpha \left( \frac{1}{\Gamma(\alpha)} + \frac{1}{\Gamma(\alpha + 1)} + \frac{1}{\Gamma(\alpha)\Gamma(2 - \beta)} + \frac{1}{\Gamma(\alpha + 1)\Gamma(2 - \beta)} \right) \\
& \leq \left[ e \lambda_i^\alpha \left( \frac{1}{\Gamma(\alpha)} + \frac{2}{\Gamma(\alpha + 1)} + \frac{1}{\Gamma(\alpha - \beta + 1)} + \frac{1}{\Gamma(\alpha)\Gamma(2 - \beta)} + \frac{1}{\Gamma(\alpha + 1)\Gamma(2 - \beta)} \right) \right. \\
& \quad \left. + e \sum_{\substack{j=1 \\ j \neq i}}^k \lambda_j^\alpha \left( \frac{1}{\Gamma(\alpha)} + \frac{1}{\Gamma(\alpha + 1)} + \frac{1}{\Gamma(\alpha)\Gamma(2 - \beta)} + \frac{1}{\Gamma(\alpha + 1)\Gamma(2 - \beta)} \right) \right] \left( \sum_{j=1}^k A_j \right) \\
& = C_i \left( \sum_{j=1}^k A_j \right),
\end{aligned}$$

where  $C_i$ ,  $i = 1, 2, \dots, k$  are given by (3.3). Hence,

$$\|Tu\|_{X^k} = \sum_{i=1}^k \|T_i u\|_X \leq \left( \sum_{i=1}^k C_i \right) \left( \sum_{j=1}^k A_j \right) < \infty, \quad (3.9)$$

so it follows that  $T$  is uniformly bounded.

Now we will prove that  $T$  is equi-continuous. For  $u = (u_1, u_2, \dots, u_k) \in \Omega$ ,  $t_1, t_2 \in [1, e]$  with  $t_1 < t_2$ , we can get

$$\begin{aligned} & |T_i u(t_2) - T_i u(t_1)| \\ & \leq \frac{\lambda_i^\alpha}{\Gamma(\alpha)} \int_1^{t_1} \left( \left( \log \frac{t_2}{s} \right)^{\alpha-1} - \left( \log \frac{t_1}{s} \right)^{\alpha-1} \right) \left| g_i(s, u_i(s), {}^H D_{1+}^\beta u_i(s)) \right| ds \\ & \quad + \frac{\lambda_i^\alpha}{\Gamma(\alpha)} \int_{t_1}^{t_2} \left( \log \frac{t_2}{s} \right)^{\alpha-1} \left| g_i(s, u_i(s), {}^H D_{1+}^\beta u_i(s)) \right| ds \\ & \quad + \frac{(\log t_1 - \log t_2)}{\Gamma(\alpha-1)} \sum_{j=1}^k \left( \frac{\lambda_j^{-1}}{\sum_{j=1}^k \lambda_j^{-1}} \right) \lambda_j^\alpha \\ & \quad \times \int_1^e \left( \log \frac{e}{s} \right)^{\alpha-2} \left| g_j(s, u_j(s), {}^H D_{1+}^\beta u_j(s)) \right| ds \\ & \quad + \frac{(\log t_2 - \log t_1)}{\Gamma(\alpha)} \sum_{\substack{j=1 \\ j \neq i}}^k \left( \frac{\lambda_j^{-1}}{\sum_{j=1}^k \lambda_j^{-1}} \right) \lambda_j^\alpha \\ & \quad \times \int_1^e \left( \log \frac{e}{s} \right)^{\alpha-1} \left| g_j(s, u_j(s), {}^H D_{1+}^\beta u_j(s)) \right| ds \\ & \quad + \frac{(\log t_1 - \log t_2)}{\Gamma(\alpha)} \sum_{\substack{j=1 \\ j \neq i}}^k \left( \frac{\lambda_j^{-1}}{\sum_{j=1}^k \lambda_j^{-1}} \right) \lambda_i^\alpha \\ & \quad \times \int_1^e \left( \log \frac{e}{s} \right)^{\alpha-1} \left| g_i(s, u_i(s), {}^H D_{1+}^\beta u_i(s)) \right| ds. \end{aligned}$$

Using (H3), we get

$$\begin{aligned} & |T_i u(t_2) - T_i u(t_1)| \\ & \leq \frac{e A_i \lambda_i^\alpha}{\Gamma(\alpha+1)} ((\log t_2)^\alpha - (\log t_1)^\alpha) + \frac{e A_i \lambda_i^\alpha}{\Gamma(\alpha+1)} (\log t_1 - \log t_2) \\ & \quad + e \sum_{j=1}^k \frac{A_j \lambda_j^\alpha}{\Gamma(\alpha)} (\log t_1 - \log t_2) + e \sum_{\substack{j=1 \\ j \neq i}}^k \frac{A_j \lambda_j^\alpha}{\Gamma(\alpha+1)} (\log t_2 - \log t_1) \\ & = e A_i \lambda_i^\alpha \left( \frac{1}{\Gamma(\alpha)} + \frac{1}{\Gamma(\alpha+1)} \right) (\log t_1 - \log t_2) + \frac{e A_i \lambda_i^\alpha}{\Gamma(\alpha+1)} ((\log t_2)^\alpha - (\log t_1)^\alpha) \\ & \quad + e \sum_{\substack{j=1 \\ j \neq i}}^k A_j \lambda_j^\alpha \left( \frac{1}{\Gamma(\alpha)} + \frac{1}{\Gamma(\alpha+1)} \right) (\log t_2 - \log t_1). \quad (3.10) \end{aligned}$$

In addition,

$$\left| {}^H D_{1+}^\beta T_i u(t_2) - {}^H D_{1+}^\beta T_i u(t_1) \right|$$

$$\begin{aligned}
&\leq \frac{\lambda_i^\alpha}{\Gamma(\alpha - \beta)} \int_1^{t_1} \left( \left( \log \frac{t_2}{s} \right)^{\alpha - \beta - 1} - \left( \log \frac{t_1}{s} \right)^{\alpha - \beta - 1} \right) \left| g_i(s, u_i(s), {}^H D_{1+}^\beta u_i(s)) \right| ds \\
&\quad + \frac{\lambda_i^\alpha}{\Gamma(\alpha - \beta)} \int_{t_1}^{t_2} \left( \log \frac{t_2}{s} \right)^{\alpha - \beta - 1} \left| g_i(s, u_i(s), {}^H D_{1+}^\beta u_i(s)) \right| ds \\
&\quad + \frac{(\log t_1^{1-\beta} - \log t_2^{1-\beta})}{\Gamma(\alpha - 1)\Gamma(2 - \beta)} \sum_{j=1}^k \left( \frac{\lambda_j^{-1}}{\sum_{j=1}^k \lambda_j^{-1}} \right) \lambda_j^\alpha \\
&\quad \times \int_1^e \left( \log \frac{e}{s} \right)^{\alpha - 2} \left| g_j(s, u_j(s), {}^H D_{1+}^\beta u_j(s)) \right| ds \\
&\quad + \frac{(\log t_2^{1-\beta} - \log t_1^{1-\beta})}{\Gamma(\alpha)\Gamma(2 - \beta)} \sum_{\substack{j=1 \\ j \neq i}}^k \left( \frac{\lambda_j^{-1}}{\sum_{j=1}^k \lambda_j^{-1}} \right) \lambda_j^\alpha \\
&\quad \times \int_1^e \left( \log \frac{e}{s} \right)^{\alpha - 1} \left| g_j(s, u_j(s), {}^H D_{1+}^\beta u_j(s)) \right| ds \\
&\quad + \frac{(\log t_1^{1-\beta} - \log t_2^{1-\beta})}{\Gamma(\alpha)\Gamma(2 - \beta)} \sum_{\substack{j=1 \\ j \neq i}}^k \left( \frac{\lambda_j^{-1}}{\sum_{j=1}^k \lambda_j^{-1}} \right) \lambda_i^\alpha \\
&\quad \times \int_1^e \left( \log \frac{e}{s} \right)^{\alpha - 1} \left| g_i(s, u_i(s), {}^H D_{1+}^\beta u_i(s)) \right| ds.
\end{aligned}$$

By (H3), we get

$$\begin{aligned}
&| {}^H D_{1+}^\beta T_i u(t_2) - {}^H D_{1+}^\beta T_i u(t_1) | \\
&\leq \frac{e A_i \lambda_i^\alpha}{\Gamma(\alpha - \beta + 1)} ((\log t_2)^{\alpha - \beta} - (\log t_1)^{\alpha - \beta}) \\
&\quad + \frac{e A_i \lambda_i^\alpha}{\Gamma(\alpha + 1)\Gamma(2 - \beta)} ((\log t_1)^{1 - \beta} - (\log t_2)^{1 - \beta}) \\
&\quad + e \sum_{j=1}^k \frac{A_j \lambda_j^\alpha}{\Gamma(\alpha + 1)\Gamma(2 - \beta)} ((\log t_1)^{1 - \beta} - (\log t_2)^{1 - \beta}) \\
&\quad + e \sum_{\substack{j=1 \\ j \neq i}}^k \frac{A_j \lambda_j^\alpha}{\Gamma(\alpha + 1)\Gamma(2 - \beta)} ((\log t_2)^{1 - \beta} - (\log t_1)^{1 - \beta}) \\
&= \frac{e A_i \lambda_i^\alpha}{\Gamma(\alpha - \beta + 1)} ((\log t_2)^{\alpha - \beta} - (\log t_1)^{\alpha - \beta}) \\
&\quad + e A_i \lambda_i^\alpha \left( \frac{1}{\Gamma(\alpha)\Gamma(2 - \beta)} + \frac{1}{\Gamma(\alpha + 1)\Gamma(2 - \beta)} \right) (\log t_1^{1 - \beta} - \log t_2^{1 - \beta}) \\
&\quad + e \sum_{\substack{j=1 \\ j \neq i}}^k A_j \lambda_j^\alpha \left( \frac{1}{\Gamma(\alpha)\Gamma(2 - \beta)} + \frac{1}{\Gamma(\alpha + 1)\Gamma(2 - \beta)} \right) (\log t_2^{1 - \beta} - \log t_1^{1 - \beta}). \quad (3.11)
\end{aligned}$$

Hence, from (3.10) and (3.11), we obtain

$$\begin{aligned}
&\|T_i u(t_2) - T_i u(t_1)\|_X \\
&\leq \frac{e A_i \lambda_i^\alpha}{\Gamma(\alpha - \beta + 1)} ((\log t_2)^{\alpha - \beta} - (\log t_1)^{\alpha - \beta})
\end{aligned}$$

$$\begin{aligned}
& + e \sum_{\substack{j=1 \\ j \neq i}}^k A_j \lambda_j^\alpha \left( \frac{1}{\Gamma(\alpha)} + \frac{1}{\Gamma(\alpha+1)} \right) (\log t_2 - \log t_1) \\
& + e A_i \lambda_i^\alpha \left( \frac{1}{\Gamma(\alpha)\Gamma(2-\beta)} + \frac{1}{\Gamma(\alpha+1)\Gamma(2-\beta)} \right) (\log t_1^{1-\beta} - \log t_2^{1-\beta}) \\
& + e \sum_{\substack{j=1 \\ j \neq i}}^k A_j \lambda_j^\alpha \left( \frac{1}{\Gamma(\alpha)\Gamma(2-\beta)} + \frac{1}{\Gamma(\alpha+1)\Gamma(2-\beta)} \right) (\log t_2^{1-\beta} - \log t_1^{1-\beta}) \\
& + e A_i \lambda_i^\alpha \left( \frac{1}{\Gamma(\alpha)} + \frac{1}{\Gamma(\alpha+1)} \right) (\log t_1 - \log t_2) + \frac{e A_i \lambda_i^\alpha}{\Gamma(\alpha+1)} ((\log t_2)^\alpha - (\log t_1)^\alpha),
\end{aligned}$$

which implies  $\|T_i u(t_2) - T_i u(t_1)\|_X \rightarrow 0$  as  $t_2 \rightarrow t_1$  and so  $\|T u(t_2) - T u(t_1)\|_{X^k} \rightarrow 0$  as  $t_2 \rightarrow t_1$ . Therefore, the operator  $T$  is equi-continuous on  $X^k$  and it follows from the Arzela-Ascoli theorem that  $T$  is completely continuous.

Define  $Q = \{(u_1, u_2, \dots, u_k) \in X^k : (u_1, u_2, \dots, u_k) = \mu T(u_1, u_2, \dots, u_k), k = 1, 2, \dots, 12, 0 < \mu < 1\}$  is bounded. Let  $(u_1, u_2, \dots, u_k) \in Q$ , then  $(u_1, u_2, \dots, u_k) = \mu T(u_1, u_2, \dots, u_k)$  and for each  $t \in [1, e]$ , we have  $u_i(t) = \mu T_i(u_1, u_2, \dots, u_i)$ ,  $i = 1, 2, \dots, k$ . Hence, from (H3), we get

$$\begin{aligned}
|u_i(t)| & \leq \mu \left[ \frac{2e A_i \lambda_i^\alpha}{\Gamma(\alpha+1)} + \sum_{\substack{j=1 \\ j \neq i}}^k \frac{e A_j \lambda_j^\alpha}{\Gamma(\alpha)} + \sum_{\substack{j=1 \\ j \neq i}}^k \frac{e A_j \lambda_j^\alpha}{\Gamma(\alpha+1)} \right] \\
& = \mu \left[ e A_i \lambda_i^\alpha \left( \frac{1}{\Gamma(\alpha)} + \frac{2}{\Gamma(\alpha+1)} \right) + e \sum_{\substack{j=1 \\ j \neq i}}^k A_j \lambda_j^\alpha \left( \frac{1}{\Gamma(\alpha)} + \frac{1}{\Gamma(\alpha+1)} \right) \right].
\end{aligned}$$

Thus,

$$\|u_i\| \leq \mu \left[ e A_i \lambda_i^\alpha \left( \frac{1}{\Gamma(\alpha)} + \frac{2}{\Gamma(\alpha+1)} \right) + e \sum_{\substack{j=1 \\ j \neq i}}^k A_j \lambda_j^\alpha \left( \frac{1}{\Gamma(\alpha)} + \frac{1}{\Gamma(\alpha+1)} \right) \right] \quad (3.12)$$

and

$$\begin{aligned}
& \left| {}^H D_{1+}^\beta T_i u(t) \right| \\
& \leq \frac{e A_i \lambda_i^\alpha}{\Gamma(\alpha - \beta + 1)} + \frac{e A_i \lambda_i^\alpha}{\Gamma(\alpha + 1)\Gamma(2 - \beta)} + \sum_{j=1}^k \frac{e A_j \lambda_j^\alpha}{\Gamma(\alpha)\Gamma(2 - \beta)} + \sum_{\substack{j=1 \\ j \neq i}}^k \frac{e A_j \lambda_j^\alpha}{\Gamma(\alpha + 1)\Gamma(2 - \beta)} \\
& = e A_i \lambda_i^\alpha \left( \frac{1}{\Gamma(\alpha - \beta + 1)} + \frac{1}{\Gamma(\alpha)\Gamma(2 - \beta)} + \frac{1}{\Gamma(\alpha + 1)\Gamma(2 - \beta)} \right) \\
& \quad + e \sum_{\substack{j=1 \\ j \neq i}}^k A_j \lambda_j^\alpha \left( \frac{1}{\Gamma(\alpha)\Gamma(2 - \beta)} + \frac{1}{\Gamma(\alpha + 1)\Gamma(2 - \beta)} \right),
\end{aligned}$$

which gives

$$\|{}^H D_{1+}^\beta u_i\| \leq \mu \left[ e A_i \lambda_i^\alpha \left( \frac{1}{\Gamma(\alpha - \beta + 1)} + \frac{1}{\Gamma(\alpha)\Gamma(2 - \beta)} + \frac{1}{\Gamma(\alpha + 1)\Gamma(2 - \beta)} \right) \right]$$



$$+e \sum_{\substack{j=1 \\ j \neq i}}^k A_j \lambda_j^\alpha \left( \frac{1}{\Gamma(\alpha)\Gamma(2-\beta)} + \frac{1}{\Gamma(\alpha+1)\Gamma(2-\beta)} \right) \Bigg]. \quad (3.13)$$

From (3.12) and (3.13), we obtain

$$\begin{aligned} & \|u_i\| + \left\| {}^H D_{1+}^\beta u_i \right\| \\ & \leq \mu \left[ e \lambda_i^\alpha \left( \frac{1}{\Gamma(\alpha)} + \frac{2}{\Gamma(\alpha+1)} + \frac{1}{\Gamma(\alpha-\beta+1)} + \frac{1}{\Gamma(\alpha)\Gamma(2-\beta)} + \frac{1}{\Gamma(\alpha+1)\Gamma(2-\beta)} \right) \right. \\ & \quad \left. + e \sum_{\substack{j=1 \\ j \neq i}}^k \lambda_j^\alpha \left[ \frac{1}{\Gamma(\alpha)} + \frac{1}{\Gamma(\alpha+1)} + \frac{1}{\Gamma(\alpha)\Gamma(2-\beta)} + \frac{1}{\Gamma(\alpha+1)\Gamma(2-\beta)} \right] \left( \sum_{j=1}^k A_j \right) \right] \\ & = \mu C_i \left( \sum_{j=1}^k A_j \right), \end{aligned} \quad (3.14)$$

where  $C_i, i = 1, 2, \dots, k$  are given by (3.3). Hence,

$$\|u\|_{X^k} = \sum_{i=1}^k \|u_i\|_X \leq \mu \left( \sum_{i=1}^k C_i \right) \left( \sum_{j=1}^k A_j \right) < \infty. \quad (3.15)$$

This indicates that the set  $Q$  is bounded. Thus, by Lemma 2.2, the operator  $T$  has at least one fixed point, which denotes that the original system (1.1) has at least one solution on  $[1, e]$ .  $\square$

## 4. Hyers-Ulam stability

Let  $\varepsilon_i > 0$ . Consider the following inequality

$$\left| {}^H D_{1+}^\alpha u_i(t) - \lambda_i^\alpha f_i(t, u_i(t), {}^H D_{1+}^\beta u_i(t)) \right| \leq \varepsilon_i, \quad t \in [1, e]. \quad (4.1)$$

**Definition 4.1.** [25] The fractional differential system (1.1) is called Ulam-Hyers stable, if there is a constant  $c_{f_1, f_2, \dots, f_k} > 0$  such that for each  $\varepsilon = \varepsilon(\varepsilon_1, \varepsilon_2, \dots, \varepsilon_k) > 0$  and for each solution  $u = (u_1, u_2, \dots, u_k) \in X^k$  of the inequality (4.1), there exists a solution  $\bar{u} = (\bar{u}_1, \bar{u}_2, \dots, \bar{u}_k) \in X^k$  of (1.1) with

$$\|u - \bar{u}\|_X \leq c_{f_1, f_2, \dots, f_k} \varepsilon, \quad t \in [1, e].$$

**Definition 4.2.** [25] The fractional differential system (1.1) is called generalized Ulam-Hyers stable, if there exists function  $\psi_{f_1, f_2, \dots, f_k} \in \mathbb{C}(\mathbb{R}^+, \mathbb{R}^+)$  with  $\psi_{f_1, f_2, \dots, f_k}(0) = 0$  such that for each  $\varepsilon = \varepsilon(\varepsilon_1, \varepsilon_2, \dots, \varepsilon_k) > 0$  and for each solution  $u = (u_1, u_2, \dots, u_k) \in X^k$  of the inequality (4.1), there exists a solution  $\bar{u} = (\bar{u}_1, \bar{u}_2, \dots, \bar{u}_k) \in X^k$  of (1.1) with

$$\|u - \bar{u}\|_X \leq \psi_{f_1, f_2, \dots, f_k}(\varepsilon), \quad t \in [1, e].$$

**Remark 4.1.** Let function  $u = (u_1, u_2, \dots, u_k) \in X^k$ ,  $k = 1, 2, \dots, 12$ , be the solution of system (4.1). If there are functions  $\varphi_i : [1, e] \rightarrow \mathbb{R}^+$  dependent on  $u_i$  respectively, then

- (i)  $|\varphi_i(t)| \leq \varepsilon_i, t \in [1, e], i = 1, 2, \dots, 12;$   
 (ii)  ${}^H D_{1+}^\alpha u_i(t) = \lambda_i^\alpha f_i(t, u_i(t), {}^H D_{1+}^\beta u_i(t)) + \varphi_i(t), t \in [1, e], i = 1, 2, \dots, 12.$

**Lemma 4.1.** Suppose  $u = (u_1, u_2, \dots, u_k) \in X^k$  is the solution of inequality (4.1). Then, the following inequality holds:

$$\begin{aligned} |u_i(t) - u_i^*(t)| &\leq \varepsilon_i e \left( \frac{1}{\Gamma(\alpha)} + \frac{2}{\Gamma(\alpha+1)} \right) + \varepsilon_j e \sum_{\substack{j=1 \\ j \neq i}}^k \left( \frac{1}{\Gamma(\alpha)} + \frac{1}{\Gamma(\alpha+1)} \right), \\ |{}^H D_{1+}^\beta u_i(t) - {}^H D_{1+}^\beta u_i^*(t)| &\leq \varepsilon_j e \sum_{\substack{j=1 \\ j \neq i}}^k \left( \frac{1}{\Gamma(\alpha)\Gamma(2-\beta)} + \frac{1}{\Gamma(\alpha+1)\Gamma(2-\beta)} \right) \\ &\quad + \varepsilon_i e \left( \frac{1}{\Gamma(\alpha-\beta+1)} + \frac{1}{\Gamma(\alpha)\Gamma(2-\beta)} + \frac{1}{\Gamma(\alpha+1)\Gamma(2-\beta)} \right), \end{aligned}$$

where

$$\begin{aligned} u_i^*(t) &= \frac{1}{\Gamma(\alpha)} \int_1^t \left( \log \frac{t}{s} \right)^{\alpha-1} z_i(s) ds \\ &\quad - \frac{\log t}{\Gamma(\alpha-1)} \sum_{j=1}^k \left( \frac{\lambda_j^{-1}}{\sum_{j=1}^k \lambda_j^{-1}} \right) \int_1^e \left( \log \frac{e}{s} \right)^{\alpha-2} z_j(s) ds \\ &\quad + \frac{\log t}{\Gamma(\alpha)} \sum_{\substack{j=1 \\ j \neq i}}^k \left( \frac{\lambda_j^{-1}}{\sum_{j=1}^k \lambda_j^{-1}} \right) \int_1^e \left( \log \frac{e}{s} \right)^{\alpha-1} z_j(s) ds \\ &\quad - \frac{\log t}{\Gamma(\alpha)} \sum_{\substack{j=1 \\ j \neq i}}^k \left( \frac{\lambda_j^{-1}}{\sum_{j=1}^k \lambda_j^{-1}} \right) \int_1^e \left( \log \frac{e}{s} \right)^{\alpha-1} z_i(s) ds, \\ {}^H D_{1+}^\beta u_i^*(t) &= \frac{1}{\Gamma(\alpha-\beta)} \int_1^t \left( \log \frac{t}{s} \right)^{\alpha-\beta-1} z_i(s) ds \\ &\quad - \frac{(\log t)^{1-\beta}}{\Gamma(\alpha-1)\Gamma(2-\beta)} \sum_{j=1}^k \left( \frac{\lambda_j^{-1}}{\sum_{j=1}^k \lambda_j^{-1}} \right) \int_1^e \left( \log \frac{e}{s} \right)^{\alpha-2} z_j(s) ds \\ &\quad + \frac{(\log t)^{1-\beta}}{\Gamma(\alpha)\Gamma(2-\beta)} \sum_{\substack{j=1 \\ j \neq i}}^k \left( \frac{\lambda_j^{-1}}{\sum_{j=1}^k \lambda_j^{-1}} \right) \int_1^e \left( \log \frac{e}{s} \right)^{\alpha-1} z_j(s) ds \\ &\quad - \frac{(\log t)^{1-\beta}}{\Gamma(\alpha)\Gamma(2-\beta)} \sum_{\substack{j=1 \\ j \neq i}}^k \left( \frac{\lambda_j^{-1}}{\sum_{j=1}^k \lambda_j^{-1}} \right) \int_1^e \left( \log \frac{e}{s} \right)^{\alpha-1} z_i(s) ds, \end{aligned}$$

and here

$$z_i(s) = \frac{h_i(s)}{s}, \quad h_i(s) = \lambda_i^\alpha f_i(s, u_i(s), {}^H D_{1+}^\beta u_i(s)), \quad i = 1, 2, \dots, 12.$$

**Proof.** From Remark 4.1, we have

$$\begin{cases} {}^H D_{1+}^{\alpha} u_i(t) = \lambda_i^{\alpha} f_i(s, u_i(s), {}^H D_{1+}^{\beta} u_i(s)) + \varphi_i(t), & t \in [1, e], \\ u_i(1) = 0, & i = 1, 2, \dots, 12, \\ u_i(e) = u_j(e), & i, j = 1, 2, \dots, 12, \quad i \neq j, \\ \sum_{i=1}^k \lambda_i^{-1} u'_i(e) = 0, & i = 1, 2, \dots, 12. \end{cases} \quad (4.2)$$

By Lemma 2.3, the solution of (4.2) can be given in the following form

$$\begin{aligned} u_i(t) = & \frac{1}{\Gamma(\alpha)} \int_1^t \left( \log \frac{t}{s} \right)^{\alpha-1} \left( z_i(s) + \frac{\varphi_i(s)}{s} \right) ds \\ & - \frac{\log t}{\Gamma(\alpha-1)} \sum_{j=1}^k \left( \frac{\lambda_j^{-1}}{\sum_{j=1}^k \lambda_j^{-1}} \right) \int_1^e \left( \log \frac{e}{s} \right)^{\alpha-2} \left( z_j(s) + \frac{\varphi_j(s)}{s} \right) ds \\ & + \frac{\log t}{\Gamma(\alpha)} \sum_{\substack{j=1 \\ j \neq i}}^k \left( \frac{\lambda_j^{-1}}{\sum_{j=1}^k \lambda_j^{-1}} \right) \int_1^e \left( \log \frac{e}{s} \right)^{\alpha-1} \left( z_j(s) + \frac{\varphi_j(s)}{s} \right) ds \\ & - \frac{\log t}{\Gamma(\alpha)} \sum_{\substack{j=1 \\ j \neq i}}^k \left( \frac{\lambda_j^{-1}}{\sum_{j=1}^k \lambda_j^{-1}} \right) \int_1^e \left( \log \frac{e}{s} \right)^{\alpha-1} \left( z_i(s) + \frac{\varphi_i(s)}{s} \right) ds \end{aligned}$$

and

$$\begin{aligned} & {}^H D_{1+}^{\beta} u_i(t) \\ = & \frac{1}{\Gamma(\alpha-\beta)} \int_1^t \left( \log \frac{t}{s} \right)^{\alpha-\beta-1} \left( z_i(s) + \frac{\varphi_i(s)}{s} \right) ds \\ & - \frac{(\log t)^{1-\beta}}{\Gamma(\alpha-1)\Gamma(2-\beta)} \sum_{j=1}^k \left( \frac{\lambda_j^{-1}}{\sum_{j=1}^k \lambda_j^{-1}} \right) \int_1^e \left( \log \frac{e}{s} \right)^{\alpha-2} \left( z_j(s) + \frac{\varphi_j(s)}{s} \right) ds \\ & + \frac{(\log t)^{1-\beta}}{\Gamma(\alpha)\Gamma(2-\beta)} \sum_{\substack{j=1 \\ j \neq i}}^k \left( \frac{\lambda_j^{-1}}{\sum_{j=1}^k \lambda_j^{-1}} \right) \int_1^e \left( \log \frac{e}{s} \right)^{\alpha-1} \left( z_j(s) + \frac{\varphi_j(s)}{s} \right) ds \\ & - \frac{(\log t)^{1-\beta}}{\Gamma(\alpha)\Gamma(2-\beta)} \sum_{\substack{j=1 \\ j \neq i}}^k \left( \frac{\lambda_j^{-1}}{\sum_{j=1}^k \lambda_j^{-1}} \right) \int_1^e \left( \log \frac{e}{s} \right)^{\alpha-1} \left( z_i(s) + \frac{\varphi_i(s)}{s} \right) ds. \end{aligned}$$

Then, we deduce that

$$\begin{aligned} |u_i(t) - u_i^*(t)| & \leq \varepsilon_i \frac{2e}{\Gamma(\alpha+1)} + \sum_{j=1}^k \varepsilon_j \frac{e}{\Gamma(\alpha)} + \sum_{\substack{j=1 \\ j \neq i}}^k \varepsilon_j \frac{e}{\Gamma(\alpha+1)} \\ & = \varepsilon_i e \left( \frac{1}{\Gamma(\alpha)} + \frac{2}{\Gamma(\alpha+1)} \right) + \varepsilon_j e \sum_{\substack{j=1 \\ j \neq i}}^k \left( \frac{1}{\Gamma(\alpha)} + \frac{1}{\Gamma(\alpha+1)} \right) \end{aligned}$$

and

$$\begin{aligned}
& |{}^H D_{1+}^\beta u_i(t) - {}^H D_{1+}^\beta u_i^*(t)| \\
& \leq \varepsilon_i \frac{e}{\Gamma(\alpha - \beta + 1)} + \varepsilon_j \frac{e}{\Gamma(\alpha + 1)\Gamma(2 - \beta)} \\
& \quad + \varepsilon_j \sum_{j=1}^k \frac{e}{\Gamma(\alpha)\Gamma(2 - \beta)} + \varepsilon_i \sum_{\substack{j=1 \\ j \neq i}}^k \frac{e}{\Gamma(\alpha + 1)\Gamma(2 - \beta)} \\
& = \varepsilon_j e \sum_{\substack{j=1 \\ j \neq i}}^k \left( \frac{1}{\Gamma(\alpha)\Gamma(2 - \beta)} + \frac{1}{\Gamma(\alpha + 1)\Gamma(2 - \beta)} \right) \\
& \quad + \varepsilon_i e \left( \frac{1}{\Gamma(\alpha - \beta + 1)} + \frac{1}{\Gamma(\alpha)\Gamma(2 - \beta)} + \frac{1}{\Gamma(\alpha + 1)\Gamma(2 - \beta)} \right).
\end{aligned}$$

□

**Theorem 4.1.** Assume that Theorem 3.1 hold, then the fractional differential system (1.1) is Ulam-Hyers stable if the eigenvalues of matrix  $A$  are in the open unit disc. There exists  $|\lambda| < 1$ , for  $\lambda \in \mathbb{C}$  with  $\det(\lambda I - A) = 0$ , where

$$A = \begin{pmatrix} \theta_1(\lambda_1^\alpha + \lambda_1^{\alpha-\beta})l_1 & \theta_2(\lambda_2^\alpha + \lambda_2^{\alpha-\beta})l_2 & \cdots & \theta_2(\lambda_{12}^\alpha + \lambda_{12}^{\alpha-\beta})l_{12} \\ \theta_2(\lambda_1^\alpha + \lambda_1^{\alpha-\beta})l_1 & \theta_1(\lambda_2^\alpha + \lambda_2^{\alpha-\beta})l_2 & \cdots & \theta_2(\lambda_{12}^\alpha + \lambda_{12}^{\alpha-\beta})l_{12} \\ \vdots & \vdots & \ddots & \vdots \\ \theta_2(\lambda_1^\alpha + \lambda_1^{\alpha-\beta})l_1 & \theta_2(\lambda_2^\alpha + \lambda_2^{\alpha-\beta})l_2 & \cdots & \theta_1(\lambda_{12}^\alpha + \lambda_{12}^{\alpha-\beta})l_{12} \end{pmatrix}.$$

**Proof.** Let  $u = (u_1, u_2, \dots, u_{12}) \in X^k$ ,  $k = 1, 2, \dots, 12$ , be the solution of the inequality given by

$$|{}^H D_{1+}^\alpha u_i(t) - \lambda_i^\alpha f_i(t, u_i(t), {}^H D_{1+}^\beta u_i(t))| \leq \varepsilon_i, \quad t \in [1, e],$$

and  $\bar{u} = (\bar{u}_1, \bar{u}_2, \dots, \bar{u}_{12}) \in X^k$  be the solution of the following system

$$\begin{cases} {}^H D_{1+}^\alpha \bar{u}_i(t) = \lambda_i^\alpha f_i(s, \bar{u}_i(s), {}^H D_{1+}^\beta \bar{u}_i(s)), & t \in [1, e], \\ \bar{u}_i(1) = 0, & i = 1, 2, \dots, 12, \\ \bar{u}_i(e) = \bar{u}_j(e), & i, j = 1, 2, \dots, 12, \quad i \neq j, \\ \sum_{i=1}^k \lambda_i^{-1} \bar{u}_i'(e) = 0, & i = 1, 2, \dots, 12. \end{cases} \quad (4.3)$$

By Lemma 2.3, the solution of (4.3) can be given in the following form

$$\begin{aligned}
& \bar{u}_i(t) \\
& = \frac{\lambda_i^\alpha}{\Gamma(\alpha)} \int_1^t \left( \log \frac{t}{s} \right)^{\alpha-1} g_i(s, \bar{u}_i(s), {}^H D_{1+}^\beta \bar{u}_i(s)) ds \\
& \quad - \frac{\log t}{\Gamma(\alpha - 1)} \sum_{j=1}^k \left( \frac{\lambda_j^{-1}}{\sum_{j=1}^k \lambda_j^{-1}} \right) \lambda_j^\alpha \int_1^e \left( \log \frac{e}{s} \right)^{\alpha-2} g_j(s, \bar{u}_j(s), {}^H D_{1+}^\beta \bar{u}_j(s)) ds
\end{aligned}$$

$$\begin{aligned}
& + \frac{\log t}{\Gamma(\alpha)} \sum_{\substack{j=1 \\ j \neq i}}^k \left( \frac{\lambda_j^{-1}}{\sum_{j=1}^k \lambda_j^{-1}} \right) \lambda_j^\alpha \int_1^e \left( \log \frac{e}{s} \right)^{\alpha-1} g_j(s, \bar{u}_j(s), {}^H D_{1+}^\beta \bar{u}_j(s)) ds \\
& - \frac{\log t}{\Gamma(\alpha)} \sum_{\substack{j=1 \\ j \neq i}}^k \left( \frac{\lambda_j^{-1}}{\sum_{j=1}^k \lambda_j^{-1}} \right) \lambda_i^\alpha \int_1^e \left( \log \frac{e}{s} \right)^{\alpha-1} g_i(s, \bar{u}_i(s), {}^H D_{1+}^\beta \bar{u}_i(s)) ds.
\end{aligned}$$

Now, by Lemma 4.1, for  $t \in [1, e]$ , we can get

$$\begin{aligned}
& |u_i(t) - \bar{u}_i(t)| \\
& \leq |u_i(t) - u_i^*(t)| + |u_i^*(t) - \bar{u}_i(t)| \\
& \leq \varepsilon_i e \left( \frac{1}{\Gamma(\alpha)} + \frac{2}{\Gamma(\alpha+1)} \right) + \varepsilon_j e \sum_{\substack{j=1 \\ j \neq i}}^k \left( \frac{1}{\Gamma(\alpha)} + \frac{1}{\Gamma(\alpha+1)} \right) \\
& + e \left( \frac{1}{\Gamma(\alpha)} + \frac{2}{\Gamma(\alpha+1)} \right) (\lambda_i^\alpha + \lambda_i^{\alpha-\beta}) l_i(t) \left( \|u_i - \bar{u}_i\| + \left\| {}^H D_{1+}^\beta u_i - {}^H D_{1+}^\beta \bar{u}_i \right\| \right) \\
& + e \left( \frac{1}{\Gamma(\alpha)} + \frac{1}{\Gamma(\alpha+1)} \right) \sum_{\substack{j=1 \\ j \neq i}}^k (\lambda_j^\alpha + \lambda_j^{\alpha-\beta}) l_j(t) \left( \|u_j - \bar{u}_j\| + \left\| {}^H D_{1+}^\beta u_j - {}^H D_{1+}^\beta \bar{u}_j \right\| \right)
\end{aligned}$$

and

$$\begin{aligned}
& \left| {}^H D_{1+}^\beta u_i(t) - {}^H D_{1+}^\beta \bar{u}_i(t) \right| \\
& \leq \left| {}^H D_{1+}^\beta u_i(t) - {}^H D_{1+}^\beta u_i^*(t) \right| + \left| {}^H D_{1+}^\beta u_i^*(t) - {}^H D_{1+}^\beta \bar{u}_i(t) \right| \\
& \leq \varepsilon_j e \sum_{\substack{j=1 \\ j \neq i}}^k \left( \frac{1}{\Gamma(\alpha)\Gamma(2-\beta)} + \frac{1}{\Gamma(\alpha+1)\Gamma(2-\beta)} \right) \\
& + \varepsilon_i e \left( \frac{1}{\Gamma(\alpha-\beta+1)} + \frac{1}{\Gamma(\alpha)\Gamma(2-\beta)} + \frac{1}{\Gamma(\alpha+1)\Gamma(2-\beta)} \right) \\
& + e \left( \frac{1}{\Gamma(\alpha-\beta+1)} + \frac{1}{\Gamma(\alpha)\Gamma(2-\beta)} + \frac{1}{\Gamma(\alpha+1)\Gamma(2-\beta)} \right) \\
& \times (\lambda_i^\alpha + \lambda_i^{\alpha-\beta}) l_i(t) \left( \|u_i - \bar{u}_i\| + \left\| {}^H D_{1+}^\beta u_i - {}^H D_{1+}^\beta \bar{u}_i \right\| \right) \\
& + e \left( \frac{1}{\Gamma(\alpha)\Gamma(2-\beta)} + \frac{1}{\Gamma(\alpha+1)\Gamma(2-\beta)} \right) \sum_{\substack{j=1 \\ j \neq i}}^k (\lambda_j^\alpha + \lambda_j^{\alpha-\beta}) \\
& \times l_j(t) \left( \|u_j - \bar{u}_j\| + \left\| {}^H D_{1+}^\beta u_j - {}^H D_{1+}^\beta \bar{u}_j \right\| \right).
\end{aligned}$$

Hence, we have

$$\begin{aligned}
& \|u_i - \bar{u}_i\|_X \\
& = \|u_i - \bar{u}_i\| + \left\| {}^H D_{1+}^\beta u_i(t) - {}^H D_{1+}^\beta \bar{u}_i(t) \right\| \\
& \leq e \left( \frac{\alpha+2}{\Gamma(\alpha+1)} + \frac{1}{\Gamma(\alpha-\beta+1)} + \frac{\alpha+1}{\Gamma(\alpha+1)\Gamma(2-\beta)} \right) \varepsilon_i
\end{aligned}$$

$$\begin{aligned}
& + e \sum_{\substack{j=1 \\ j \neq i}}^k \left( \frac{\alpha+1}{\Gamma(\alpha+1)} + \frac{\alpha+1}{\Gamma(\alpha+1)\Gamma(2-\beta)} \right) \varepsilon_j \\
& + e \left( \frac{\alpha+2}{\Gamma(\alpha+1)} + \frac{1}{\Gamma(\alpha-\beta+1)} + \frac{\alpha+1}{\Gamma(\alpha+1)\Gamma(2-\beta)} \right) \\
& \times (\lambda_i^\alpha + \lambda_i^{\alpha-\beta}) l_i(t) (\|u_i - \bar{u}_i\|_X \\
& + e \sum_{\substack{j=1 \\ j \neq i}}^k \left( \frac{\alpha+1}{\Gamma(\alpha+1)} + \frac{\alpha+1}{\Gamma(\alpha+1)\Gamma(2-\beta)} \right) \lambda_j^\alpha + \lambda_j^{\alpha-\beta}) l_j(t) \|u_j - \bar{u}_j\|_X \\
& = \theta_1 \varepsilon_i + \sum_{\substack{j=1 \\ j \neq i}}^k \theta_2 \varepsilon_j + \theta_1 (\lambda_i^\alpha + \lambda_i^{\alpha-\beta}) l_i(t) \|u_i - \bar{u}_i\|_X \\
& + \sum_{\substack{j=1 \\ j \neq i}}^k \theta_2 (\lambda_j^\alpha + \lambda_j^{\alpha-\beta}) l_j(t) \|u_j - \bar{u}_j\|_X,
\end{aligned}$$

where

$$\begin{aligned}
\theta_1 &= e \left( \frac{\alpha+2}{\Gamma(\alpha+1)} + \frac{1}{\Gamma(\alpha-\beta+1)} + \frac{\alpha+1}{\Gamma(\alpha+1)\Gamma(2-\beta)} \right), \\
\theta_2 &= e \left( \frac{\alpha+1}{\Gamma(\alpha+1)} + \frac{\alpha+1}{\Gamma(\alpha+1)\Gamma(2-\beta)} \right).
\end{aligned}$$

Then we have

$$\begin{aligned}
& (\|u_1 - \bar{u}_1\|_X, \|u_2 - \bar{u}_2\|_X, \dots, \|u_{12} - \bar{u}_{12}\|_X)^T \\
& \leq G(\varepsilon_1, \varepsilon_2, \dots, \varepsilon_{12})^T + A(\|u_1 - \bar{u}_1\|_X, \|u_2 - \bar{u}_2\|_X, \dots, \|u_{12} - \bar{u}_{12}\|_X)^T,
\end{aligned}$$

where

$$G_{12 \times 12} = \begin{pmatrix} \theta_1 & \theta_2 & \cdots & \theta_2 \\ \theta_2 & \theta_1 & \cdots & \theta_2 \\ \vdots & \vdots & \ddots & \vdots \\ \theta_2 & \theta_2 & \cdots & \theta_1 \end{pmatrix}.$$

Then, we can get

$$(\|u_1 - \bar{u}_1\|_X, \|u_2 - \bar{u}_2\|_X, \dots, \|u_{12} - \bar{u}_{12}\|_X)^T \leq (I - A)^{-1} G(\varepsilon_1, \varepsilon_2, \dots, \varepsilon_{12})^T.$$

Let

$$H = (I - A)^{-1} G = \begin{pmatrix} h_{1,1} & h_{1,2} & \cdots & h_{1,12} \\ h_{2,1} & h_{2,2} & \cdots & h_{2,12} \\ \vdots & \vdots & \ddots & \vdots \\ h_{12,1} & h_{12,2} & \cdots & h_{12,12} \end{pmatrix}.$$

Obviously,  $h_{i,j} > 0$ ,  $i, j = 1, 2, \dots, 12$ . Set  $\varepsilon = \max\{\varepsilon_1, \varepsilon_2, \dots, \varepsilon_{12}\}$ , then we can get

$$\|u - \bar{u}\|_{X^k} \leq \left( \sum_{j=1}^k \sum_{i=1}^k h_{i,j} \right) \varepsilon. \quad (4.4)$$

Thus, we have derived that system (1.1) is Ulam-Hyers stable.  $\square$

**Remark 4.2.** Making  $\psi_{f_1, f_2, \dots, f_k}(\varepsilon)$  in (4.4). We have  $\psi_{f_1, f_2, \dots, f_k}(0) = 0$ . Then by Definition 4.2, we deduce that the fractional differential system (1.1) is generalized Ulam-Hyers stable.

## 5. Example

The benzene graph we studied in the system (1.1) can be extended to other types of graphs. For example, star graphs and chord bipartite graphs provide a theoretical basis for physics, computer networks and other fields. Here we only discuss the fractional differential system on the star graphs ( $i = 1, 2, 3$ ). We discuss the solution of a fractional differential equation on a formic acid graph and the approximate graphs of solutions are presented by using iterative methods and numerical simulation.

**Example 5.1.** Consider the following questions

$$\begin{cases} {}^H D_{1+}^{\frac{5}{2}} u_1(t) = \left(\frac{1}{4}\right)^{\frac{5}{2}} \left( \frac{t}{(t+7)^4} \left( \sin(u_1(t)) + \frac{|{}^H D_{1+}^{\frac{3}{2}} u_1(t)|}{1 + |{}^H D_{1+}^{\frac{3}{2}} u_1(t)|} \right) \right), \\ {}^H D_{1+}^{\frac{5}{2}} u_2(t) = \left(\frac{1}{2}\right)^{\frac{5}{2}} \left( \frac{t}{(t^3+3)^5} \left( \sin(u_2(t)) + \frac{|{}^H D_{1+}^{\frac{3}{2}} u_2(t)|}{1 + |{}^H D_{1+}^{\frac{3}{2}} u_2(t)|} \right) \right), \\ {}^H D_{1+}^{\frac{5}{2}} u_3(t) = \left(\frac{3}{4}\right)^{\frac{5}{2}} \left( \frac{3t^2}{1000} |\arcsin(u_3(t))| + \frac{3t^2 |{}^H D_{1+}^{\frac{3}{2}} u_3(t)|}{1000(1 + |{}^H D_{1+}^{\frac{3}{2}} u_3(t)|)} \right), \\ u_1(1) = u_2(1) = u_3(1) = 0, \\ u_1(e) = u_2(e) = u_3(e), \\ \left(\frac{1}{4}\right)^{-1} u'_1(e) + \left(\frac{1}{2}\right)^{-1} u'_2(e) + \left(\frac{3}{4}\right)^{-1} u'_3(e) = 0, \end{cases} \quad (5.1)$$

We obtain

$$\alpha = \frac{5}{2}, \quad \beta = \frac{3}{2}, \quad k = 3, \quad \lambda_1 = \frac{1}{4}, \quad \lambda_2 = \frac{1}{2}, \quad \lambda_3 = \frac{3}{4}.$$

Coordinate systems with  $u_1$ ,  $u_2$ ,  $u_3$  are established respectively on the formic acid graph with 3 edges (Figure 4).

For  $t \in [1, e]$ ,

$$\begin{aligned} g_1(t, u_1(t), {}^H D_{1+}^{\beta} u_1(t)) &= \frac{1}{(t+7)^4} \left( \sin(u_1(t)) + \frac{|{}^H D_{1+}^{\frac{3}{2}} u_1(t)|}{1 + |{}^H D_{1+}^{\frac{3}{2}} u_1(t)|} \right), \\ g_2(t, u_2(t), {}^H D_{1+}^{\beta} u_2(t)) &= \frac{1}{(t^3+3)^5} \left( \sin(u_2(t)) + \frac{|{}^H D_{1+}^{\frac{3}{2}} u_2(t)|}{1 + |{}^H D_{1+}^{\frac{3}{2}} u_2(t)|} \right), \end{aligned}$$



Figure 3. A sketch of  $C_2H_2O$ .

Figure 4. Formic acid graph with labeled vertices.

$$g_3(t, u_3(t), {}^H D_{1+}^\beta u_3(t)) = \frac{3t}{1000} |\arcsin(u_3(t))| + \frac{3t |{}^H D_{1+}^{\frac{3}{2}} u_3(t)|}{1000(1 + |{}^H D_{1+}^{\frac{3}{2}} u_3(t)|)}.$$

For any  $x, y, x_1, y_1$ , it is clear that

$$\begin{aligned} g_1(t, x, y) - g_1(t, x_1, y_1) &\leq \frac{1}{(t+7)^4} (|x - x_1| + |y - y_1|), \\ g_2(t, x, y) - g_2(t, x_2, y_2) &\leq \frac{1}{(t^3+3)^5} (|x - x_2| + |y - y_2|), \\ g_3(t, x, y) - g_3(t, x_3, y_3) &\leq \frac{3t}{1000} (|x - x_3| + |y - y_3|). \end{aligned}$$

So we get

$$l_1 = \sup_{t \in [1, e]} |l_1(t)| = \frac{1}{4096}, l_2 = \sup_{t \in [1, e]} |l_2(t)| = \frac{1}{1024}, l_3 = \sup_{t \in [1, e]} |l_3(t)| = \frac{3e}{1000},$$

$$B_1 = 22.3235, B_2 = 23.7510, B_3 = 25.7324,$$

and

$$(B_1 + B_2 + B_3)(l_1 + l_2 + l_3) = 0.6735 < 1.$$

Therefore, by Theorem 3.1 system (5.1) has a unique solution on  $[1, e]$ .

Meanwhile,

$$\begin{aligned} \theta_1 &= 8.0140, \quad \theta_2 = 4.4778, \\ A &= \begin{pmatrix} 5.5028e-04 & 2.9594e-03 & 0.0452 \\ 3.0747e-04 & 5.2966e-03 & 0.0452 \\ 3.0747e-04 & 2.9594e-03 & 0.0809 \end{pmatrix}. \end{aligned}$$

Let

$$\det(\lambda I - A) = (\lambda - 0.0824)(\lambda - 0.0003)(\lambda - 0.0036) = 0,$$

so we have

$$\lambda_1 = 0.0824 < 1, \quad \lambda_2 = 0.0003 < 1, \quad \lambda_3 = 0.0036 < 1.$$

It follows from Theorem 4.1 that system (5.1) is Ulam-Hyers stable, and by Remark 4.2, it will be generalized Ulam-Hyers stable.


Ultimately, the simulate iterative process curve and approximate solution to the fractional differential system (5.1) are carried out by using the iterative method and numerical simulation. Let  $u_{i,0} = 0$ , the iteration sequence is as follows,

$$u_{1,n+1}(t)$$




$$\begin{aligned}
&= \frac{(\frac{1}{4})^{\frac{5}{2}}}{\Gamma(\frac{5}{2})} \int_1^t \left( \log \frac{t}{s} \right)^{\frac{3}{2}} \frac{1}{(t+7)^4} \left( \sin(u_{1,n}(t)) + \frac{|{}^H D_{1+}^{\frac{3}{2}} u_{1,n}(t)|}{1 + |{}^H D_{1+}^{\frac{3}{2}} u_{1,n}(t)|} \right) ds \\
&\quad + \frac{(\frac{1}{4})^{\frac{5}{2}} (\frac{1}{2})^{-1} (\log t)}{\left( (\frac{1}{4})^{-1} + (\frac{1}{2})^{-1} + (\frac{3}{4})^{-1} \right) \Gamma(\frac{5}{2})} \int_1^e (1 - \log s)^{\frac{3}{2}} \frac{1}{(t^3 + 3)^5} \left( \sin |u_{2,n}(t)| \right. \\
&\quad \left. + \frac{|{}^H D_{1+}^{\frac{3}{2}} u_{2,n}(t)|}{1 + |{}^H D_{1+}^{\frac{3}{2}} u_{2,n}(t)|} \right) ds + \frac{(\frac{1}{4})^{\frac{5}{2}} (\frac{1}{2})^{-1} (\log t)}{\left( (\frac{1}{4})^{-1} + (\frac{1}{2})^{-1} + (\frac{3}{4})^{-1} \right) \Gamma(\frac{5}{2})} \int_1^e (1 - \log s)^{\frac{3}{2}} \\
&\quad \times \left( 0.003t |\arcsin(u_{3,n}(t))| + \frac{3t |{}^H D_{1+}^{\frac{3}{2}} u_{3,n}(t)|}{1000 + 1000 |{}^H D_{1+}^{\frac{3}{2}} u_{3,n}(t)|} \right) ds \\
&\quad - \frac{(\frac{1}{2})^{-1} (\frac{1}{4})^{\frac{5}{2}} (\log t)}{\left( (\frac{1}{4})^{-1} + (\frac{1}{2})^{-1} + (\frac{3}{4})^{-1} \right) \Gamma(\frac{5}{2})} \int_1^e (1 - \log s)^{\frac{3}{2}} \frac{1}{(t+7)^4} \left( \sin(u_{1,n}(t)) \right. \\
&\quad \left. + \frac{|{}^H D_{1+}^{\frac{3}{2}} u_{1,n}(t)|}{1 + |{}^H D_{1+}^{\frac{3}{2}} u_{1,n}(t)|} \right) ds - \frac{(\frac{1}{4})^{\frac{5}{2}} (\frac{3}{4})^{-1} (\log t)}{\left( (\frac{1}{4})^{-1} + (\frac{1}{2})^{-1} + (\frac{3}{4})^{-1} \right) \Gamma(\frac{5}{2})} \int_1^e (1 - \log s)^{\frac{3}{2}} \\
&\quad \times \frac{1}{(t+7)^4} \left( \sin(u_{1,n}(t)) + \frac{|{}^H D_{1+}^{\frac{1}{2}} u_{1,n}(t)|}{1 + |{}^H D_{1+}^{\frac{1}{2}} u_{1,n}(t)|} \right) ds - \frac{(\frac{1}{4})^{\frac{5}{2}} (\frac{1}{4})^{-1} (\log t)}{\left( (\frac{1}{4})^{-1} + (\frac{1}{2})^{-1} + (\frac{3}{4})^{-1} \right)} \\
&\quad \times \frac{1}{\Gamma(\frac{3}{2})} \int_1^e (1 - \log s)^{\frac{1}{2}} \frac{1}{(t+7)^4} \left( \sin(u_{1,n}(t)) + \frac{|{}^H D_{1+}^{\frac{1}{2}} u_{1,n}(t)|}{1 + |{}^H D_{1+}^{\frac{1}{2}} u_{1,n}(t)|} \right) ds \\
&\quad - \frac{(\frac{1}{2})^{\frac{5}{2}} (\frac{1}{2})^{-1} (\log t)}{\left( (\frac{1}{4})^{-1} + (\frac{1}{2})^{-1} + (\frac{3}{4})^{-1} \right) \Gamma(\frac{3}{2})} \int_1^e (1 - \log s)^{\frac{1}{2}} \frac{1}{(t^3 + 3)^5} \left( \sin |u_{2,n}(t)| \right. \\
&\quad \left. + \frac{|D_{1+}^{\frac{3}{2}} u_{2,n}(t)|}{1 + |D_{1+}^{\frac{3}{2}} u_{2,n}(t)|} \right) ds - \frac{(\frac{3}{4})^{\frac{5}{2}} (\frac{3}{4})^{-1} (\log t)}{\left( (\frac{1}{4})^{-1} + (\frac{1}{2})^{-1} + (\frac{3}{4})^{-1} \right) \Gamma(\frac{3}{2})} \int_1^e (1 - \log s)^{\frac{1}{2}} \\
&\quad \times \left( 1 + 0.003t |\arcsin(u_{3,n}(t))| + \frac{3t |{}^H D_{1+}^{\frac{3}{2}} u_{3,n}(t)|}{1000 + 1000 |{}^H D_{1+}^{\frac{3}{2}} u_{3,n}(t)|} \right) ds.
\end{aligned}$$


The iterative sequence of  $u_{2,n+1}$  and  $u_{3,n+1}$  is similar to  $u_{1,n+1}$ . After several iterations, the approximate solution of fractional differential system (5.1) can be obtained by using the numerical simulation. Figure 5 is the approximate graph of the solution of  $\overrightarrow{u_1 u_0}$  after iterations. Figure 6 is the approximate graph of the solution of  $\overrightarrow{u_2 u_0}$  after iterations. Figure 7 is the approximate graph of the solution of  $\overrightarrow{u_3 u_0}$  after iterations.



**Figure 5.** Approximate solution of  $u_1$ .



**Figure 6.** Approximate solution of  $u_2$ .



**Figure 7.** Approximate solution of  $u_3$ .

## Acknowledgements

We declare that all authors conceive of the study and carry out the proof. All authors of this paper read and approve the final manuscript.

## References

- [1] A. Z. Abdian, A. Behmaram and G. H. Fath-Tabar, *Graphs determined by signless Laplacian Spectra*, Akce. Int. J. Graphs. Co., 2020, 17(1), 45–50.
- [2] B. Ahmad, M. Alghanmi, A. Alsaedi, et al., *Existence and uniqueness results for a nonlinear coupled system involving Caputo fractional derivatives with a new kind of coupled boundary conditions*, Appl. Math. Lett., 2021, 116, 107018.
- [3] B. Ahmad, A. Alsaedi, S. K. Ntouyas, et al., *Hadamard-Type Fractional Differential Equations, Inclusions and Inequalities*, Springer, Switzerland, 2017.
- [4] Y. Cui, W. Ma, et. al, *New uniqueness results for boundary value problem of fractional differential equation*, Nonlinear. Anal-Model., 2018, 23(1), 31–39.
- [5] L. Debnath, *Fractional integrals and fractional differential equations in fluid mechanics*, Frac. Calc. Appl. Anal., 2003, 6, 119155.
- [6] M. Faieghi, S. Kuntanapreeda, H. Delavari, et al., *LMI based stabilization of a class of fractional order chaotic systems*, Nonlinear. Dynam., 2013, 72(1), 301–309.
- [7] D. G. Gordeziani, M. Kupreishvli, H. V. Meladze, et al., *On the solution of boundary value problem for differential equations given in graphs*, Appl. Math. Lett., 2008, 13, 80–91.
- [8] J. R. Graef, L. Kong and M. Wang, *Existence and uniqueness of solutions for a fractional boundary value problem on a graph*, Fract. Calc. Appl. Anal., 2014, 17(2), 499–510.
- [9] J. Klafter, S. C. Lim and R. Metzler, *Fractional Dynamics in Physics*, World Scientific, Singapore, 2011.
- [10] V. Lakshmikantham and A. S. Vatsala, *Basic theory of fractional differential equations*, Nonlinear. Anal., 2008, 69, 267–782.
- [11] A. Mahdy, M. S. Mohamed, K. A. Gepreel, et al., *Dynamical characteristics and signal flow graph of nonlinear fractional smoking mathematical model*, Chaos Soliton Fract., 2020, 141, 110308.
- [12] V. Mehendiratta, M. Mehra and G. Leugering, *Existence and uniqueness results for a nonlinear Caputo fractional boundary value problem on a star graphs*, J. Math. Anal. Appl., 2019, 477(2), 1243–1264.
- [13] R. Metzler and J. Klafter, *Boundary value problems for fractional diffusion equations*, Phys A., 2000, 278(1–2), 107–125.
- [14] J. Nan, W. Hu, Y. Su, et al., *Existence and stability of solutions for a coupled hadamard type sequence fractional differential system on glucose graphs*, J. Appl. Anal. Comput., 2024, 14(2), 911–946.
- [15] V. Pivovarchik, *Inverse problem for the Sturm-Liouville equation on a star-shaped graph*, Math. Nachr., 2007, 280(13), 1595–1619.

- [16] S. Z. Rida, A. M. A. El-Sayed and A. A. M. Arafa, *Effect of bacterial memory dependent growth by using fractional derivatives reaction diffusion chemotactic model*, J. Stat. Phys., 2010, 140(4), 797–811.
- [17] N. A. Sheikh, M. Jamil, D. Ching, et al., *A generalized model for quantitative analysis of sediments loss: A Caputo time fractional model*, J. King. Saud. Univ. Sci., 2020, 33(1), 101179.
- [18] A. Sun, Y. Su and Q. Yuan, *Existence of solutions to fractional differential equations with fractional order derivative terms*, J. Appl. Anal. Comput., 2021, 11(1), 486–520.
- [19] W. Sun, Y. Su, et al., *Existence and simulation of positive solutions for  $m$ -point fractional differential equations with derivative terms*, Open. Math., 2021, 19(1), 1820–1846.
- [20] W. Sun, Y. Su and X. Han, *Existence of solutions for a coupled system of Caputo-Hadamard fractional differential equations with  $p$ -laplacian operator*, J. Appl. Anal. Comput., 2022, 12(5), 1885–1900.
- [21] J. Tariboon, S. K. Ntouyas, S. Asawasamrit, et al., *Positive solutions for Hadamard differential systems with fractional integral conditions on an unbounded domain*, Open. Math., 2017, 15(1), 645–666.
- [22] D. Tripathil, S. Pandey and S. Das, *Peristaltic flow of viscoelastic fluid with fractional maxwell model through a channel*, Appl. Math. Comput., 2010, 215, 3645–3654.
- [23] A. Turab and W. Sintunavarat, *The novel existence results of solutions for a nonlinear fractional boundary value problem on the ethane graph*, Alex. Eng. J., 2021, 60(6), 5365–5374.
- [24] J. R. Wang, Y. Zhou and M. Medve, *Existence and stability of fractional differential equations with Hadamard derivative*, Topol. Method. Non. Anal., 2013, 41(1), 113–133.
- [25] W. Zhang and W. Liu, *Existence and Ulam's type stability results for a class of fractional boundary value problems on a star graphs*, Math. Method. Appl. Sci., 2020, 43, 8568–8594.
- [26] Y. Zhang, Y. Su and Y. Yun, *Existence and stability of solutions for Hadamard type fractional differential systems with  $p$ -Laplacian operators on benzoic acid garphs*, AIMS Math., 2025, 10(4), 7767–7794.

Received December 2024; Accepted April 2025; Available online April 2025.